

Report on Diamond Drilling Pacific North West Capital Corp.

Belford Township, Timmins Area, Ontario
Porcupine Mining Division

NTS 42B-9

October 06, 2005



2 . 3 0 6 5 2

By
Joan Barry, P.Geo.

Table of Contents

INTRODUCTION

PROPERTY LOCATION AND ACCESS

PREVIOUS WORK

CURRENT WORK PROGRAM

PERSONNEL

REFERENCES

Table of Figures

PROPERTY LOCATION MAP

DRILL HOLE PLAN

DRILL HOLE SECTION

Appendix

DIAMOND DRILL HOLE LOG

INTRODUCTION

Pacific North West Capital Corp. carried out a diamond drilling program to test an AEM conductor that was confirmed by ground Max Min EM. Hole WTM_05-01 (E) was drilled to a depth of 210 metres. The hole was started on September 28 and completed on September 30, 2005. The drilling was completed by Benoit Drilling of Val d'Ore, P.Q.

PROPERTY LOCATION AND ACCESS

The property consists of one unpatented minig claim # 3006250 (16 claim units) which Pacific North West Capital Corp. has optioned from Falconbridge Limited.

The property is located approximately 65 km west north-west of Timmins, Ontario. It can be accessed by travelling west from Timmins along Highway 101 for approximately 10 km and then proceeding west and north along the Mallette Lumber Road for approximately 75 km. Various bush roads give additional good access.

PREVIOUS WORK

In November 2004, AeroQuest Limited carried out a helicopter survey on behalf of Falconbridge Limited on the West Timmins Property near Timmins, Ontario. An AeroTEM II time domain helicopter electromagnetic system was used in conjunction with a high-sensitivity cesium vapor magnetometer. This survey covered claim # 3006250.

In September, 2005 Pacific North West Capital Corp. cut a 16 line km Grid 2 on the property over which ground HLEM and Magnetic surveys were conducted to ground truth the airborne anomaly. This work has not yet been compiled.

CURRENT WORK PROGRAM

Pacific North West Capital Corp.

Hole WTM_05-01 (E) was drilled to a depth of 201 metres to test the EM conductor. The hole was started on September 28 and completed on September 30, 2005. The drilling was carried out by Benoit Drilling of Val d'Ore, P.Q.

The hole was collared at 400W, 25 E on Grid 2 to intersect an HLEM conductor flanked by a magnetic high. It was anticipated that the conductor would be intersected at 127 metres down hole. Several zones of graphitic schist up to 2 metres in thickness were encountered in the hole however the likely source of the conductor is 6.05 metre graphitic schist with a zone of up to 5% pyrite and pyrrhotite. Assays are pending.

PERSONNEL

The following persons were involved in the planning, supervision, performance and report writing of this work:

John Londry, P.Geo.

Vice President of Exploration
259 Fielding Road, Unit 3B
Lively, Ontario
P3Y 1L8
Work phone: 705-674-5888

Michel Leblanc, P.Geo

Project Geologist
259 Fielding Road, Unit 3B
Lively, Ontario
P3Y 1L8
Work phone: 705-674-5888

Jennifer Berger

Geologist
259 Fielding Road, Unit 3B
Lively, Ontario
P3Y 1L8
Work phone: 705-674-5888

Leo Levac

Field Technition
259 Fielding Road, Unit 3B
Lively, Ontario
P3Y 1L8
Work phone: 705-674-5888


Joan Barry, P.Geo.


Geologist, Office Manager
259 Fielding Road, Unit 3B
Lively, Ontario
P3Y 1L8
Work phone: 705-674-5888

REFERENCES

AeroQuest Limited, 2004. Report on a Helicopter-Borne AeroTEM II
Electromagnetic and Magnetometer Survey. Aeroquest Jon # 04028, Montcalm
Area - Area of Interest, Timmins Area, Ontario.

Respectfully submitted,


Joan M. Barry, B.Sc., P.Geo.
Geologist



Grid 2

WTM-05-01
Az: 225
Dip: -45
200m

1000W

400W

0W

BLO

P3006250

P3006252

P3006251

0 300 600
metres


Scale: 1:12,500
Projection: Nad 83 Zone 17
October, 2005

Pacific North West Capital Corp.

West Timmins Project
Montcalm Grid 2 and
Diamond Drilling
Belford Township



Watson



Belford

Scale: 1:100,000
Projection: Nad 83 Zone 17
October, 2005

Pacific North West Capital Corp.

West Timmins Project
Montcalm Grid 2 and
Diamond Drilling
Belford Township

WTC
WTM-05-01
P3006250

P3006252

4203045
4203046

4202914
4203047

4207723
4207721

4207722

P3008918

P3008919

P3008917

P3008916

P4206316

P3008915

P4206355

P3006302

P3008924

4207724

P3008925

P4206313

P3006303

P4206359

P3006304

P4206356

P3006305

P3008921

P1169587

P3008922

P3008920

P3008923

P1169588

P1169589

P1169594

P4206309

P1169586

P4206308

P4206350

P4206354

P3008927

P4206351

P4206352

P4206353

P1169582

P1169581

P3008926

P1169590

P3008913

P3008914


P3008911

P3008912

4203045
4203046

WTC
WTM-05-01
P3006250

P3006252



- WCK - WACKE
- GAB - GABBRO
- FP - FELDSPAR PORPHYRY
- ID - INTERMEDIATE DYKE
- FEL - FELSIC
- BIO - BIOTITE
- SIL - SILICEOUS
- GNT - GARNET
- QV - QUARTZ VEIN
- PYRR - PYRRHOTITE
- PY - PYRITE

SYMBOL: PFN
 TSX
 TORONTO STOCK EXCHANGE
 OTCBB: PAWEF

PACIFIC NORTH WEST
 CAPITAL CORP.

West Timmins Project: Montcalm	
Hole Number:	WTM-05-01 (E) Grid 2 L400W-25S
Azimuth:	225
Inclination:	-45
Length:	Prop EOH @ 200m → Actual EOH 201m
Date:	Oct 13, 2005
Scale:	1:1000

APPENDIX

Hole ID: WTM_05-01 (E)
 Property: West Timmins Project (Montcalm)
 Claim No:
 Casing: Left in hole
 Core Size: NQ
 Azimuth: 225
 Inclination: -45
 Length (EOH): 201 meters
 Date Started: Sept. 28, 2005
 Completed: Sept. 30, 2005
 Logged By: J. Berger

Township:
 Grid Number: 2
 Grid East: 400 West
 Grid North: 25 South
 UTM East: 397323 E
 UTM North: 5387113 N
 UTM Elevation:
 Survey Method: Handheld GPS
 Drill Contractor: Benoit Drilling

Surveys/Tests:

Depth	Azimuth	Inclination
39m	223.5	-45.1
90m	224.9	-44.9

Remarks: 108.85-114.35: Felsic/Silicified Sediments, good Au potential
 117.00-123.05: Graphite Schist, locally up to 10% zPyrr + Py (likely the conductor)

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
0.00	25.00	CASING	Overburden				
25.00	28.00	WACKE	Fine grained, dark grey, locally up to 10% garnet porphyroblasts, locally weak actinolite alteration, wk				
			25.10-25.45: 20% garnet porphyroblasts, 20-25% qtz veins/flooding, 5% Pyrite + trace Pyrrhotite blebs/disseminated, some rusty fracture surface	25.00	25.70	0.70	94651
			26.00-26.40: Banded section, 30% carbonate/felsic bands (mm-cm scale banding), pale grey/white color, banding at 52, locally rusty fracture surfaces	25.70	27.10	1.40	94652
			27.00-27.65: 40% Siliceous/felsic bands (3-15cm wide), light grey color, fine grained, banding at 56				
			27.65-28.00: 20% qtz veins/flooding, minor plagioclase, 10% rusty blebs + vugs, 10% Actinolit (patchy), 5-10% garnet porphyroblasts, 3% Pyrite + 1% Pyrrhotite disseminated + blebs	27.10	28.00	0.90	94653
28.00	32.55	BIOTITE GNEISS	Fine/medium grained, 15-20% biotite, moderate foliation at 52, medium grey color, trace disseminated Pyrite, locally weak actinolite alteration, minor chlorite, <5% qtz-plagioclase veinlets/veins (2mm-2cm wide)				
32.55	43.90	GABBRO	Medium grained, moderate actinolite alteration throughout, greenish dark grey, minor qtz-plagioclased veinlets/fracture fill, trace disseminated Pyrite + Pyrrhotite				
			43.10-43.51: 50% qtz-plagioclase veins/flooding				
			43.36-43.51: qtz vein, minor plagioclase, contacts at 39				
			43.80-43.90: Medium grained qtz-plagioclase vein, which forms the contact of the feldspar porphyry unit, minor rust, no visible sulfides, contact at 36				

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
43.90	45.80	FELDSPAR PORPHYRY	Purplish brown color, 20-25% plagioclase phenocrysts (round, 2mm-1cm in size), 15% Biotite, weak foliation at 50, trace disseminate Pyrite, locally minor rust associate with fracture surfaces and in veinlets, 2 faults healed with qtz (each showing about 1cm of displacement) Sharp contact at 45.80				
45.80	52.21	GABBRO	Medium grained, dark greenish grey, weak-moderate actinolite alteration throughout, <5% qtz vein's, locally minor chlorite alteration along fracture surfaces, foliation at 41				
52.21	54.10	INTERMEDIATE VOLCANICS	Fine grained, medium grey, 10% Hornblende phenocrysts, uniform, trace disseminated Pyrite, sharp but irregular contacts				
54.10	56.10	BIOTITE GNEISS	Strong foliation at 40, 30-40% Biotite, 20% felsic bands locally crenulated, rare qtz veinlets, sharp but irregular contacts				
56.10	58.00	GABBRO	Medium grained, massive, dark grey				
58.00	59.65	BIOTITE GNEISS	25% Biotite bands, 10-15% felsic layers/bands (<1cm wide), fine grained, banding at 52, rare garnets, 3cm qtz-plag vein associated with strong chlorite alteration, vein at 32				
59.65	61.10	GRAPHITE SCHIST	20% Biotite bands, <5% garnets in bands, 5% qtz-plagioclase veinlets (<1cm wide, dark grey/black, 5% Pyrrhotite>Pyrite in blebs and veins, sharp contact with the BIO GNEISS at 32, banding at 36	59.65	61.10	1.45	94654
61.10	65.00	GRAPHITE SCHIST	Dark grey/black, fine grained, 10% Biotite banding, 3% Pyrrhotite>Pyrite as fracture fill, veins (0.5cm wide) and blebs; compositional banding at 51	61.10	62.10	1.00	94655
				62.10	63.10	1.00	94656
				63.10	64.10	1.00	94657
				64.10	65.00	0.90	94658
65.00	65.90	FELSIC DYKE	Fine grained, pale grey, 50% of core is broken, 1% Pyrite>Pyrrhotite disseminated + 1% on fracture surfaces, 10% Biotite bands at 50 (brownish color)	65.00	65.90	0.90	94659
65.90	68.85	GRAPHITE SCHIST	As above, banding at 51	65.90	66.90	1.00	94660
			67.90-68.85: 25-30% Biotite bands, 3% Pyrite>Pyrrhotite as blebs + disseminated, banding at 48	66.90	67.90	1.00	94661
				67.90	68.85	0.95	94662
68.85	73.88	GNEISS	Fine grained, compositional banding, predominantly felsic bands (60-70%), 10% Biotite bands, banding at 55, 1-2% Pyrite+Pyrrhotite on fracture surfaces, disseminated and as veinlets (<0.5cm				
73.88	74.20	INTERMEDIATE DYKE	Uniform, medium grey, fine grained, sharp contacts at 60				

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
74.20	78.80	GRAPHITE SCHIST	Dark grey/black, fine grained, 10% Biotite bands, rare qtz-plagioclase vein's (<0.5cm wide), locally vuggy (<5%); 2-3% Pyrrhotite>Pyrite as veinlets, diss, blebs + on fracture surfaces, banding at 49, 73.10-73.50: Felsic Dyke, fine grained, light grey, sharp contacts at 60 77.10-78.20: 5% Pyrrhotite>Pyrite	74.20	75.10	0.90	94663
				75.10	76.10	1.00	94664
				76.10	77.10	1.00	94665
				77.10	78.20	1.10	94666
				78.20	78.80	0.60	94667
78.80	81.40	GABBRO/ AMPHIBOLITE	Coarse Grained, moderate/strong Actinolite alteration, trace disseminated Pyrite 79.90-80.30: Banded section, 20% felsic bands 80.60-81.40: 3% Pyrrhotite>Pyrite as blebs up to 1cm in size, on fracture surfaces and associated with the contacts of a 1cm wide qtz vein (qtz vein contacts at 16)	80.60	81.40	0.80	94668
81.40	84.00	BIOTITE GNEISS	30% Biotite bands, pale grey to brown, banded, <5% qtz veinlets, <1% Pyrite along fracture surfaces, bands at 5				
84.00	84.45	GABBRO/ AMPHIBOLITE	Massive-weakly foliated, moderate actinolite alteration, dark greenish grey, coarse grained, sharp contacts at 57				
84.45	85.65	BIOTITE GNEISS	As Above				
85.65	86.00	FELSIC DYKE	Pale grey, fine grained, uniform, minor Pyrite at contacts, sharp contacts at 60				
86.00	87.65	BIOTITE GNEISS	20-25% Biotite bands, pale grey-brownish color, banded at 60, 5% qtz veinlets (<2cm wide), <1% Pyrite along fracture surfaces				
87.65	97.15	GABBRO	Medium grained, massive-weak foliation, moderate actinolite alteration, <5% qtz-plagioclase veinlets (<0.5cm wide)				
			91.55-97.15: Moderately foliated gabbro, fine/medium grained, 10% biotite bands, 5-10% qtz and qtz-plagioclase veinlets				
97.15	98.65	GARNET BIOTITE GNEISS	Banded, 25% biotite bands, 5-10% garnet porphyroblasts (<1cm in size), 10% graphitic bands(dark black), 10% qtz-plagioclase veins (<1cm wide), BANDS AT 56 99.55-99.75: Qtz flooded section				
98.65	101.45	GABBRO	Medium-Coarse grained, weak-moderate actinolite alteration, moderate foliation at 55, 10-15% biotite bands				
101.45	102.80	GRAPHITE SCHIST	40% graphitic bands (dark black), 10% biotite bands, rare qtz veinlets; 1% Pyrite + Pyrrhotite as veinlets (2mm wide) and on fracture surface, locally weak chlorite alteration, bands at 64	101.45	102.80	1.35	94669

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
102.80	103.90	GABBRO	As Above, foliation at 60				
103.90	104.15	GRAPHITE SCHIST	Dark grey + black bands, minor qtz veinlets; 3-5% Pyrrhotite > Pyrite disseminated, blebs and veinlets; sharp contacts at 55, bands at 60				
104.15	104.50	INTERMEDIATE DYKE	Fine grained, sharp contacts at 60, massive, uniform, light grey				
104.50	104.70	GRAPHITE SCHIST	As Aove, banding at 61				
104.70	105.90	INTERMEDIATE DYKE	Fine grained, locally foliated at 52, 10-15% Hornblende phenocrysts elongated, sharp contacts at 52, up to 2% Pyrite and Pyrrhotite disseminated				
105.90	107.70	GRAPHITE SCHIST	Dark grey-black, 10% biotite, 5% Pyrrhotite + Pyrite disseminated, blebs, veins, and on fracture surfaces, 10% qtz-plagioclase veinlets (<0.5cm wide), bands at 56	105.90	106.80	0.90	94670
				106.80	107.70	0.90	94671
107.70	108.85	CONTACT	Gradational contact, 60% felsic layers ad 30% black graphitic layers, 10% qtz-plagioclase veins (<2cm wide), 1-1cm Tourmaline vein adjacent to 2 qtz-plagioclase veins, strongly crenulated, weak chlorite alteration of the more felsic layers, 2-3% Pyrrhotite as blebs and disseminated				
108.85	114.35	SILICEOUS SEDIMENTS	Fine grained, banded at 55, very siliceous unit, pale greenish grey color, moderate chlorite alteration, 5% qtz-plagioclase veins (0.5-10cm wide), inor actinolite alteration, 3% Pyrrhotite>Pyrite in veins (<0.5 cm wide) and disseminated	108.85	109.90	1.05	94673
			109.10-109.20: Qtz-feldspar vein, fine grained, minor k-feldspar, sharp contacts at 30	109.90	110.90	1.00	64674
			111.75-112.05: Graphite schist, crenulated, 2% Pyrrhotite and minor Pyrite, 80% graphite bands/layers, minor qtz and actinolite	110.90	111.75	0.85	94675
				111.75	112.80	1.05	94676
				112.80	113.70	0.90	94677
				113.70	114.35	0.65	94678
114.35	115.60	GRAPHITE SCHIST	Banded, 40% graphitic bands-dark black, <5% qtz and qtz-plagioclase veinlets, 5% strongly actinolite altered bands, 3% Pyrrhotite and Pyrite mainly as veins (<0.5cm wide), bands at 60	114.35	115.60	1.25	94679
115.60	117.00	SILICEOUS SEDIMENTS	As above, bands at 57	115.60	116.30	0.70	94680
				116.30	117.00	0.70	94681
117.00	123.05	GRAPHITE SCHIST	Dark grey-black, banded at 60, 2 phases of qtz and qtz-plagioclase veins (<1cm wide) cross-cutting one another, 10% biotite, sulphides vary from section to section				

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
			117.00-119.80: 5% Pyrite and Pyrrhotite mainly in veins (about 0.5cm wide), 1-1cm pyrrhotite pod, minor sulphides disseminated and in blebs	117.00	118.00	1.00	94682
				118.00	119.00	1.00	94683
				119.00	119.80	0.80	94684
			119.80-120.60: Up to 10% Pyrrhotite > Pyrite as veins (up to 1cm wide), pods/blebs and	119.80	120.60	0.80	94685
			120.60-121.80: 3% Pyrrhotite > pyrite in blebs and veins (about 0.5cm wide), only 30% graphitic bands	120.60	121.80	1.20	94686
			121.80-122.50: 1-2% Pyrite + Pyrrhotite, blebs + disseminated, only 10% graphitic bands	121.80	122.50	0.70	94687
			122.50-123.05: Up to 10% Pyrrhotite > Pyrite as veins (up to 1.5cm wide), blebs and (minor)	122.50	123.05	0.55	94688
123.05	131.15	SILICEOUS SEDIMENTS	Banded at 60, fine grained, pale grey to pale greenish grey, locally weak chlorite alteration, rare qtz-plagioclase veins (<1cm wide); 3% Pyrrhotite + Pyrite as veins (0.5cm wide), blebs and disseminated	123.05	124.00	0.95	94689
				124.00	125.00	1.00	94690
				125.00	126.00	1.00	94691
				126.00	127.00	1.00	94692
				127.00	128.00	1.00	94693
				128.00	129.00	1.00	94694
			129.65-129.85: Graphite rich section, 2 0.5cm qtz-plagioclase veins	129.00	129.65	0.65	94695
			129.85-130.20: 10% garnet porphyroblasts (<0.5cm in size)	129.65	130.20	0.55	94696
				130.20	131.15	0.95	94697
131.15	132.95	GRAPHITE SCHIST	30% Graphitic layers/bands, black to dark grey, bands at 60, 10% qtz-plagioclase veins (2mm wide), 3% Pyrite + Pyrrhotite in blebs and veins (about 0.5cm wide), sharp contacts with the surrounding units	131.15	132.00	0.85	94698
				132.00	132.95	0.95	94699
132.95	134.90	SILICEOUS SEDIMENTS	Pale grey, uniform, no distinctive banding, fine grained, rare garnet porphyroblasts (<0.5cm in size)				
134.90	144.60	GARNET BIOTITE GNEISS	Fine grained, 15% garnet porphyroblasts (0.5cm in size), 10% hornblende laths (<1m in size), moderate actinolite alteration, brown to grey color, no distinctive preferred orientation, rare qtz veins (<1cm wide), trace Pyrite, 10% biotite rich bands				
144.60	145.10	GRAPHITE SCHIST	20% graphitic bands (<cm wide), 30% qtz veins/flooding, fine grained, pale greenish grey layers that have been weakly chloritized, 1% pyrrhotite specks, banding at 57				
145.10	149.55	BIOTITE GNEISS	15% biotite laths, banded, 15-20% felsic layers, rare garnet porphyroblasts (2mm in size), fine grained, rare qtz and qtz-plagioclase veins (about 1cm wide), locally up to 1% pyrite associated with the felsic layers/bands, foliation at 60, trace disseminated pyrrhotite				
149.55	156.80	GARNET BIOTITE GNEISS	Banded, 15% garnet porphyroblasts (<0.5cm in size), fine grained, 20% light grey felsic layers/bands which lack garnets but locally contain up to 20% amph laths (<1cm in size), rare qtz veins (<1cm wide), trace disseminated pyrite and pyrrhotite				

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
156.80	158.70	SHEAR ZONE	Banded, purplish brown color, 30% qtz veins (<1cm wide), 40-50% biotite rich bands, 15% actinolite bands, 5% plagioclase, 1% pyrite specks and trace disseminated pyrrhotite, rusty fracture surfaces,	156.80	157.80	1.00	94700
				157.80	158.70	0.90	94701
158.70	159.10	GRAPHITE SCHIST	40% Graphitic layers/bands at 33, dark black, 5% k-feldspar alteration, 30% qtz-plagioclase veins/flooding, locally disked, 10% biotite rich bands, 3% Pyrite>Pyrrhotite blebs	158.70	159.60	0.90	94702
159.10	162.35	BIOTITE GNEISS	Layered/Banded, fine grained, rare graphitic bands, 5% qtz-plagioclase veins (<2cm wide), 5% qtz-plagioclase healed fractures, bands at 44; 3% pyrite>pyrrhotite blebs, veins and occasionally associated with the healed fractures	159.60	160.60	1.00	94703
				160.60	161.50	0.90	94704
162.35	168.50	WACKE	Layered/banded, 15-20% biotite, grey to pinkish grey, fine grained, 50-60% of the layers have a pinkish color due to k-feldspar, 10% of the layers contain garnet porphyroblasts (<0.5cm in size), trace disseminated pyrite and pyrrhotite throughout, <5% qtz-plagioclase healed fractures, moderate to strong foliation at 47 163.45-164.20: 2 qtz-veins (12cm + 25cm) at 50, 3% Pyrite specks and blebs + 5% pyrrhotite specks and blebs along the contacts of the veins and within the veins themselves	163.45	164.20	0.75	94705
168.50	180.65	WACKE	Fine grained, medium grey, locally banded/layered, fresh surfaces are fairly felsic (qtz-rich), rare qtz-plagioclase healed fractures, 5% biotite, <1% disseminated pyrite 170.45-172.90: Siliceous section, light grey, 10% pinkish colored k-feldspar rich bands (<2cm wide), 15-20% qtz-plagioclase rich layers-almost white in color, 5% pyrite disseminated and specks, banded at 57, gradual contacts with the wacke 174.90-177.00: Becoming more medium grained, strong foliation, 25-30% hornblende laths, med grey color, >2% pyrite along fracture surfaces as blebs, foliation at 48, rare qtz veins (.1cm wide), <5% qtz-plagioclase-k feldspar healed fractures, fresh surfaces are fairly felsic (qtz rich), trace pyrite disseminated and specks	170.45	171.60	1.15	94706
				171.60	172.90	1.30	94707
180.65	182.90	FELSIC SEDIMENTS	Light grey, siliceous, rare pinkish bands (k-feldspar rich), fine grained, banded at 48, 10-15% weakly chloritized bands, no visible biotite, trace pyrite 182.00-182.70: 20% qtz-plagioclase healed fractures, <5% garnet porphyroblasts (<0.5cm in size)				
182.90	186.00	BIOTITE GNEISS	Pale to medium grey color with 60% purplish colored biotite rich bands, fine grained, mafic bands are slightly greenish due to weak actinolite alteration, trace disseminated pyrite, bands at 33 184.70-185.00: Intermediate dyke, fine grained, greenish grey color, sharp contacts at 60				
186.00	188.40	CONTACT	Gradational contact, zone of assimilation, banded pale grey and purplish, very siliceous, 5% qtz veins (<2cm wide), 10% qtz-plagioclase veins/flooding, 5% chlorite streaks, local specks of a bright green mineral (looks similar to malachite)				

From	To	Rock Type	Description (Texture, Structure, Alteration, Mineralization)	Sample			
				From	To	Length	Sample #
			187.00-187.10: 5% talc as specks and streaks				
188.40	191.55	GRANITE	Light grey to white, 15% biotite laths, weak to moderate foliation at 55, 5% qtz and calcite fracture fill 190.30-190.40: Angular fragments, possibly gabbro xenoliths				
191.55	201.00	GABBRO	Fine to medium grained, dark grey, massive, rare qtz and calcite healed fractures 191.55-194.20: 5% qtz veins (<5cm wide), 5% qtz-plag-calcite veins (<1cm wide), weak foliation at 40 192.70-192.80: Granitic dyke (as above), contacts at 25				
			END OF HOLE AT 201.00m				