

Report on 2004 Summer Drill Program

Hunter Mine Property

ValGold Resources Ltd.

Whitney Township

Porcupine Mining District

Ontario

2 - 30149



**Dr. Derek McBride P.Eng.
Kiam Jensen P.Geol.
Toronto, Ontario
December 30, 2004**

Table of Contents

Summary	i
Introduction	1
Location	1
Geological Mapping	1
2004 Drilling	4
Geology of the Mine Section based on drilling	13
Interpretation	13
Conclusions	15
Appendix 1 Summary Drill Logs	17
Appendix 2 Assay Certificates	18

List of Figures

1	Location Map	2
2	2004 Drill Holes Hunter Property	3
3	Drill Hole Section HM-04-28, 29, 30 & 39	6
4	Drill Hole Section HM-04-31, 32 & 33	8
5	Drill Hole Section HM-04-34 & 35	9
6	Drill Hole Section HM-04-36	11
7	Drill Hole Section HM-04-37 & 38	12
8	Interpretation of Mine Horizon	14

List of Tables

1	Drill Hole Summary	5
2	List of Assays over One Gram, 2004 Drilling	5
3	Proposed Drilling Budget 2005 Program	15
4	Project Costs 2004 Drilling	16

Summary

In June 2004, ValGold Resources Ltd. undertook a drilling program on their Hunter Mine Property. Drilling was carried out on Porcupine Lake by Benoit Drilling of Val d'Or, Quebec using their barge. Twelve holes were drilled to test the gold-bearing shoots indicated by winter drilling in 1985-1986.

Drill results indicated an anomalous gold-bearing horizon that was traced for 200 metres along strike, 150 metres vertical and remains open at depth and along strike in both directions. Within this horizon is a shoot that carries values in the 11 to 18 gram per tonne range over one and a half metres or better. This shoot plunges from surface in the shaft area in a north, northeast direction at an angle of approximately -30 degrees.

A drill program is recommended in 2005 to trace this shoot northeastward and at depth. Additional shoots were indicated in the old work in the shaft area; holes will test their extent. The final part of this program will investigate the horizon to the southwest where it strikes towards the Dome Superpit and has never been investigated. A budget of \$500,000 is proposed for this work.

Introduction

In early summer 2004, a drilling program was undertaken to evaluate the Hunter Mine mineralization at depth. To perform this program, the drilling was carried out from the barge of Benoit drilling of Val d'Or. Twelve holes tested the mineralized horizon from the 130 metre to the 250 metre level.

Location

The Hunter Mine Property is situated south of Highway 101 in Porcupine, Ontario along the east shore of Porcupine Lake all the way to the Dome pump house east of South Porcupine (Figure 1). A separate claim block extends the property to within the town limits of South Porcupine and two kilometers of the Dome Superpit. Drilling was on claim HR 1009.

Geological Mapping (Map 1 in pocket)

Mapping was conducted in the fall of 2003 over the Hunter Mine Claims and along the shore of Porcupine Lake. Outcrops on the property lie mainly along the shore of Porcupine Lake with the best exposures being adjacent to the shaft. This large stripped area is underlain by a sequence of bedded carbonate-rich exhalitic sedimentary rocks. Individual beds can be easily identified and traced for considerable distances. A sugary quartz vein (band) has been pitted and the pit on the lakeshore marks the first discovery pit in the Timmins' area.

From footwall to hanging wall (east to west) these rocks form a sequence consists of a basal calcium carbonate rich weakly sericitic exhalitic unit that is pale grey in colour and weathers to a pale brown because of some contained iron carbonate. The top of this unit is a narrow (<0.3m) band of medium grey fine clastic sediments. Conformable with the top of this band is the 20 centimetre sugary quartz band that forms the core of the mineralized horizon. The hanging wall is a thick sequence of iron carbonate-rich sedimentary rocks that in outcrop are diagnostic by the pervasive quartz-filled tension fractures. South of the main outcrop area these rocks are overlain by dark blue-black, very soft ultramafic rocks. North of the main outcrop the exhalitic sequence can be traced to just south of the property boundary, Highway 101. East of the main road south from Porcupine, Haileybury Cres., clumps of trees mark very old pits. These are now badly caved, but the southerly one contains fine-grained medium grey bedded clastic sedimentary rocks.





Figure 2
ValGold Resources Ltd.
Hunter Mine Property
 Whitney Township, Porcupine Mining District
 Ontario

**2004 Summer
 Drill Program**



D. McBride, January 2005

Drilling in 1985 expanded the mine section to show that the mine horizon is flanked by ultramafic units which are in turn flanked by clastic sedimentary rocks. Bedded felsic lapilli and ash fall tuffs were observed in the southeast corner of the property.

2004 Drilling (Table 1)

Drilling commenced on June 11, 2004 with hole HM 04-28 (Figure 3) which was designed to duplicate previous hole S-16 which had returned an intersection of 24.41g per tonne over 2.74m (0.7136oz per ton over 9ft). The hole was collared at the calculated location of hole S-16 based on earlier maps. When the barge was attempting to move to this location it hit the casing of S-16. The new hole was located approximately 13 meters ahead of S-16 on section. The lake bottom was hit at 2 metres and the casing continued to 55 metres through lake-bottom clays and basal till. Ultramafic talcose fragmental volcanic rocks compose the next 29 metres. These rocks abruptly change into finely laminated soft ultramafic tuffs at 84.1 metres. Below, the contact with the laminated carbonate-rich tuffs is quite sharp and is marked by a rapid hardening of the rock and a change in colour from a medium grey to a brownish grey at 105.1 metres.

The next section is composed of laminated carbonate exhalitic sedimentary rocks that are historically referred to as the altered zone. It is at the top of this unit that the gold mineralization is located; an assay value of 35.56g per tonne over 0.78 metres from 105.2 to 105.9 metres shows the presence of significant gold beyond the historical mine workings and confirms the results from the 1985-1986 drilling (Table 2). Below the mineralized intersection the rock continues as a laminated sequence to the end of the hole at 195.5 metres (Appendix 1 for summary log and 2 for detailed log). Extensive assaying of these rocks did not detect additional mineralized horizons.

The second hole, HM04-29, was drilled from the same location at -75 degrees to intersect the mineralized horizon at the 180 metre level or 35 metres below S-16. It intersected overburden to 55.5 metres followed by the ultramafic talcose fragmental volcanic rocks to 104.7 metres. These rocks are the same as described in the previous hole. At approximately 132 metres, the transition to the exhalitic sedimentary rocks occurs. Sampling of this section is in progress. Continuing down section, the rocks continue their exhalitic sedimentary character commonly showing a pale buff colour and locally the yellow-buff of leucoxene. Bedding is evident throughout the sequence and becomes prominent lower down. From 245.8 to 252.4 metres a section of medium to dark grey argillites is present. Bedding in this section is 75 to 80 degrees to the core axis. Below this argillite the rocks are pale grey siliceous clastic sediments to the end of the hole at 303.2 metres.

Hole HM04-30 was drilled from approximately the same location as the previous two holes to further test this horizon on this section at the 250 metre level. It was cased to 55 metres before entering the talcose ultramafic volcanic rocks. These continue to 107 metres before rapidly grading into the laminated variety of ultramafic tuff. The laminated tuff continues to 139.5 metres where it grades into the bedded exhalitic, tuffaceous sediments. This sequence goes to the end of the hole at 258 metres becoming more

Table 1

**DRILL HOLE: Summary Summer Program
2004**


HUNTER MINE PROJECT

Hole No.	U T M			Direction	Dip	Length m	Acc. Length m	Started	Finished
	Long.	Lat.							
HM04-28	5370930.4	487135.0		105 Az.	-50	195.5	195.5	June 11	June 14
HM04-29	5370930.4	487135.0		105 Az.	-75	303	498.5	June 14	June 19
HM04-30	5370931.0	487130.1		105 Az.	-85	258	756.5	June 22	June 28
HM04-31	5370891.3	487116.9		105 Az.	-50	243	999.5	June 28	July 2
HM04-32A	5370891.3	487116.9		105 Az.	-80	63	1062.5	July 2	July 2
HM04-32B	5370891.3	487116.9		105 Az.	-80	255	1317.5	July 2	July 6
HM04-33	5370904.0	487073.0		105 Az.	-80	237	1554.5	July 6	July 9
HM04-34A	5370818.4	487097.3		105 Az.	-67	117.8	1672.3	July 10	July 12
HM04-34B	5370818.4	487097.3		105 Az.	-67	167.9	1840.2	July 12	July 15
HM04-35	5370818.4	487097.3		105 Az.	-85	249	2089.2	July 15	July 19
HM04-36	5370884.7	486989.3		105 Az.	-90	234	2323.2	July 20	July 24
HM04-37	5370768.3	487021.0		105 Az.	-66	249	2572.2	July 24	July 28
HM04-38	5370768.3	487021.0		105 Az.	-80	219	2791.2	July 28	July 30
HM04-39	5370982.0	486941.0		105 Az.	-75	309	3100.2	July 30	August 3

Table 2

**List of Assays over One Gram per Tonne
From 2004 Drill Program**

1	Hole	Sample	From	To	Width	Au (g/t)	Au			Metallic
							check	Au (2nd)	Au check	
5	HM04-28	44512	105.12	105.90	0.78	32.88	37.85	37.03	34.49	37.52
148	HM04-30	44607	146.10	147.41	1.31	1.51	1.49			2.06
195	HM04-31	44654	113.00	113.60	0.60	2.61	1.89	2.25		2.86
197	HM04-31	44656	114.70	115.80	1.10	2.06		2.06	2.266	2.28
198	HM04-31	44657	115.80	116.30	0.50	36.48	36.82	36.65	18.325	31.60
210	HM04-31	44669	127.05	127.67	0.62	1.17	0.90	1.035		1.14
331	HM04-32	44790	142.00	142.64	0.64	2.32	2.59			3.33
350	HM04-32	44809	158.95	159.36	0.41	1.06	0.98			0.59
522	HM04-34A	44981	130.58	131.75	1.17	1.03	0.79			0.81
523	HM04-34A	44982	131.75	132.00	0.25	1.86	1.95			1.71
616	HM04-35	43075	156.00	156.47	0.47	0.22		VG		0.61
629	HM04-35	43088	165.74	166.28	0.54	1.40	0.96			1.10
872	HM04-37	35131	171.74	172.28	0.54	1.18	1.13			0.61
874	HM04-37	35133	172.78	173.43	0.65	0.89	1.16			1.17
937	HM04-37	35196	227.42	227.72	0.30	1.01	0.94			0.01
976	HM04-38	35235	146.87	147.76	0.89	1.13	1.05			0.06
1036	HM04-38	35295	187.58	188.50	0.92	1.85	1.87			2.15



D. McBride, January 2005

Note: Plane of section is 15° to UTM Grid


sedimentary, down section. A medium to dark grey argillite section occurs from 216.7 to 227.8 metres and probably correlates with the section at 245.8 metres in the previous hole. The mineralized horizon is present in this hole from 146.1 to 147.41 metres which returned 1.5 g per tonne over 1.31 metres.

Drilling continued 30 metres to the south on the section of Hole S-15; this hole contained an assay of 24.5 g per tonne over one metre (0.761 oz per ton /3.3ft.). Hole HM04-31 (Figure 4) was designed to intersect the mineralized horizon above S-15. It was cased to 64.3 metres before entering the talcose ultramafic tuffs. They continue to 82.7 metres where they meet the laminated variety which continues to 113 metres. Below these ultramafics are the exhalitic sedimentary rocks; they continue to the end of the hole at 242.2 metres. This hole returned 4.03 g per tonne over 3.3 metres which included 36.65 g per tonne over 0.50 metres just below the upper contact with the laminated ultramafic rocks. 11 metres below a second mineralized section gave 1.04 g over 0.62 metres. Below these values the gold values drop and seldom exceed 0.10 gram per tonne. The next hole, HM04-32 was designed to the zone at the 180 metre level; it passed through the lake and overburden to 58.4 metres before intersecting the talcose ultramafic fragmental volcanic sequence. These ultramafics continue to 87.4 metres where they rapidly grade into the laminated type which terminates at 138.1 metres. Below these rocks the exhalitic sedimentary section continues to the end of the hole at 255 metres. The medium to dark gray or graphitic argillite is found from 228 metres to the end of the hole. The mineralized horizon is present from 142.0 to 142.64 metres and ran 2.46 g per tonne over 0.64 metres. A second zone is present from 158.95 to 159.36 metres and contains 1.02 g per tonne over 0.41 metres.

Hole HM04-33 was drilled on the same section to intersect the mineralization at the 250 metre level. It was cased to 56.8 metres before entering the talcose ultramafics. They continue to 87.4 metres and rapidly grade into the laminated variety which in turn grade into the exhalitic sediments at 134.1 metres. These exhalitic sediments continue to 230 metres where they grade into dark gray argillites to the end of the hole at 237 metres.

Holes HM04-34 and 34B (Figure 5) were drilled below S-14, 100 metres south of S-16. The first attempt was lost at 117.8 metres; but the second attempt was able to penetrate the exhalitic sediments. It was cased to 59.7 metres and penetrated the talcose ultramafics to 75 metres. Below 75 metres it passed through the laminated tuffs to the exhalitic sediments at 128.8 metres. This sequence continues to the end of the hole at 167.9 metres. The mineralized horizon consisted of a 1.42 metre intersection which returned 1.4 grams of gold per tonne from 130.58 to 132.0 metres.


Hole HM04-35 continued to test this section at the 250 level. It was cased to 75 metres where it encountered the talcose ultramafic rocks; these continued to 94.5 metres. At 94.5 they contact the laminated ultramafic tuffs that extend to 133.8 metres where they contact the lower exhalitic sedimentary sequence. The exhalitic sedimentary sequence continues to the end of the hole at 248.9 metres. Dark gray argillites that form the lower marker horizon extend from 237.5 to 240 metres. Visible gold occurs in the section from 156.0 to



D. McBride, January 2005

Note: Plane of section is 15° to UTM Grid

32.88/0.78 Grade (g/t Au) / width (metres)



D. McBride, January 2005

Note: Plane of section is 15° to UTM Grid


156.47 metres, but the assayed value was 0.22 grams per tonne. A metallic assay returned 0.61 gram per tonne. A second gold-bearing zone is located from 165.74 to 166.28 metres and returned a value of 1.18 grams per tonne over 0.54 metres. This hole completed the investigation of the Hole S-14 section.


The next hole, HM04-36 (Figure 6), was drilled on section of hole S-10 and was designed to cut the mineralized horizon at the 250 metre level. It was cased to 57 metres and penetrated the talcose ultramafic rocks for 60 metres to 117.6 metres. The rocks rapidly change to the laminated ultramafic tuff which goes to 138.9 metres. Beyond is the exhalitic sedimentary sequence which continues to the end of the hole at 243 metres. Graphitic argillite of the lower marker horizon is located from 221 to 224 metres. Mineralization was not observed in this hole, but a one metre section from 140.18 to 141.18 averages 0.43 grams per tonne. It probably represents the mineralized horizon.

The next two holes were the most southerly drilled; they tested the Hole S-20 section about 40 metres north of the shaft or 150 metres south of Hole S-16. They were designed to test the down dip continuation of the 4.24 gram per tonne over 0.67 metre intersection in Hole S-20. Hole HM04-37 was aimed at the 180 metre level; it was cased to 63 metres and continued in talcose ultramafic rocks to 105 metres. Below these rocks the laminated ultramafic tuffs go to 167.8 metres and contain a siliceous feldspar porphyry band near its base. Exhalitic sedimentary rocks continue to the end of the hole at 248.8 metres. The last three metres is the dark gray graphitic argillite. Mineralization is weak in this hole and is represented by a 2.12 metre section, from 171.3 to 173.4 metres, which ran one gram per tonne.

Hole HM04-38 (Figure 7) probed this section at the 250 metre level. After casing to 59.3 metres it penetrated talcose ultramafics for 36 metres to 95.6 metres where it contacted the laminated ultramafic tuffs. These rocks extend to 143.5 metres. Below, the exhalitic sedimentary sequence extends to the end of the hole at 218.6 metres. The lower marker graphitic argillite starts at 213.3 metres and continues to the end of the hole. The mineralized horizon is present from 146.35 to 147.76 metres as a 1.4 metre section containing one gram per tonne.

The last hole in the program was designed to test the S-16 section at depth. It was collared approximately 100 metres west of the first ring of holes. According to the interpretation of Kirwin, 1988 (OGS Assessment File T2664) this hole should have intersected the exhalitic sedimentary rocks at between 130 and 200 metres. The hole was cased to 49 metres before entering the talcose ultramafic fragmental tuffs and agglomerates. It continued in these rocks to 275.8 metres and then passed through a laminated ultramafic band before returning to the talcose rocks to 305 metres. The last 4 metres to 309 metres was a medium to dark gray highly sericitic schist that was probably a fine dark mud. The trend of this unit appears to be within 20 degrees of the core axis. It is unlike any of the other rocks seen in the adjacent holes. The writer is interpreting these sediments to be a band within the talcose ultramafic rocks and the absence of the normal mine section suggests that the dip is much steeper than previously thought.





D. McBride, January 2005

Note: Plane of section is 15° to UTM Grid

After the drill program was completed, all samples containing greater than one gram of gold were re-assayed using the metallic assay. This method includes coarse gold in the sample and gives a total gold assay for the sample. The right column shows the metallic assays; for the higher values, the metallic assays are similar or slightly elevated. They show that the nugget effect is neutral or slightly positive.

Geology of the Mine Section as based on 2004 Drilling


The lowest rock in the mine section is the exhalitic sedimentary sequence. It can be divided into a lower middle and upper sequences. The lower lies below the graphitic argillite and consists predominantly of pale gray siliceous clastic sedimentary rock with a high calcium carbonate content. Mineralization is not known in these rocks. Above these rocks the graphitic argillite sequence marks a transition from predominantly clastic sedimentary rock to more exhalitic and tuffaceous ones. It is a two to five metre thick series of beds that can be traced between holes.

The main exhalitic sedimentary section or upper series of beds is fairly well bedded and contains abundant carbonate. It is a pale brownish gray in colour and is best differentiated from the other lithologies by its colour and high carbonate content. Gold is found within the top few metres of the unit. On surface at the discovery pit this rock weathers light brown from the iron in the carbonate. A quartz band marks the upper limit of the band and contains the gold in the discovery pit. From the drill intersections, the gold mineralization is essentially conformable with the upper contact of the exhalitic unit.

Above this contact is the laminated ultramafic tuff. It is a well laminated rock with dark talcose beds and paler gray carbonate ones. Up section the talcose bands increase relative to the carbonate and take on the appearance of striped black and pale gray rock. The transition to the overlying talcose ultramafic rocks is sharp. These rocks form a sequence of bedded dark green, soft tuffaceous rocks and agglomerates. Lapilli, fragments and bedding are diagnostic of this rock as is its very soft matrix. It forms the upper most unit seen in all drill holes and in Hole HM04-39 has a core length of more than 300 metres.

Interpretation (Figure 8)

The 2004 summer drill program has extended the mineralized horizon to the 250 metre level. Gold values very near the top of the exhalitic sedimentary sequence show the continuity of this gold-bearing horizon as a conformable band in the stratigraphy. These observations agree with those mapped on the lakeshore outcrop north of the shaft. Within this gold-bearing horizon, a higher grade ore shoot with visible gold can be traced from the old stopes as a -35 degree plunging structure approximately 100 metres long. Old hole S-18, 30 metres north of the most northerly hole in this program, returned a 3.2 metre section grading 0.44 grams of gold with visible gold observed in the core. A further 60 metres north a 1.6 gram gold assay over 0.3 metre indicates that the gold-bearing horizon continues north. Unfortunately the geology seems complicated from the old log and the core is no longer available. Hole HM04-39 tried to test this horizon well below Hole HM04-30. It failed to penetrate through the talcose ultramafic rocks before being



terminated at 309 metres and shows that the horizon is steeper than previously interpreted.

Conclusions

The drilling program in the summer of 2004 has succeeded in defining a 100 metre long gold shoot within an extensive horizon that contains more than a gram of gold over more than a metre. This shoot can be traced from surface northerly with a plunge of 35 degrees to below the 180 metre level beyond the northern limit of the program. Future drilling should attempt to trace the shoot to the north and down plunge.

South of this shoot the lower values that the horizon continues and an old value in Hole S-12 of 40.9 grams over 0.46 metres (1.195 oz per ton over 1.5 ft.) indicates that an additional gold-bearing shoot may be present in the shaft area. In addition only four S series holes tested the horizon south of the shaft and only for 300 metres. Gold values were in the trace range, but much of the sampling did not include the gold-bearing horizon as it is now understood. More drilling is warranted in this area and along the horizon to the south to the end of the claims at the Dome Mines pump house.

This recommended program will test the three principle target areas on the property with 3500 metres of drilling. 2000 metres will test the continuity of the north gold shoot, 700 the shaft gold shoot and 800 the southern extension. The total estimated cost of this program is \$500,000 and is detailed in Table 3.

**Table 3
Proposed Drilling Budget 2005**

Drilling 3500 metres: down dip & extension \$100.00 per metres	\$350,000.00
Supervision and technical analysis	\$40,000.00
Magnetic and EM surveys for entire property	\$15,000.00
Local Field Assistant 60 days at \$150.00 per	\$9,000.00
Travel and accommodations 1 and a half months	\$12,000.00
Field office and storage	\$10,000.00
Supplies, services and assaying	\$10,000.00
Mob., demob. of drill and barge	\$20,000.00
Reporting and government filings	\$10,000.00
Contingency	\$24,000.00
Total	\$500,000.00

The project cost totaled \$430,624.96. Table 4 shows the breakdown from the accounting department of ValGold.

Table 4
Table of costs

Drilling contract costs Benoit Drilling	\$	340,638.87
Assaying Swastika and Acurrassay Labs.	\$	16,631.00
Core storage, sampling and hole surveying	\$	17,582.13
Project geologist	\$	38,675.32
Project management	\$	17,097.64
Total Project Cost	\$	430,624.96

APPENDIX 1
Drill Logs

Hunter Mine - Diamond Drill Log

Property:	Hunter Mine	Hole Dip	-50	Page No.:	1 of 8	Hole HM-04-28				
Location:		Hole Azimuth:	105o	Date Started:	11/06/2004					
Claim No:	HR 1009	Hole Length	195.5m	Date Finished:	11/06/2004					
Elevation:	Porcupine Lake	Purpose:	test zone	Drill Co.:	Benoit					
UTM Coords.:	5370930. 4N, 487135.0E			Logged by:	K. Jensen					
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
0.0	60.0	Casing - water depth 7'								
60.0	102.5	Talcose Carbonated Brecciated UM								
		note 60.0-63.0 reamed core recovery 1.56								
		med., deep bluish tint, very talcose, fg, black to greenish black, carbonated, soft to moderately soft 60-73.6 to moderately soft 73.6 poorly magnetic with intensely carbonated sections to weakly magnetic, brecciated healed with carbonate, rare qc stringers <0.5cm CA-30o, contorted foliation to CA-50o, 60.0-73.6 intensely carbonated, rare buff green elongated fragments, 61.87 5mm qc CA-30o, pale grey white mg elongated fragments 67.16, 67.35, carbonated brecciated and irregular masses, 72.50 2cm qc veinlet CA-75o, 73.6-74.40 intensity of carbonate brecciated decreases to <10%, and hairlike to 1-2mm carbonate, 74.40-77.15 foliated locally contorted possibly tuffaceous carbonated UM, CA-58o see pyrite								
77.15	79.4	at 74.89, contorted 76.43-77.15, 77.15 ground contact.								
		Lamprophyre fg mg, salt and pepper brownish altered hornblende masses and carbonated, locally bluish tint, carbonate stringers, CA-30-35o with bluish hue, moderately soft, carbonated moderately, non magnetic, nil to poorly developed foliation locally pale buff greenish yellow sericite altn, random x-cut orientation, rare qc or c, qtz stringers, trace sulphides, 79.40 ground contact, 79.37 3cm								
79.4	79.9	crumbly core, possible fault/shear.								
		Lamp and Carbonate UM tuff, similar to lamp but looks like alteration contact recrystallization, weak alignment foliation CA-68o, mottled to salt and pepper texture of black to black green and pale greenish white carbonate, discontinuous foliated gashes < 2mm, x-cutting foliation, rare scattered cg py.								
79.9	83.87	moderately soft, non magnetic, moderately devel. foliation, 79.90 contact CA-60o. TCS carbonated UM tuff talcose, fg, laminated with whitish to whitish grey carbonate, uniform, moderate soft, non magnetic, well developed of bedding CA-40o foliation with locally contorted folded with displacements 1.5cm 81.18-82.44 CA-80o chlorite fol., scattered mg py associated with discontance or carbonate bands <0.5% to trace, 81.18 3cm x-cut bed, V shaped light brownish qc veinlet								

Property		Hole No.	28		Sheet No.	2				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au Check	Au (2nd)	Au check
79.9	83.87	CA-80o+60o to 83.87 contorted and folded bedded tuff, 83.87 3cm shearing intense foliation with carbonated stringer CA-70o.								
83.87	90.03	Massive to locally Bx UM talcose, fg, black green, moderately soft, non magnetic, massive, uniform, randomly orientated 1-3mm carbonated ff, nil to weak level of schistose or foliation, void of stringers qtz or qc or c, nil to trace sulphides usually py cg py at 83.04 associated with discontinuous carbonated, 84.30 1.2cm pyrite, 84.45 mg py, 88.39-88.66 brecciated carbonated healed, 88.66-89.11 very talcose, 89.11-89.19 very schistose, sheared zone CA-75o+85o, 89.19-90.03 intense brecciated carbonated healed 50%.								
90.03	90.3	silicified carbonated altn felsic, fg light grey brown, non magnetic, weakly to moderately soft, massive, uniform, possible contact alteration, 90.30 broken contact, chlorite ff small and discontance.								
90.3	91.02	Altered UM, similar to 77.15-79.40, fg, brownish tint to black green, well developed talcose, soft to moderately soft, crushed zone, nil to very weak developed foliation/schistose, nil stringers, trace sulphides, carbonated, talcose, 91.02 contact CA-40o overall sineous sharp.								
91.02	91.23	qtz vein, white, milky with pale green talcose and black green UM fragments inclusions, 91.23 contact sharp sinuous overall 45o.								
91.23	94.94	Tuff UM, carbonated, talcose, fg, massive to tuffaceous, greyish altered, siliceous 91.23-91.53, black to black green with carbonate greenish white, increasing development of bedding 1-2mm laminated, moderately soft, non magnetic, bed at 92.57 CA-72o, 93.79 CA-70o, 94.45-94.64 contorted bedding, trace sulphides, 94.94 contact sharp CA-75o.	44501		94	94.94	0.94	0.02		
94.94	96.62	Grey felsic dike, ophanitic, light to medium grey, hard, siliceous, scattered medium to dark green x-cut chlorite, non magnetic, non carbonated, few scattered 1-2mm white qtz stringers CA-60-85o from 95.46-95.94, scattered light grey qtz veinlets 1cm CA-60o irregular at 96.14, white qtz vein 2-3cm irregular CA-80o at 95.41, scattered fg py with few mg py overall 1% locally 1-2% from 94.94-96.14, 96.14-96.62 similar to 94.94-96.14 except nil to trace sulphides, 96.42 increasing greyish white q veinlets to 96.62, contact CA-50o	44502		94.94	96	1.06	0.03	0.04	
96.62	97.2	FP or silicified massive UM, aphanitic medium grey with locally 1mm phenocrysts, siliceous and several greyish with qtz veinlets, 1-1.5cm, hard,	44503		96	97	1	0		

Property		Hunter Mine		Hole No.	28		Sheet No.	3			
Meterage		Description		Sample			Assay				
From	To			No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
96.62	97.2	non magnetic, rare sections of <0.5% fg py, nil level foliation, 97.20 contact sharp CA-80o.									
97.2	97.43	TCS UM tuff, fg, black green, carbonated foliation CA-55o, moderately soft, non magnetic, talcose, possible inclusions, void of stringers, nil sulphides, 97.39 contact sharp CA-60o.		44504	97	98	1	0			
97.43	97.55	Felsic dike, light to medium grey, aphanitic, massive uniform, nil level foliation void of stringers, void of sulphides, fg black green clots similar to 94.94-96.62 96.62 contact sharp CA-65o.									
97.55	98.12	Silicified zone, felsic dike, massive greyish white qtz veining with inclusions of medium grey with black green clots FP similar to 97.39-97.50, void of sulphides, hard, non magnetic, non carbonated, void of second generation stringers, 98.04 contact sharp CA-50-55o.									
98.12	98.69	Silicified zone, UM, massive ultramafic intruded by weakly carbonated veins and silicified ultramafic altered to light grey veins white black green core, qtz veinlets random, nil sulphides, 98.04-98.43 silicified 98.43 CA-70o, 98.43-98.62 silicified foliated black green UM, 98.62 CA-50-60o irregular.		44505	98	99	1	0			
98.69	98.95	Felsic dike and qtz vein, similar to 97.43-97.55 with <1mm white phenocrysts hard, nil sulphides, 98.69-98.74 whitish qtz vein CA-75o, 98.78 1/2cm CA-75o, 98.84-98.95 qtz vein with chlorite UM and talc inclusions 98.95 CA-55o no sulphides.									
98.95	105.13	Silicified UM tuff and Bx, fg, blackish green to black, fine laminations, moderately hard to hard silicified sections, talcose nil to weakly carbonated altered to medium grey to blackish grey, irregular discontinuous carbonated and qtz masses, well developed bedding with local contorted bedding small folds, locally brecciated, 99.29-99.44 white qtz vein CA-70o+55o no sulphides, 98.95-99.57 brecciated, 99.57-99.91 tuffaceous bed CA-55o minor kinkling, 99.91-100.02 qtz vein white CA-80o + 40-45o irregular, 100.02-100.85 80o silicified light medium grey, brecciated, 100.85-101.33 55o brecciated, folded, black green, irregular, 101.33-101.49 greyish qtz vein with altered UM and talc CA-75o overall, 101.49-101.94 brecciated black green, with irregular greyish qtz masses, 101.94 ground contact, 101.94-102.49 silicified brecciated UM CA-65o,		44506	99	100	1	0			
				44507	100	101	1	0	0		
				44508	101	102	1	0.01			
				44509	102	103	1	0.02			
				44510	103	104	1	0.03			
				44511	104	105.12	1.12	0.16			

Property		Hunter Mine		Hole No.	28			Sheet No.	4		
Meterage		Description		Sample			Assays				
From	To			No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
102.49	105.13	Laminated Ultramafic Tuff									
98.95	105.13	black green and greyish qtz carbonated, weakly carbonated to moderately, well developed bedding CA-65o at 102.8, scattered x-cutting chlorite fol. CA-23o, randomly 1-2mm whitish qtz stringer CA-20o, 55o & 40o opposite, bed at 104.16 CA-60o, 104.3 1 1/2cm brownish siliceous aphanitic felsic dike CA-45o, 104.43 3-4mm qtz whitish to pale green stringer x-cut bedding, CA-15-20o, 104.68-104.90 irregular greyish qtz veinlet with whitish ankerite carbonate on contacts low angle and II to bed, 105.13 contact CA-40-45o sineous.									
105.13	195.5	Exhalitic Sedimentary Rocks and Tuffs									
105.13	105.28	Qtz vein, pale brownish tint qtz vein, UM inclusions, minor py at 105.13, contact 105.28 CA-30o overall sinuous, pin prick VG at 105.23, inclusions rimmed with chocolate brown tourmaline.		44512	105.1	105.9	0.78	32.88	37.85	37.03	34.49
105.28	107.44	Bleached felsic tuff, aphanitic to fine grained, light buff to greyish tan, random rare 1-2mm qtz eyes, well developed laminations tuffaceous, siliceous, hard, non magnetic, 3 local spots with green fuchsite, 1 as small fragment, qtz and qtz ankerite stringers II to bedding CA-35-45o, vfg to fg pyrite associated mostly with tuff minor in stringers, scattered fg chalcopyrite, overall 1-2% pyrite, 105.28-105.54 greyish buff with qtz stringers and tourmaline scattered chalcopyrite, 1-2% vfg to fg pyrite in tuff, scattered VG at 105.34 pin prick, 105.43 2 on both sides of qtz veinlet and tourmaline CA-30o 1 spot 0.5mm long cluster and qtz and ankerite stringers x-cut bedding CA-20-40o, 105.54-107.44 buff felsic tuff, 3 fuchsite 1 fragment, well developed bedding, random chocolate brown tourmaline hairlike usually II to bedding, locally x-cut by qtz ankerite stringers, brighter density from 105.54-106.05 and 107-107.44, 105.73 1-2mm cluster of VG associated with grey qtz stringer CA-10-30o x-cut bedding CA-45o with brown tourmaline on contacts, 107.39-107.44 greyish buff silicified contacts with II brown tourmaline.		44513 44514	105.9 106.7	106.7 107.4	0.75 0.79	0.22 0.16	0.21		
107.44	116.17	UM tuff, similar to 102.49-105.13 minor kinkle folding, discontinuous and qtz stringers II to bedding Ca-42-45o, 107.44-112.10 bedding x-cut by chlorite stringers with usual small displacements CA-25-30o 1-5mm up to 20/meter, about 50:50 x-cuts and displaced qtz stringers II to bedding 10-15/meter, scattered to trace vfg fg pyrite <0.5% overall, 111.08-111.15 greyish buff felsic tuff band, 2-3% vfg pyrite CA-80o, 111.24 contorted felsic band II to bedding but		44515 44516 44517	107.4 108.4 115.0	108.4 109.4 116.2	1 1 1.17	0.01 0.1 0.24			

Property: Hunter Mine		Hole No.	28		Sheet No.	6				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		siliceous nil to weak development of schistosity or bedding, 1% fg mg py, 122.57 sharp contact CA-70-75o.								
122.57	123.12	Felsic dike, aphanitic to fg, pale brownish buff to pink tint buff, well developed foliation with chlorite not laminated, random qtz stringers 2mm CA-12o x-cuts foliation CA-65o and qtz stringer CA-45-50o scattered fg pyrite <0.5%, 123.12 CA-80o irregular.	44524	122.24	123.12	0.88	0.43			
123.12	124.36	Chlorite silicified UM tuff, same as above, blackish green, tuff, silicified, hard, non carbonated, non magnetic, trace sulphides, irregular qtz veinlets and sections, well level bedding CA-irregular, 123.21-123.40 qtz zone, minor vein 123.10-123.29, CA-60o with inclusions, 123.60 1-2cm qtz vein CA-85o, 123.40 123.81 contorted bedding, large fold at 123.72, 123.81-123.91 qtz section CA-60o, 124.10-124.36 qtz carbonated vein with numerous chlorite, tuff inclusions scattered sulphides CA-70-80o irregular.	44525	123.12	124.36	1.26	0.02			
124.36	134.03	Unaltered tuff, fg, hanlike to 0.5mm lamination, dark green to black green and greyish white silica, hard, non carbonate, silicified locally greyish green, bedding well level CA-65o 124.82, trace to nil sulphides, void of stringers, 124.36-124.78 contorted and low angle bedding CA-35-55o, scattered fg py, 126.07-126.54 <0.5mm carbonated phenocrysts poor to weak foliation/schist, 126.54-126.69 pale buff to light brown altn of tuff several 0.5cm qtz stringer, altered zone 1% fg py cotact CA-80+85o, 128.20-128.40 minor small kinkle folding, 128.94 1cm white with pale brown tourmaline CA-75o, 130.54-130.58 qc veinlet CA-70o ll to bed, 132.0-132.16 silicified and qtz vein zone, small stringer 1mm bands of py CA-80o, fragments sericitic altn, 132.16 contact of vein CA-15o, 134.03 contact sharp CA-75-80o.								
134.03	143.63	Silicified tuff to fragmental tuff, similar to above tuff but more qtz and qc random sections 1-2mm, well level bedding CA-65-70o, massive uniform, scattered vfg to fg pyrite locally up to 1% overall <0.5%, rare qc stringers always ll to bed 135.20-135.25, 136.24 0.7cm, 138.08 5mm, 139.08 5mm, 140.42 1cm, 141.79 5mm, 141.80 1.2cm, 142.04-142.44 silicified with 7 qtz veinlets 1/2-1 1/2cm buff altn with minor chocolate brown tourmaline, scattered pyrite in tuff but higher % beside veinlets locally 1-2% fg to mg pyrite, 143.10-143.14 slightly bright green altn almost fuchsite, 143.63 contact CA-65o.	44526	141.5	142	0.5	0.07			
			44527	142	142.5	0.5	0.02			
			44528	142.5	143.6	1.1	0.05	0.06		
143.63	144.52	Qtz vein system and Bx zone, siliceous aphanitic grey massive uniform void	44529	143.6	144.52	0.92	0			

Property: Hunter Mine		Hole No.	29		Sheet No.	2				
Meterage		Description	Sample				Assays		Assays	
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
123.32	303.24	Exhalitic Sedimentary Rocks and Tuffs								
123.32	123.6	chocolate brown, siliceous felsic like x-cut by grey opaque fol., to 5mm, usually at CA-35o, 30o, 20o, 123.67 CA-35o.								
		contorted tuff, tuff fragmental schistose bedding II to CA to 60o-70o, contorted bedding, kink folding to slips planes at axis of folds CA-10o-20o.								
		silicified zone, tuff, dark grey, CA-50o +70o scattered < 0.5% fg py.								
131.34	132.9	blackish green laminated tuff, silicified, qtz II to bedding, vein contorted, moderately soft.								
132.9	134.15	laminated fg tuff, moderately hard, bed CA-65o 134.0, minor kink folding 134.15 CA-48o-50o contact.								
134.15	136.87	very contorted, s+z folds to slips planes II to fold axis CA 20o + 30o intensely qtz stringer flooding, moderately soft, scattered pyrite elongated vfg blobs II to bedding 136.87 contorted CA-65o								
136.87	137.13	fg well developed bed, fine laminated stretched qtz fragments, light med grey greenish tint.								
137.13	138.91	similar to 134.15-136.87 without contorted folding, black green, moderately soft scattered fg py. 137.90-138.20 minor grinding, light brownish tint. 138.57-138.91 silicified crushed zone bx 138.91 contact CA-65o								
138.91	139.53	siliceous fragment to bx zone in black green and medium brown altn silic flooding 139.53 contact CA-70o to 7mm q st displaced by 1 mm qv CA-25o.	44556	138.5	139.5	1	0.03			
139.91	140.5	similar to 137.13-138.91 minor folding, moderately soft, black green local qtz flooding, moderately soft, trace sulphides, 140.50 sharp contact CA-55o 140.50-140.63 silicified contact altn, hard, 140.63 CA-60o small angular x-cut folding.	44557	139.5	140.5	0.99	0.02			
140.63	141.54	aphanitic light grey, opaque qtz vein hard, inclusions of altn tuff, bleached greenish buff to buff tan 1-2mm qtz II CA inclusions CA-30o-35o nil to scattered fg py, 141.54 irregular contact CA 65o-70o gradational.	44558	140.5	141.7	1.16	0			
141.54	142.38	silicified contact altn irregular qtz ff scattered py, blackish green with light brown to medium brown siliceous intrusions irregular qv, 142.38 contact CA	44559	141.7	142.3	0.64	0.01			
142.38	143.2	foliated light grey felsic dike chlorite CA 65o-70o intruded lg silic light brownish felsic dike massive aphanitic silicified (felsic dike?) CA silicified zone?	44560	142.3	143.2	0.87	0.01			
143.2	146.25	medium to dark green locally black green, moderately soft, good devel. bedding fragmental tuff to scattered fragments, qtz flooding usually II to bed CA 65o	44561	143.2	144.3	1.11	0			
		143.50 up to 144.30, 14430-146.25 x-cut bedding folded, usually light grey	44562	144.3	145.5	1.19	0			
			44563	145.5	145.9	0.39	0			

Property: Hunter Mine		Hole No.	29		Sheet No.	3				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		opaque .5-1.5 ore irregular + contorted bedding, 145.48-145.58 whitish qtz translucent to white milky carbonated, inclusions, minor talcose altn, light green to brown tint contacts CA 70o-35o 145.58-145.84 veined silicified fragmental -	44564	145.9	146.8	0.87	0.03			
143.2	146.25	tuff 145.84 irregular contact.								
146.25	147.37	qtz flood fragmental tuff similar to 143.20-146.25 but moderately hard, scattered large pale greyish green to pale greenish grey several cm, 147.37 contact CA-60o.	44565	146.75	147.37	0.62	0			
147.37	147.94	tuff greyish green well laminated x-cut by black green chlorite qtz 2-4mm CA 20o,25o,30o,40o moderately hard, 147.94 contact to bedding CA 80o.	43201	147.37	148.4	1.03	0.01			
147.94	149	light to medium grey, fg, laminated, well devel. bed minor 1-2mm chlorite ll x-cut at bed, bed qtz flooding and altn to pale brownish grey, CA-50o contact	43202	148.4	149	0.6	0.02			
149	149.14	same as 147.94-149.00 intensely silicified contact altn, chlorite and dark brown altn 149.14 contact always irregular CA-58o ll to bedding.	43203	149	149.5	0.5	0.11			
149.14	149.35	light greyish to medium grey siliceous aphanitic felsic dike, very hard, non magnetic, void of stringers, vfg to fg pyrite <0.5% 149.35 sharp contact CA-45o, 149.35-149.48 contact CA-45o								
149.48	150.05	mafic dike, vfg to aphanitic, dark grey, void of foliation, massive uniform, hard to very hard, non magnetic, non carbonated few scattered 1mm qtz ll at CA-60o,80o,70o to 2-4 mm new lower contact, void of sulphides 150.05 contact 50o.	43204	149.5	150.22	0.72	0.01			
150.05	153.85	fragmental tuff contorted, fragmental tuff, moderately hard, laminated light dark grey faint pale brown altn, chlorite sections of tuff, small fragments usually stretched well devel. bedding usually contorted, 150.05-152.20 contorted.	44566	150.22	151	0.78	0.04			
		150.21-150.58 qv brecciated tuff very hard, qtz flooded, contacts CA-50o,40o new ll to bed 153.90 contact CA-40o.	44567	151	152	1	0.01			
			44568	152	153.13	1.13	0			
			44569	153.13	153.67	0.54	0			
			44570	153.67	154.15	0.48	0			
154.15	155.5	minor fragmental tuff, well devel. bed CA-60o light to medium grey, chlorite laminated chloritic bands, hard to moderately hard, non carbonated siliceous to silicified, rare veining 0-2 per 3m scattered to localized fg py >0.5%.	44571	154.15	154.9	0.75	0			
			44572	154.9	155.51	0.61	0			
155.5	156.63	fragmental tuff medium grey - medium green laminated minor tuff, kink greyish white qv at 156.32 5-7mm 156.42-156.63 sericitic altn minor fuchite 156.42 contact CA-62 slips, 156.63 contact CA-65 slips, 156.63-158.35	44573	155.51	155.63	0.12	0.17			
		light grey + and light medium green laminated fragmental tuff, local tuff altn, scattered py chlorite altn, bedding variable 50o-72o, qtz flooding moderate	43205	156.63	158	1.37	0.05			
			43206	158	159	1	0.11			
			43207	159	160.38	1.38	0.07			

Property: Hunter Mine		Hole No.	29		Sheet No.	4				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		158.35-159.39, 157.49 3 cm opaque white qv CA-35o x-cut bed 158.35-160.39 small fragments locally fuchite altn, 159.12-159.28 v wedge white opaque qv with 1 cm chlorite v on lower contact CA irregular x-cut bedding -								
155.5	156.63	locally <CA35o II to bed, locally qtz flood, large fragments, minor kink folding 160.39 contact bed CA-50.	44574	160.39	161.14	0.75	0.03			
160.39	169.36	sericitic tuff, pervasive uniform, pale yellow green, weak to moderate sericitia altn, minor grey 1-2mm qtz eyes minor qtz st usually II to bed, in altn and x-cut in less altn chlorite tuff 162-162.96 locally scattered fg pyrite locally small stretch blobs II to bed, locally cg at 161.35-161.67, rare stringers bed 159.5 CA-60o, 166 bed CA-60o 167.40-167.56 kink folding slip at CA-28o x-cut bedding CA-70o-75o, 167.9 bedding CA-65o, 169.17 1cm grey white qv CA-63o II to bed.	44575	161.14	161.94	0.8	0			
			44576	161.94	163	1.06	0			
			44577	163	164	1	0			
			44578	164	165	1	0.01	0.01		
			44579	165	166	1	0			
			43208	166	167	1	0.04			
			43209	167	168	1	0.02			
169.36	176.87	weak to moderate patchy sericitic altn tuff kink green to medium greenish grey to pale tuff green well level bedding sections of fragmental tuff, 171-171.70 scattered 1cm greyish white qv usually II to bed CA-55o-57o, 171.91-171.98 milky white and grey qv in tuff inclusions CA- Irregular x-cut bed CA-80o + II to bed CA-63o, 172.43-172.62 qtz vein with tuff inclusions CA-57o+75o, 172.77-172.99 qtz vein with tuff inclusions CA-80o+60o, 173.07-173.09 qtz greyish CA-40o x-cut bed 70o, 175.35-176.87 kink folding.	43210	168	169.35	1.35	0.07			
			43211	169.35	170.16	0.81	0.01			
			43212	170.16	171	0.84	0.11			
			44580	171	171.91	0.91	0			
			44581	171.91	173.12	1.21	0.01			
			44582	173.12	174	88	0.03			
			43213	174	175.5	1.5	0.01			
176.87	181.82	fragmental tuff, sericitic weak to moderate pervasive uniform as above, buff to pale buff greenish yellow, 176.87-177.51 scattered greyish qtz eyes fragments, 177.51-177.73 brecciated grey white qtz healed, 178 bedding CA 60o well developed, 178.49-179.71 qtz flooded locally brecciated scattered py II to bed, 179.39-179.45 white qv locally CA-65 x-cut folded bedding at CA 20o 179.94 bedding CA-60o 180.94 1cm white qv CA-35o x-cut bed CA-50o 181.10-181.64 kink folding small, 181.61 fuchsite fragment, 181.64 ground contact.	43214	175.5	176.86	1.36	0.04			
			44583	176.86	177.74	0.88	0			
			44584	177.74	178.5	0.76	0			
			44585	178.5	179.14	0.64	0			
			44586	179.14	179.72	0.58	0			
			44587	179.72	180.66	0.94	0.01			
			43215	180.66	180.83	0.17	0.03			
			43216	180.83	181.46	0.63	0.02			
181.82	182.22	nil to weak sericitic altn medium grey green fg laminate tuff elongated oval qtz II to bed, 182.22 bed contact CA-65o	43217	181.66	183.5	1.84	0			
			43218	183.5	184.6	1.1	0.18			
182.42	194	weak moderate pervasive sericitic tuff 182.42-184.59 qtz flooding, strongly pervasive sericitic altn, py, 184.59-185.92 weak and patch sericitic altn, sile scattered py, qtz flooded irregular + II to bed, 185.92-189.27 weak to weak moderate sericitic tuff minor locally strong qtz flooding	43219	184.6	186	1.4	0.03			
			43220	186	187.5	1.5	0			
			43221	187.5	189	1.5	0.01			
			43222	189	190.5	1.5	0.01			

Property: Hunter Mine		Hole No.	29		Sheet No.	5				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		scattered fg py, 186.34-186.40 white opaque qtz vein CA-60o ll to bed,	43223	190.5	192	1.5	0			
		186.66-186.79 qtz veining ll to bed, 189.27-194 moderate sericitic altn tuff,	43224	192	193.5	1.5	0			
		well devel. bedding, 189.64-189.69 qv ll to bed CA-60 + scattered py fg,	43225	193.5	195	1.5	0.01			
182.42	194	190.15-190.19 rounded qtz massive bedding curved at 190.15 and really								
		flat at 190.19, 191.39-191.65 qtz veining at py fg ll to bed, 192.0-192.23	43226	195	196.5	1.5	0			
		crenulation axis CA-25o x-cut bed 70o.	43227	196.5	198	1.5	0.04	0.06		
194	204.08	weak and patchy sericitic altn tuff majority locally small stretched angular to	43228	198	199.5	1.5	0.01			
		carbonated angular fragments, blackish green to pale yellow tuff, very well	43229	199.5	201	1.5	0.02			
		level bedding CA-75o consistant, void of stringers and qtz flooding, patchy	43230	201	202.5	1.5	0.01			
		locally fg py <0.5% hard siliceous.	43231	202.5	204.08	1.58	0.05			
204.08	204.97	medium grey fg tuff, laminated void of stringer, scattered py <1% + ll to	44588	204.08	204.97	0.89	0.07			
		bedding 204.97 contact CA-70o.								
204.97	205.56	weak patchy sericitic tuff.	43232	204.97	206	1.03	0.03			
205.56	211.34	fragmented tuff, rare patches of weak sericitic altn for 1m, fg matrix medium	43233	206	207.34	1.34	0.01			
		grey to dark grey, more fragments, 207.33-207.74 qtz breccia zone, milky	44589	207.34	207.74	0.4	0			
		white, nil to trace py, contact CA-60o irregular, 207.94-209.84 fragment	43234	207.74	208.82	1.08	0			
		tuff, light kink to pale brownish green, 208.48-208.77 chlorite slips, CA	43235	208.82	209.84	1.02	0			
		40o x-cut bed, 209.84-210.78 qtz flooding, 210.36-210.78 brecciated qtz	43236	209.84	210.78	0.94	0			
		healed zone, 210.78-211.34 hard to moderate hard, siliceous, medium	43237	210.78	211.94	1.16	0			
		brownish green, tuff fragments.								
211.34	217.87	Ultra mafic fragmental tuff, fg, black green light buff green to grey green,	43238	211.94	213	1.06	0			
		211.34-211.94, moderate hard, overall moderately soft to moderately hard,	43239	213	214.5	1.5	0			
		weakly carbonated, good devel. laminae, locally silicified bed to 2-5mm qtz	43240	214.5	216	1.5	0.02			
		stringer ll to bed well devel'ed schistosity, CA-65o-67o, nil to trace sulphides	43241	216	217.5	1.5	0			
		217.56-217.87 qtz veining ll to contorted bedding 217.87 sharp contact 80o	43242	217.5	217.85	0.35	0			
217.87	303.24	Sedimentary Sequence								
217.87	231.36	carbonated argillite vfg fg, medium grey to dark grey few blackish argillite	43243	217.85	219	1.15	0			
		bands, calcareous non magnetic carbonated, hard sections, massive with	43244	219	220	1	0	0		
		little signs of bedding, scattered py in sections vfg and ll to bedding, 219.96-	43245	220	221	1	0			
		220.15 several qtz veinlets ll to bed .5-1.5cm, 218.53-218.74 several qtz	43246	221	222	1	0			
		veinlets ll to bed, 223.92-224.89 scattered 1-2% fg py locally 3-5% ll to bed	43247	222	223	1	0.01			
		CA-70o 224.89- rare to nil veining, 231.23-231.36 black aphanitic to fg	43248	223	223.94	0.94	0			
		graphitic argillite.	44590	223.92	224.89	0.97	0.29	0.22		

Property: Hunter Mine		Hole No.	30			Sheet No.	2			
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		light medium green, talcose contact CA-25 LC crenulated 1-3mm qc stringers.								
107.08	139.37	Laminated Ultramafic Tuffs								
107.08	117.63	foliated UM tuff with minor tuffaceous fragmented, schistose CA to II to CA-38o at 108.5, 58o at 114.7, scattered carbonated stringers underly orientated.								
117.63	118.97	contorted qc stringers 2-7mm with chocolate brown tourmaline.	44569	117.6	119.0	1.34	0.03			
118.97	128.73	tuff - tuff fragmental UM, scattered qc and carbonated stringers 10-15m, schistose CA-28 to II to CA, 122.84 tight fold, axis II to CA, 123.80-126 crumulations to chlorite slip planes CA-15o schistose random & folded, 126.1 schistose 50o, 126.48-126.84 massive porphyritic & foliated CA-45o-15o, 126.84 contact CA-42o, 126.48 contact CA-49o sharp to 4cm grey qv, 126.84-127.26 moderately hard, 127.26-127.72 schistose tuff bed CA-30o minor small folds, 127.72-128.73 very contorted folded, crenulations chlorite II slip planes CA-20o-30o tuff.								
128.73	129.32	massive, foliated, mafic dike? Contacts sharp CA-28o with 1cm qc st porphyritic foliated to 128.97, hard massive CA-58o, 129.32 contacts 70o+50o.								
129.32	138.13	contorted tuff to fragmented tuff, 129.32-131.28 schistose II to CA, 130.37 fold nose, 131.28-136.89 very contorted, CA-40o at 131.9, 35o at 134, 135-136.89 crumulated, folded, chlorite II slips planes (136.89-137.52), 138.13 sharp contact CA-75o-78o.	44597	137.0	138.1	1.13	0.01			
138.13	138.49	mafic dike, fg, dark green altn at contacts to pale green buff, massive, uniform, nil level of foliation, non magnetic, non carbonated, siliceous, x-cut by random 1-2 mm white qtz ff stringers, nil to trace sulphides, LC ground.	44598	138.1	138.5	0.36	0			
138.49	139.37	carbonated schistose UM, as above well level of schist CA-60o trace sulphides minor crenulations to chlorite slip, 139.37 sharp contact CA-55o.	44599	138.4	139.4	0.93	0.06			
139.37	268	Exhalitic Tuffs and Sedimentary Rocks								
139.37	140.15	felsic dike, fg aphanitic, light brown to buff brown, hard, siliceous, nil level of foliation, brownish siliceous ff (tourmaline) and random q st 1-2mm, scattered fg py <0.5%, 139.82 5mm white qv CA-80o, 140.05 5mm white qv CA-70o-75o, 140.15 ground contact.	44600	139.4	14.2	0.78	0.01			
140.15	141.55	foliated felsic dike, aphanitic to fg buff to pale greyish buff to black green chlorite elongated phenocrysts to foliated bands 1mm, hard, siliceous few 1-2 mm q st usually near II to CA, few at 25o +55o foliation, trace to scattered fg to vfg py <0.5%, 141.05-141.16 1/2cm and wedge shaped 2-6cm	44601	140.2	141.6	1.4	0.04	0.04		

Property:		Hole No.	30		Sheet No.	3				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		chocolate brown siliceous veins ie brecciated felsic dike fragments, 141.55 contact CA-50o.								
141.55	142.06	um tuff fragmented, as above, 141.55-141.88 buff altn fragments white dark green, matrix, silicified, 141.88-142.06 fg grained weak level of bedding or schist, hard medium brownish green to pale green buff, possible felsic dike weak foliated, contacts sharp 60o+60o.	44602	141.6	142.6	1.03	0.02			
142.06	144.6	qtz flood wk altn, ultra mafic tuff fragments, flood II to schist CA-45-50o to crenulated, minor displacements along axis, grey white with pale brown tint, trace to scattered vfg fg py, 142.18-142.41 irregular white qtz vein mass.	44603 44604	142.58 143.6	143.6 144.6	1.02 1	0 0			
144.6	145.66	medium grey to medium grey green, um tuff 1-2mm laminated, patchy trace of sericitic altn.	44605	144.6	145.66	1.06	0			
145.66	146	aphanitic to fg, black green to dark grey laminations tuff to siliceous seds, scatter to <0.5% pyrite, pyrrhotite II to bed CA-50o in more chlorite sections 145.92 to 146.0.	44608	145.66	146.1	0.44	0.01			
146	146.1	massive chlorite with irregular pale grey white opaque qv, scattered py po and chalcopyrite, ground contact.								
146.1	258	Exhalitic Sedimentary Rocks and Tuffs								
146.1	147.41	fg grey green laminated + crenulated tuff grading to chlorite orthi fragmental 146.95-147.41, few 3-5mm grey q st x-cut bed, 146.42-146.68 crenulation axis CA-15o chlorite II, 147.05 bed CA-57o, 147.14-147.41 several 5mm qtz translucent qtz to minor carbonate II to + x-cut bedding, 147.41 contact CA 35o.	44607	146.1	147.41	1.31	1.51	1.49		
147.41	148.64	aphanitic medium grey, very hard, non magnetic, no carbonate, siliceous, massive uniform, ie grey white opaque qtz st near II to bedding CA -55o, scattered 1-2% fg py usually in qtz ff, 148.64 contact CA-60o.	44608	147.41	148.64	1.23	0			
148.6	149.5	light brownish green to pale greenish light brown, baked altn by greenish brown <1mm black phenocrysts dike form 148.88-149.12 on side of core + x-cut bedding, 149.12-149.5 fragments altn to buff to greenish tone, 149.24-149.50 qtz flooding & irregular whitish opaque qv, 149.50 contact CA -75o.	44609	148.64	149.5	0.86	0			
149.5	150.23	crenulated locally, tuff, light grey to buff pale green, trace to scattered vfg py, 150.23 siliceous contact CA-60o	44610	149.5	150.23	0.73	0.01			
150.23	151.64	brecciated qtz flooded fragmental tuff, altn hard silicified zone, light grey, pale grey, buff, tan, dark coloured fragments, locally chocolate brown silicified sections, with irregular patches + masses of white qv, bedding x-cut by	44611	150.23	151.64	1.41	0.1			

Hunter Mine - Diamond Drill Log HM-04-31

Property:	Hunter Mine	Hole Dip:	-50	Page No.:	1 of 9	Hole HM-04-31				
Location:		Hole Azimuth:	105	Date Started:	June 28,2004					
Claim No:	HR 1009	Hole Length:	242.15m	Date Finished:	02-Jul-04					
Elevation:	Porcupine Lake	Purpose:		Drill Co.:	Benoit					
UTM Coords.:	5370891.3N, 487116.9E			Logged by:	K. Jensen					
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
0.0	64.3	Casing								
64.3	82.7	Ultramafics								
64.3	68.6	very talcose, very soft, nil to very weak magnetic, fg, light to medium grey to blackish grey blue hue, nil to very weak carbonated, massive uniform in qts stringer, locally breccia chlorite healed, scattered mg to cg pyrite + masses, 68.45 to 68.60 crushed crumbly core, shear fault zone.								
68.6	72	massive locally brecciated, rare veining.								
72	73.56	porphyritic texture, moderately soft, UC broken LC 20-25o, 1/2 to 1cm pyrite masses + x-cut, few irregular qc stringers random CA-20o to 45o.	44644	72	73.5	1.5	0			
73.56	75.68	possible flow top breccia, minor random qc veining LC 35o-40o, 74.95-75.06 irregular qtz masses.	44645	73.5	75	1.5	0			
75.68	76.54	massive, crushed zone, 76.44-76.54 qc zone of distortion CA-80o+65o LC sharp.								
76.54	80.81	tuffaceous to tuff fragments, moderate well develop bedding bedding CA-45o local x-cut chlorite ff CA-20o, locally crenulated, scattered qc il to bed, trace to <0.5% pyrite fg mg, 80.50 5cm qc irregular vein mass in black tournalline x-cut CA-50o. 80.81 shearing CA-70o-80o.								
80.81	82.65	pale grey green black, very soft, talcose, moderately carbonated, crushed zone, weak level of sch to nil, CA-55o LC.								
82.65	112.42	Laminated Ultramafic Tuffs								
82.65	83.5	tuff fragmental.								
83.5	86.61	brecciated massive, LC gradational 85.43-85.64 crushed zone.								
86.61	87.53	tuffaceous, minor fragments LC-60o.								
87.53	88.03	massive intrusive UM dike, non carbonated kinkle qtz ff stringers, nil to trace sulphides, black fg, moderately soft, non magnetic, LC sharp CA-40o x-cut bedding CA-60o.								
88.03	98	moderate to level bedding tuff to tuff fragmental moderately soft more carbonated to moderately hard more siliceous sections, minor crenulations, scattered fg mg pyrite to trace, 93.56-93.97 intrusive carbonate ff stringers CA-80o + LC shearing								

Property		Hunter Mine	Hole No.	31		Sheet N	2			
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		ll to bed, 90.15-90.31 greyish qtz vein in chlorite ll, contacts UC 27o x-cut bed CA 60o-65o, LC-55o, 94.60-96.90 blobs + masses of qtz discontinuous stringers,								
88.03	98.00	96.90-98.00 more tuffaceous, bedding locally crenulation 60o.								
98	99.12	massive, moderately soft to moderatly hard, nil to poorly develop. schistosity, minor fg, scattered 1-2mm qtz stringers deformed, 40 60o LC 55o.								
99.12	99.73	tuff.								
99.73	99.81	aphanitic to fg, light grey, felsic dike, level, chlorite ll, trace to vfg py contacts CA- 75 + 55o sharp.								
99.81	100.28	vfg tuff to lapilli ash, medium dark green greyish white fragments, nil to trace sulphides, hard silicified.	44646	99.7	100.3	0.55	0.08			
100.28	100.7	felsic dike, same as 94.73-99.81, scattered to 1% vfg fg pyrite, chlorite ff, 100.40-100.66 regular chlorite ff almost laminated, 100.66-100.70 massive chlorite LC CA-70o in 7mm grey opaque qv.	44647	100.3	100.7	0.42	0.09			
100.7	102.27	tuff, fragmental, well level bed, locally contorted chlorite filled slip planes CA-50o x-cut bed 60o locally crenulated.	44648	100.7	102.3	1.57	0.02			
102.27	103.09	same as 99.81-100.28, foliated medium green, hard, x-cut white 1-2mm stringer qtz, grey light bleaching altn on contacts, trace to 0.5% py, locally 102.66-102.88 2-3% mg py. 102.48-102.55 light grey felsic dike in white qtz ff st only 1 x-cut into + past contact. 102.88-103.09 greyish felsic dike in light brown tint scattered py.	44649	102.3	103.1	0.82	0.05	0.08		
103.09	103.94	qtz stringer contorted bedding tuff laminated UC 50 LC-65o gradational, trace to scattered py.	44650	103.1	103.9	0.85	0.02			
103.94	104.32	greyish felsic dike in chloritic qtz tuff inclusions, 1-2% fg pyrite LC-50o irregular in contact tuff.	44651	103.9	104.3	0.38	0.03			
104.32	113	blackish green grading to light greyish green talcose, silicified, moderately hard to hard, contorted bedding tuff in fragmentals, to locally talcose 111.42-111.94, 108.15-109.08 qtz flooding, random stringers, 108.90-109.08 opaque irregular qv in inclusions, pale green tint, LC CA-75o-80o, nil to scattered fg pyrite locally 1-2%, 112.42 5cm greyish + brownish white qv CA-32o irregular.	44652	108.2	109.1	0.93	0.03			
			44653	112.0	113.0	1	0.41			
113	242.15	Exhalitic Tuffs and Sedimentary Rocks								
113	113.6	aphanitic to fg, buff tan to pale yellowish tint, hard, siliceous, qtz stringer ff, with irregular qtz vein at contacts, 113.00-113.16 qv CA-80o+75o vfg scattered pyrite 113.36-113.50 qtz stringers, inclusion of tuff, fg ng py, 113.50-113.60 qv CA-40	44654	113.0	113.6	0.6	2.61	1.89	2.25	
113.6	115	same as 104.32-113.00 fragmental tuff with white opaque qtz stringers altn gradational contact.	44655	113.6	114.7	1.1	0.29		0.29	
115	115.8	weak sericitic altn, fragmental tuff LC irregular.	44656	114.7	115.8	1.1	2.06		2.06	
115.8	116.12	qv greyish to brownish older qv x by milky white opaque qv UC CA-32o 1cm	44657	115.8	116.3	0.5	36.48	36.82	36.65	

Property: Hunter Mine		Hole No.	31		Sheet N	3			
Meterage		Description	Sample			Assay			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)
		LC CA-45o sinuous near ll to bed.							
116.12	116.3	fragmental sub angular weak sericitic.							
116.3	116.88	chloritic fragmental tuff LC CA-55o veining.	44658	116.3	117.14	0.84	0.05		
116.88	117.14	buff felsic fragmental to brecciated in vfg ground mass, nil to very weak devel. bedding, scattered qtz 1-4mm CA-20o-30o x-cut bedding LC CA-50o.							
117.14	118.18	chloritic to very weak sericitic altn, contorted bedding, scattered qtz 1-4mm CA-20o-30o x-cut bedding LC CA-70o-75o.	44659	117.14	118.18	1.04	0		
118.18	119.56	felsic dike aphanitic to fg, massive uniform, hard, silicious non magnetic, non carbonated, chlorite + qtz ff, random orientation pale buff tan to pale brown, 118.18-118.41 brecciated grey silicia healed minor chlorite tuff inclusion, 118.41-118.83 massive chlorite + qtz ff, 118.83-119.06- 5cm ground + lost core at 119.94 chlorite massive 3cm and fragmental tuff, 119.06-119.56 felsic in weak to moderate foliation of discont. chlorite ll lination CA-55o, UC irregular LC-70o displaced by q st CA-10o-15o sharp.	44660	118.18	119.56	1.38	0.04		
119.56	124.6	aphanitic pale grey to pale brown, in well devel. of foliation CA-40o-65o usually chlorite, hard, non carbonated inclusions, non magnetic, non carbonated, massive uniform, random greyish white qtz ff in altn on contacts 119.56-120.82, med grey, random grey qtz ff + py CA-45o, 119.56-119.60 altn fragmental tuff, 119.60-120.82 med grey, scattered cg py, 120.38-120.47 altn bleached brecciated chlorite qtz healed, 120.59 4mm blackish brown to dark brown vein 35o in buff inclusion, 120.82-120.90 altn bleached breccia to qtz st, 120.90-123.0 pale brown to buff tan + chlorite brecciation, scattered to trace py, rare qtz ff st, 123.0-124.60 light to medium grey+chlorite brecciation few qtz ff st in bleaching on contacts, 124.60 sharp contact CA-75o-80o	44661	119.56	120.9	1.34	0.01	0.02	
			44662	120.9	122	1.1	0.02		
			44663	122	123	1	0.01		
			44664	123	124	1	0.05		
			44665	124	124.6	0.6	0.02		
124.6	127.67	mafic tuff fragmental tuff, 124.60-126.50 weak pervasive sericitic altn tuff, locally deformed bedding in minor chlorite slips planes up to 126.0 CA-30o, few fragments 125.30-125.42, 126.50 contact 5mm grey qtz st CA-70o x-cut tuff but ll to chlorite fragmental tuff, 126.50-127.06 chlorite altn fragmental tuff in qtz ll to bedding 70o alternating light green + dark green, scattered py <0.5%, 127.06-127.13 greyish qtz zone in bleached buff inclusions contacts CA90-70o straight, sharp, 127.13-127.67 aphanitic pale grey, medium grey	44666	124.6	125.6	1	0.01		
			44667	125.6	126.51	0.91	0		
			44668	126.51	127.05	0.54	0.1		
			44669	127.05	127.67	0.62	1.17	0.9	1.035
124.6	127.67	in vfg pyrite overall 1-2% locally 3-5%, LC CA-80o locally buff altn due to silicification.							
127.67	128.98	light to medium green, to buff patchy sericitic altn, fg tuff, random orientated translucent greyish qtz st and masses 127.67-128.78 vfg py 1-2% LC slips CA-60o.	44670	127.67	128.98	1.31	0.54		

Property		Hunter Mine		Hole No.	31		Sheet N	4			
Meterage		Description		Sample			Assay				
From	To			No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
128.98	132.09	weakly altn sericitic and pervasive, fg tuff pale buff to light medium grey green occasional fragment, scattered vfg py usually ll to bed 77o at 129.3 65o		44671	128.98	130	1.02	0.03			
		at 131.25, few scattered whitish to whitish pink qc st 2-3mm x-cut bedding usually defined 40%, 60% ll to near ll to bedding, 130.11 1cm vuggy pick qc st CA-90o x-cut bed, 131.42-131.63 whitish qv to buff inclusions 65o ll bed 75o x-cut bed scattered to trace pyrite vfg fg, locally 1-2%, 132.09 contact of qtz v CA-60o x-cut bed.		44672	130	131	1	0.01			
				44673	131	132.09	1.09	0.01			
132.09	132.66	pale green to pale buff altn silicified qtz flood zone, scattered py fg white opaque and translucent stringers ll to bedding 70o, 40o and greyish black CA-30o x-cut bed, overall 1-2% pyrite, 132.66 CA-72o.		44674	132.09	132.66	0.57	0.02			
132.66	133.72	patchy sericitic altn, tuff few greyish qtz st CA-30o x-cut bed, buff at light green trace sulphides 133.72 sharp contact 90o.		44675	132.66	133.72	1.06	0.02			
133.72	134.62	chloritite white grey qtz aphanitic ash to tuff, void of fragments, possible chloritite argillite? Scattered fg to cg pyrite 1-2% sharp contact 134.62 CA-75o.		44676	133.72	134.62	0.9	0.03			
134.62	137.62	weak pervasive sericitic altn, thin to 1-2mm laminated tuff, pale to medium grey, pale green, well level bedding CA-135-75o, 135.9 CA-70o, few scattered greyish qtz st 5mm ll to bedding and pyrite, 135.08-135.27 qtz flooded, 136.24 136.34 white opaque to pale buff altn inclusion CA-55o x-cut bedding, 136.52-137.03 2-5mm irregular sil. qtz grey stringer near ll to CA 1-2% locally vfg py in tuff, 137.62 contact in 3mm black baked contact CA-70-75o.		44677	134.62	136.12	1.5	0.01			
				44678	136.12	137.62	1.5	0.01			
137.62	138.25	felsic dike massive, uniform, level, siliceous, as above aphanitic light to pale greenish to pale brownish grey with greyish kinkle qtz st and straight grey qtz in white brecciated qtz in middle 7mm CA-50o, 138.25 sharp CA-55o.		44679	137.62	138.25	0.63	0.08	0.09		
138.25	140.67	mafic tuff, fg grey to pale greenish grey laminated tuff to pale grey siliceous bands, locally contorted bedding, sections qtz flooding, hard to very hard, scattered fg py <0.5%, 140.67 contact to qtz vein CA-50o irregular.		44680	138.25	139.45	1.2	0.01			
				44681	139.45	140.67	1.22	0			
140.67	140.78	qv white opaque in chlorite tuff inclusion, 140.78 CA irregular 70-75o nil sulphides.		44682	140.67	141.45	0.78	0.15	0.14		
140.78	142.06	felsic dike, as above, light grey, qtz flooded, sections aphanitic in no visible foliations at bedding, sections in black brown ff kinkle stringers, 140.78-141.45 scattered vfg fg 1-3% pyrite, 141.45-141.95 aphanitic, greyish, silicified + siliceous irregular white opaque random orientated qtz veinlets 2-3cm each, contact sharp 65o+60o, 141.95-142.06 similar to 140.78-141.45 foliated pale chlorite and greyish brown ff, 142.06 sharp CA-50o.		44683	141.45	142.06	0.61	0.11			
142.06	142.62	qtz flood tuff, 142.13-142.35 white opaque qv + pale buff inclusion and chlorite CA-20o, trace sulphide to <0.5%, contorted bedding in white opal qtz stringers with 0.5 to 1mm chocolate brown bands possibly tourmaline or		44684	142.06	142.62	0.56	0.17			

Property:		Description	Hole No.	Sample			Sheet No.			Assay		
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check		
184.28	185.85	patchy sericitic altn tuff, 184.75-184.80 grey white to grain tint qv opaque CA II to bed 75o, 185.17-185.20 grey white qv opaque CA-65-65o, 185.44-185.85 nil to weak patchy crenulated tuff, medium grey to dark grey, laminated CA-65o, 185.85 altn contact II to bed 65o.	44729	184.28	185.85	1.57	0.02					
			44730	185.85	186.33	0.48	0					
186.33	187.33	felsic dike, aphanitic, pale buff to light tan, very hard, siliceous nil to very weak level foliated, scattered disseminated vfg to fg pyrite 1-2% up to 3% locally, 1-2mm grey discont. qtz stringers random, very weak carbonated, 186.62-186.77 grey qv opaque CA-75+70o, 187.33 LC CA-65o.	44731	186.33	187.33	1	0.03	0.03				
187.33	188.47	fragmental tuff felsic, aphanitic pale buff to light tan fragments elongated in aphanitic, med grey matrix of tuff + greyish white siliceous bands, scattered pyrite 0.5-1%, well develop bed, 187.5 CA-55o+70o, 188.0 CA-38, 188.2 CA-60o, 188.47 CA-60o.	44732	187.33	188.47	1.14	0.02					
188.47	189.96	weak moderate sericitic tuff, same as above, locally crenulated, scattered py, weak moderate pervasive sericitic altn.	44733	188.47	189.96	1.49	0					
189.96	195.94	fragmental tuff, nil to weak patchy sericitic altn <10-20% local crenulation, blackish tuff bands, buff to light tan to light brown, scattered fragments, scattered 0.5-1% vfg fg py, 2 qtz stringer 2-3 mm at 19.26+194.43 ivory 20+	44734	189.96	191.46	1.5	0.01					
			44735	191.46	193	1.54	0					
			44736	193	194.5	1.5	0.04	0.06				
189.96	195.94	400 x-cut bed, 189.96-191.95 crenulated tuff 193.25 small fragments, 193.34 195.69 qtz flood tuff to fragmental tuff, 195.69-195.94 fragmental tuff + up to 3-5% fg py, 195.94 contact sharp CA-75.	44737	194.5	195.94	1.44	0.13	0.1				
195.94	196.12	grey felsic dike, aphanitic, light grey, massive, uniform, hard to very hard, dike, II to bedding contacts 75o+60o, nil sulphides to trace.	44738	195.94	196.72	0.78	0.09					
196.12	202.67	fragmental tuff to tuffaceous fragmental, fg, light to medium grey, silicified, to siliceous fragments tuff, moderate level of bedding CA-50-65o, hard, non carbonate, non magnetic, 196.72-196.87 grey opaque qv CA-82+55o 1-2% 2mm patches of fg pyrite, 197.17-197.21 v-shaped brownish pink felsic dike 197.34-197.48 v-shape brownish pink felsic CA-40o irregular in opposite directions, 198.23 CA-60o, 198.23-202.67 fragmental tuff + qtz flood, kinkle green, siliceous, irregular qtz st, non carbonate, 202.67 contact 58o.	44739	196.72	196.87	0.15	0					
			44740	196.87	197.48	0.61	0					
			44741	197.48	198.23	0.75	0					
			44742	198.23	199.5	1.27	0					
			44743	199.5	201	1.5	0					
			44744	201	202.5	1.5	0.01					
202.67	202.98	porphyritic in black x-cut, weak level of foliation, LC in irregular 1cm qst x-cut foliation CA 60 and next unit.	44745	202.5	203.1	0.6	0					
202.98	205.6	qtz flooded fragmental tuff, grey green, bedding near II to CA up to 204.63, 204.63-205.25 CA-30o-50o local hole, + 1% py 205.25-205.60 near II to CA 203.10-203.42 qv greyish pink white 30o at uc 1.5cm contacts LC 55o-65o grey breccia qv to 203.48.	44746	203.1	203.48	0.38	0					
			44747	203.48	204.63	1.15	0					
			44748	204.63	205.6	0.97	0					
205.6	206.46	minor qtz flooded, tuff in crenulations.	44749	205.6	206.46	0.86	0					
206.46	207	fragmental tuff in minor qtz flooding 68o.	44750	206.46	207.5	1.04	0					

Property: Hunter Mine		Hole No.	32		Sheet No.	2				
Meterage		Description	Sample			Assays				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
125.59	127.22	crenulated tuff very siliceous fragmental to chlorite ll axis of crenulations, blackish to medium grey to dark grey, qtz stringer ll to bedding.	44778	129.0	130.5	1.5	0.05			
127.22	133.6	fragmental tuff, scattered <0.5% fg pyrite locally cg cubes, moderately soft to moderately hard, blackish fg, gradational contact.	44779	130.5	132.0	1.5	0			
			44780	132.0	133.5	1.5	0.02			
133.6	136.94	tuff, laminated very tight S folds kinkle folding, rare to 1 stringer/meter, scattered to trace pyrite, moderate soft, very weakly carbonated, moderately hard silicified 136.81-136.94 UC 80o, 136.94 contact CA-85o.	44781	133.5	135.0	1.5	0			
			44782	135.0	136.0	1	0.01			
			44783	136.0	136.9	0.94	0.03			
136.94	137.69	qtz vein, fg, blackish grey, hard, silicified fragmental tuff, blackish and light brown, 3-5% fg py few ng, 134.44-137.53 inclusion of carbonate TCS fg tuff, 137.53-137.69 light grey to pale brown qv in inclusions, 137.69 contact CA-85o.	44784	136.9	137.7	0.75	0			
137.69	138.15	fragmental tuff as above, contorted folded bedding, trace sulphides 138.15 CA-50	44785	137.7	138.2	0.46	0			
138.15	138.41	qtz vein, medium brown, minor inclusions <1% grading to 3-5% vfg, 138.41 CA-60o slightly sinuous.	44786	138.2	138.4	0.26	0			
138.41	142.64	tuff, fg contorted shapes S folds tight, numerous chlorite slip planes CA-20-30o grading to laminated tuff form 140.57-140.73, 139.45- 139.65 1/2 cm brownish light felsic bands contorted to match bedding to ll to CA, 140.73-142.64 slightly to weakly contorted bedding minor fragments, very siliceous, laminated, 142.64 contact CA-35o.	44787	138.4	139.0	0.59	0			
			44788	139.0	140.5	1.5	0			
			44789	140.5	142.0	1.5	0			
			44790	142.0	142.6	0.64	2.32	2.59		
142.64	143.08	qv, 142.64-142.75 pale brown, opaque, minor chocolate brown tourmaline, 142.75-142.81 medium brown altn tuff, 142.81-143.02 white opaque qv inclusions white carbonate, 143.02 contact CA-60o, 143.02-143.08 pale greenish yellow bleached felsic dike silicified, vfg py <0.5 to trace 143.08 CA-70o x-cut bed 55o	44791	142.6	143.1	0.44	0.2			
143.08	143.87	fg, grey grain, silicified, qtz flooded, trace py, 143.87 CA 60o.	44792	143.1	143.9	0.8	0.51	0.46		
143.87	146.5	felsic tuffaceous fragmental, pale buff to tan in kinkle green chlorite laminated locally buff to pale buff fragments, trace py, siliceous, 143.87-144.51 felsic tuff buff fragments, 144.51-144.59 white opaque to translucent qv CA-30o irregular 144.59-144.87 felsic tuff + fragments, 144.87-145.20 felsic tuff in kinkle chlorite lamination, 145.20-145.84 felsic fragments in chloritic matrix, contorted bedding S folding LC 60-65o, 145.84-146.50 felsic tuff with chlorite laminations of small kinkle grading to healed chlorite lamination LC CA-50o sinuous.	44793	143.9	144.9	1	0			
			44794	144.9	145.9	0.97	0			
			44795	145.9	146.5	0.65	0.13			
146.5	146.88	altn tuff, altn ultramafic chlorite tuff fragmental, light buff to tan medium green 3 white opalesant qtz st 1cm CA-50o.	44796	146.5	147.0	0.5	0			
146.88	147	tuff fragmental, as above, grey green LC CA-35-40o.								
147	147.55	qv, pale greyish to pale brownish white opaque, nil to trace py, LC CA-80o.	44797	147.0	147.6	0.55	0			
147.55	151.18	silicified altn buff green to light brown, scattered py vfg, 151.18 CA-80-85o.								
151.18	151.88	qv, white opaque with pale brown patches associated in altn tuff inclusions and chocolate brown ll near LC, trace sulphides LC CA-65o.	44801	151.18	151.88	0.7	0.19			

Property		Hunter Mine		Hole No.	32		Sheet No.	3			
Meterage		Description		Sample		Assay					
From	To			No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
151.88	152.13	fragmental tuff to tuff, hard to moderately hard, same as above, 151.88-152.13 altn silicified, pale brownish green x-cut by 1cm q st white CA-50o 158.24 at 152.04.		44802	151.88	153	1.12	0			
152.13	153.62	light grey green tuff trace py, rare 1/2cm grey qtz st LC 70o distinct.		44803	153	153.61	0.61	0			
153.62	156.93	blackish to black green um tuff with qtz lamination, locally contorted folding bedding, moderately hard, scattered 0.5% py fg locally up to 1-2% as 5mm blobs 156.0-156.40, 156.74-156.93 moderately soft, 156.93 LC 70o sinuous.		44804	153.61	155	1.39	0			
				44805	155	156	1	0			
				44806	156	156.93	0.93	0			
156.93	158.24	massive dike mafic, fg, light to medium grey, moderately hard, non carbonate non magnetic, massive, uniform, x-cut by numerous 1-3mm qtz stringers scattered disseminated fg to mg occasional blob of pyrite overall <1%, sharp CA-65o near II to bedding.		44807	156.93	158.24	1.31	0			
158.24	158.39	chlorite tuff as above, moderately soft medium greenish grey, gradational contact.		44808	158.24	158.95	0.71	0			
158.39	158.95	qtz stockwork, grey green tuff qtz stockwork greyish opaque q st x-cut by white translucent qtz st, random occasionally trace pyrite LC CA-72o.									
158.95	159.36	chlorite tuff, laminated tuff with chlorite fragments, scattered fg py in bands overall 1-2% upper part altn to light medium grey, bed CA-80o, 159.36 CA-75		44809	158.95	159.36	0.41	1.06	0.98		
159.36	159.79	siliceous zone, fg light grey to pale grey, hard siliceous to silicified, no distinct veining, fragmental to brecciated, LC CA-75o II to bed tuff chlorite trace pyrite.		44810	159.36	159.79	0.43	0.52	0.54		
159.79	160.23	chlorite tuff, same as 158.95-159.36, trace pyrite LC-60o sharp vein.		44811	159.79	160.23	0.44	0.02			
160.23	161.3	fg, hard to very hard, siliceous, pale grey to grey creamy opaque, with silicified tuff inclusions or fragments. 160.36-160.40 brecciated sections with fragments in medium brown siliceous matrix CA-53o slips, 160.69-160.77 silicified tuff fragments inclusions, few scattered grey white 1-3mm stringers low angular to II to CA, trace py to nil, 161.30 sharp CA-70o.		44812	160.23	161.3	1.07	0			
161.3	162.48	silicified fragmental, extremely silicified, nil to very poor pervasive bedding light to medium grey altn, void of stringers, non carbonate trace to <0.5 fg py, 162.48 sharp CA-80o-85o.		44813	161.3	162.48	1.18	0.02			
162.48	162.62	vein felsic dike, light grey, extremely fractured dark brown folding, hard, siliceous, non carbonate, trace sulphides LC CA-55o		44814	162.48	163.35	0.87	0.01			
162.62	163.23	similar to 161.30-162.48 shade greener, silicified, 163.23 chloritic inclusion of tuff fragmental LC irregular.									
163.23	163.35	vein qtz, pale grey opaque irregular center 40o sinuous to py.									
163.35	165.1	fragmental tuff, same as above, light grey green, local tuffaceous sections, small crenulation, few greyish qtz stringers 2-5mm near II to bedding + x-cut bedding, nil to trace pyrite, 164.54-164.87 grey fg felsic dikelet CA-75-70o		44815	163.35	165.1	1.75	0.01			

Property		Hunter Mine	Hole No.	32		Sheet No.	4			
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		165.10 CA-70-75o.								
165.1	166.28	fragmental brecciated healed with chocolate brown siliceous to 165.54 CA-70 with 3cm white opaque qv at 165.26 CA-42o and x-cut grey white 0.5-1cm qtz string CA 5-7o terminated or ends at 165.54, 165.54-166.19 brecciated tuff + fragments healed with grey translucent qtz and 1mm qtz grey CA 15o in opposite, 166.19-166.28 irregular white qtz st 70o with inclusions, trace pyrite.	44816	165.1	166.28	1.18	0.01			
166.28	167.21	tuff fragmental, as above, buff and medium green laminations, buff to tan fragments, trace py, void of qtz stringers, well level bed 70o, 167.21 slips contact CA-70o.	44817	166.28	167.21	0.93	0			
167.21	167.57	tuff fragmental altn, fg grey silicified tuff fragments, very hard, silicified, 1% scattered ff to vfg py, 167.96 sharp CA-65-70o.	44818	167.21	167.57	0.36	0.17	0.18		
167.57	167.96	qtz stockwork veining, fragmental buff healed with chocolate brown siliceous x-cut bg dominately greyish white translucent qtz st and a few white qtz st CA 40o up to 167.68, scattered 0.5-1% vfg py, 167.96 sharp CA-65-70o.	44819	167.57	167.96	0.39	0.01			
167.96	172.64	fragmental tuff, 167.96-168.25 light greyish buff to greyish buff, no stringers, gradation, 168.25-168.59 pale grey to light grey, siliceous disseminated 1-2% py 6mm cluster of vfg, 168.59-169.10 pale grey to pale greenish tint grey, more tuffaceous, few fuchsite patches, trace to scattered <0.5% py, LC sharp 60o, 169.10-169.15 grey felsic dikelet or qtz CA-85o LC, 169.15-169.17 tuff as 168.59-169.10, 169.17-169.35 fragmental brecciated healed with chocolate brown silicic matrix + x-cut by grey translucent qtz 2+5mm, CA-40o which is x-cut by 1-1.5 white opaque qtz CA-25o x-cuts LC 70o, 169.35-169.80 greyish	44820	167.96	169.1	1.14	0			
			44821	169.1	169.35	0.25	0			
			44822	169.35	170.73	1.38	0.02			
			44823	170.73	172.12	1.39	0			
167.96	172.64	fragmental tuff, 169.80-170.37 pale buff weak devel. sericitic altn FT, 170.37-172.12 patchy weak sericite FT, bed 80o, 172.12-172.64 moderate pervasive sericitic altn.	44820	167.96	169.1	1.14	0			
			44821	169.1	169.35	0.25	0			
			44822	169.35	170.73	1.38	0.02			
172.64	173.02	weak sericitic patchy <5%, buff to light + medium grey tuff,.	44823	170.73	172.12	1.39	0			
173.02	174.67	pervasive moderate sectional altn, qtz flooding scattered to brecciated vfg py 0.5-1% contacts sharp 70o+70o.	44824	172.12	173.02	0.9	0.01			
			44825	173.02	173.9	0.88	0.05			
174.67	181.02	light grey, medium grey, light grey with greenish tint tuff, well devel. bedding not deformed, fine laminations, crenulation minor at 179.77-179.82. Very rare to nil stringers, minor st qtz fine, 180.48-181.02 broken contact.	44826	173.9	174.67	0.77	0.29			
			44827	174.67	176	1.33	0.01			
			44828	176	177.5	1.5	0			
181.02	181.79	weak to moderate pervasive sericitic altn locally fine laminations of pyrite, 181.02-181.24 rusty brown fixture filling.	44829	177.5	179	1.5	0			
			44830	179	180	1	0			
181.79	185.98	same as 181.02-181.79 qtz flooding few scattered patches of fuchsite, 182.62 grey 1cm gash qtz white, 182.71 ivory 1cm white traces qst st 50o x-cut bed, 182.80 straight 1cm white traces qst st 40o x-cut bed, 183-minor small crenulations, 183.81-183.89 qtz vein with inclusions II to bed 70o	44831	180	181.02	1.02	0			
			44832	181.02	181.79	0.77	0			
			44833	181.79	183	1.21	0.04			
			44834	183	184.5	1.5	0			

Property Hunter Mine		Hole No:	32		Sheet No.	5				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		183.89-185.39 in distinct veining, qtz flood, 185.39-185.98 scattered .5-1cm grey white traces qtz st usually ll to bed LC 40o.	44835	184.5	185.98	1.48	0			
185.98	187.19	brecciated grey + brownish healed x-cut by large 1cm-2.5cm white traces qtz veinlets 40, 20o qtz flooded LC irregular 30-35o	44836	185.98	187.19	1.21	0			
187.19	191.77	fragmental tuff, scattered pyrite 0.5-1%, strongly pervasive sericitic altn, 187.19-191.93 qtz flooded, 190.6 fuchsite, 189.34 2cm black chlorite qtz st CA-55o, local crenulations at 188.74.	44837	187.19	188.5	1.31	0.06			
			44838	188.5	190	1.5	0			
			44839	190	191.5	1.5	0			
191.77	193.33	tuff, minor crenulations, trace to scattered py locally 1%, 191.77-192.40 minor qtz, good bed 80o, 192.40 1/2cm light grey felsic dike x-cut bed 88o	44840	191.5	192.5	1	0			
		192.40-193.33 qtz flooding 30%, 193.09 fuchsite.	44841	192.5	193.33	0.83	0			
			44842	193.33	194.5	1.17	0.05	0.04		
193.33	202.65	tuff, patchy random sericitic altn, buff to medium grey well laminated tuff, qtz ll to bed well level bed CA-75o at 195.3, 80o at 199, patchy to scattered py, 194.40 black qtz chlorite st CA-70o x-cut bed, minor crenulations with moderate grey sections 201.40-201.60.	44843	194.5	196	1.5	0.02			
			44844	196	197.5	1.5	0			
			44845	197.5	199	1.5	0			
			44846	199	200.5	1.5	0.01			
202.65	205.47	patchy weak to moderate sericitic altn, minor crenulations in medium grey sections, rare fragments, void st scattered <0.5% py LC 70o ll to bed.	44847	200.5	201.5	1	0.01			
			44848	201.5	202.65	1.15	0			
205.47	209.29	pervasive moderate sericitic altn, tuff rare fragments, qtz ll to bed, scattered to trace py, LC 75o, void of stringers.	44849	202.65	204	1.35	0			
			44852	204	205.47	1.47	0.01			
209.29	211.01	patchy to weak patchy sericitic altn, buff to light greenish buff to medium greenish, laminated tuff rare fragments, 210.44-210.54 qtz flood boudinage 211.01 broken ground contact.	44851	205.47	207	1.53	0.01			
			44852	207	208.5	1.5	0.02			
			44853	208.5	210	1.5	0.02	0.02		
211.01	212.09	altn tuff to fragmental tuff, fg to mg, massive, uniform, moderate hard, silicified nil to very weak devel. of foliation with schistosity to weak level bedding, 211.59-212.09 scattered fragments, 211.75-212.09 trace sulphides from white to grey white qtz st LC 80o pervasive weak sericitic altn.	44854	210	211.01	1.01	0.01			
			44855	211.01	212.09	1.08	0			
212.09	214.63	qtz flood fragmental tuff, kinkle to light greenish buff to medium greenish buff, qtz vein random greyish white translucent to white opaque, trace to <0.5 vfg py, LC irregular.	44856	212.09	213.3	1.21	0			
			44857	213.3	214.63	1.33	0			
214.63	215.82	fragmental tuff as above, medium grey green, to medium green, kinkle 215.20 bed 75o scattered py, very weakly carbonated, siliceous, moderate hard to hard, LC irregular x-cut by qtz flood.	44858	214.63	215.82	1.19	0			
215.82	217.91	qtz flood um fragmental tuff, pale greenish kinkle to medium green, to green grey, moderate qtz flooding, trace py, gradual contact, 216.12-216.29 ng grey, moderate carbonate, moderate soft to moderate hard, argillite with 2-3% fg -mg py, up C irregular LC sharp 75o.	44859	215.82	217	1.18	0			
			44860	217	217.91	0.91	0			
217.91	221.47	um fragmental tuff (lower um), carbonated, fg, black to black green, qtz flood fragmental to fragmental tuff, trace sulphides, 218.78-218.83 medium brown	44861	217.91	219	1.09	0			
			44862	219	220.15	1.15	0			

Hunter Mine - Diamond Drill Log HM-04-33

Property:	Hunter Mine	Hole Dip:	-80	Page No.:	1 of 7	Hole HM-04-33				
Location:		Hole Azimuth:	105	Date Started:	July 6, 2004					
Claim No:	HR 1009	Hole Length:	237m	Date Finished:	July 8, 2004					
Elevation:	Porcupine Lake	Purpose:		Drill Co.:	Benoit					
UTM Coords.:	5370904.0N, 487073.0E			Logged by:	K. Jensen					
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
0.0	56.8	Casing								
56.8	89.9	Massive Ultramafic Rocks								
56.8	89.9	very talcose sections, talcose, fg, black to black green, massive uniform, moderately soft to soft, non magnetic to locally magnetic to moderate, locally moderate hard, weak moderate carbonate, qtz + carbonate stringers, rare veinlets 3-4cm, locally magnetite xls, 56.82-60 massive weak devel. schistose random qtz veining 1-3mm + 1cm CA-20-60o random 5-15m, scattered mg cg pyrite 2-3%, 60-63 very broken core, 63-71.46 fragmental tuff UM, weak moderate level schistose bedding schistose random to II CA, local brecciated small section 63.85-63.90 qc vein CA-25o, 64.74-64.89 qc inclusions near CA-25-30o, 65.08-65.13 qc inclusions near CA-30o, 71.46-75.0 brecciated 80% healed qc, irregular fragments, small sections of massive <5cm, rare distinct veins, 75-76.13 massive 75.38-75.54 brecciated qv CA-35o, 76.13-81.20 brecciated large angular fragment qc healed, 81.20-82.19 brecciated small fragments, crushed zone, 82.19 CA-67o shear, 82.19-83.52 brecciated massive white qtz healed, 83.52-83.68 tuff, bed CA-55o, shear 83.61-83.68, 83.68-85.04 massive brecciated, flow breccia, 85.04-85.63 crushed zone, small 3-4mm fragments CA-40 LC, 85.63-86.53 tuff moderate well bedded CA-60-70o, fold more at 85.73, 86.53-87.54 brecciated massive LC 75o, 87.54-89.92 mafic dike UM, massive, fg, weak to nil carbonate weak to moderately magnetic, moderate hard, 87.84-88.73 tuffaceous with 1/2-1 cm py x-cut, with 1-3mm qc st 55o, minor crenulations UC 50o.								
89.92	134.11	Laminated Ultramafic Tuffs								
89.92	93.44	tuff to tuff fragmental, fg, medium grey to greyish black, weak carbonate, minor veining to 90.94, 90.94-92.62 moderate veining scattered fg ng py bed near II to CA at 91.5, 92.62-93.44 fragmental tuff, LC CA-45o sharp.								
93.44	96.76	massive dike, void of all types of stingers nil to trace sulphides, nil foliation or schistosity, moderately soft to soft, non magnetic, weak carbonate, 96.37-96.76 crumbly, shear at 96.72-96.76.								
96.76	97.54	tuffaceous to bedded fragments, shearing II to bed CA 58o.								
97.54	98.28	brecciated massive, white qc healed, 98.08 irregular 1.2-1.5cm qc CA-40o LC65								
98.28	100.72	black green, locally brecciated fragmental weak to moderately magnetic, qtz								

Property: Hunter Mine		Hole No.	33		Sheet No.	3				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
130.58	131.16	distorted crenulated tuff LC50o								
131.16	131.35	grey opaque qtz vein with inclusions CA LC-35o.								
131.35	132.75	crenulated tuff with slips 35o 132.11-132.36, rather qtz fragmental tuff bed 60o 132.75 CA-50o.	44884	131.25	132.75	1.5	0			
132.75	134.11	qtz flooded black to black green tuff, white opaque qtz st LC 65o trace pyrite.	44885	132.75	134.11	1.36	0			
134.11	237	Exhalitic Tuffs and Sedimentary Rocks								
134.11	134.93	grey to light grey crenulated fragmental tuff contorted bedding 134.93-134.94 qtz stringer discont. @ CA-40o, 0o then 90o.	44886	134.11	135.28	1.17	0.03			
134.9	135.28	dark green to black green fragmental tuff with small crenulation scattered mg cg py 1-2% bed 30o LC 70o.								
135.28	136	pale grey to faint green tint, tuff to 135.72 and fragmental to 136 LC 35o. 135.28-135.29 grey qtz st CA 70o.	44887	135.28	136	0.72	0.02			
136	137.45	light grey massive felsic tuff void of stringers, siliceous, hard, broken LC bed 49o, scattered 1% fg, to locally 2-3% ng.	44888	136	137.45	1.45	0			
137.45	139.61	pale grey to pale green well level bed, thin laminations CA-45o ore qtz grey traces to white opaque qtz st from 138.42-139.61, 0.5-1% py with qtz stringer 2-3% fg py in tuff, all qtz st ll to bed LC CA-50o.	44889 44890	137.45 138.42	138.42 139.61	0.97 1.19	0 0.21			0.2
139.61	139.89	grey felsic to siliceous material with grey white trace qtz stringers with fragments and chlorite tuff inclusions scattered py, masses 1-2mm near LC LC-48o.	44891	139.61	139.89	0.28	0			
139.89	144.51	greenish to greyish medium green tuff fragmental fg py, 139.99-140.11 white grey opaque qv with chocolate brown tourmaline and scattered py CA 35-37o 140.93-141.14 white opaque qv with tuff inclusions contacts 40o, 35o, 141.35 white grey q st 2cm 20o irregular, 141.45 white grey q st 2cm 40o irregular sinuous, 141.79-141.84 white opaque minor brown CA-50o, 142.33-142.46 ground lost core, 144.06-144.51 locally patchy white and light to medium grey qtz st usually ll to bed 45o, lost 10cm irregular near ll to CA LC-30o	44892 44893 44884 44895	139.89 140.93 141.84 143	140.93 141.84 143 144.51	1.04 0.91 1.16 1.51	0 0.04 0.03 0			
144.51	145.23	qtz vein altn tuff, fg light brownish grey to dark brown and buff fragments inclusions white opaque to pale grey translucent qv, vein LC 35o vfg py <1%	44896	144.51	145.23	0.72	0.09			0.15
145.23	145.42	light grey to pale green tuff.	44897	145.23	146	0.77	0.02			
145.42	145.62	moderate pervasive sericitic altn tuff.								
145.62	146	patchy sericitic altn tuff.								
146	146.88	light to medium grey to medium grey green tuff, few fragments, well level bedding CA-55o LC 57o.	44898	146	146.88	0.88	0			
146.88	147.45	strong pervasive to string patch sericitic altn tuff, qtz flood from 147.07-147.45 with minor py fg associated with qtz.	44899	146.88	147.5	0.57	0			
147.45	148.28	patchy moderate sericitic altn with 147.77 minor fuchsite, trace to scattered fg py LC 50o.	44900	147.45	148.25	0.8	0.01			

Property:		Hunter Mine		Hole No.	33		Sheet No.	4				
Meterage		Description		Sample		Assay						
From	To	No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check			
		144.43-144.51 crenulated tuff.										
148.28	152.82	pale grey to light pale greenish grey to minor medium grey, trace to very weak patchy sericitic altn <5%, minor small crenulations, whitish and grey qtz string		44901	148.28	149.78	1.5	0.02				
				44902	149.78	151.3	1.52	0				
				44903	151.3	152.82	1.52	0				
		Il to bed 38o-40o at 149.83-149.88 x-cuts 1/2cm grey white opaque CA-65o, 150.15-150.26 laminated qtz stringers with buff altn, 150.90-151.21 laminated greyish white qtz stringer 2mm-1cm, scattered vfg py, bleached bands at 152.38-152.41 and 152.77 1/2cm contact CA sharp 55o.										
152.82	153.68	moderate pervasive to locally medium grey band <5%, scattered fg py, 153.21-153.57 greyish qtz bands with minor trace fg py LC 60o.		44904	152.82	153.68	0.86	0.01				
153.68	155	light grey to medium grey bands laminated tuff, minor grey and white opaque qtz stringers.		44905	153.68	155	1.32	0.03	0.04			
155	155.8	patch moderate sericitic altn, scattered 1-2% fg py locally 3-5% fg ng py, minor fuchsite LC-55o.		44906	155	155.8	0.8	0.02				
155.8	157.6	pale greyish medium green tuff with small fragments, mg tuff, to buff green, minor qtz stringers usually Il to bed, soft 156.95 1/2 grey CA-80o, 157.02 veining 1cm white opaque 80o-90o CA LC 40o bed 50-53o.		44907	155.8	156.7	0.9	0.01				
				44908	156.7	157.6	0.9	0				
157.6	158.89	qtz flood, moderate pervasive to locally patch, minor fuchsite flecks, scattered py LC-45o.		44909	157.6	158.89	1.29	0				
158.89	162.08	weak patchy sericitic altn, light green to locally dark green, 160.63 1cm medium fragment felsic dike 25o 160.11-160.34 pervasive sericitic altn veining white grey qtz flood, Il to bed, 160.84-161.88 ivory qtz stringer Il to CA deform.		44910	158.89	160.11	1.22	0				
				44911	160.11	160.34	0.23	0.02				
				44912	160.34	160.84	0.5	0				
162.08	164.55	medium to dark green tuff, scattered qtz stringer usually Il to bed, 163.06-163.09 whitish grey buff inclusions CA-60o x-cut bed.		44913	160.84	162.08	1.24	0				
				44914	162.08	163.3	1.22	0				
164.55	165.7	brecciated fragmental tuff, matrix, medium to chocolate brown with bleached		44915	163.3	164.55	1.25	0.03				
164.55	165.7	pale grey to buff frag., several white opaque and grey translucent 1/2-1cm qtz veinlets CA-50-65-40o grey, 165.63-165.70 brownish siliceous sections void of inclusions LC-55o.		44916	164.55	165.7	1.15	0	0			
165.7	168.68	grey green fragmental tuff, locally 3-5cm not qtz flooded, very weak sericitic altn patchy, grey white opaque masses + veinlets, local buff fragments trace pyrite LC 50o.		44917	165.7	166.7	1	0.01				
				44918	166.7	167.7	1	0.01				
				44919	167.7	168.68	0.98	0.01				
168.68	171.36	qtz flood, weak patchy sericitic altn to light medium green fragments, white grey opaque qtz rare distinct white qtz st CA-75o.		44920	168.68	169.7	1.02	0				
				44921	169.7	170.7	1	0				
171.36	171.85	light medium green to medium grey tuff minor crenulations small, 171.67 1cm white opaque qtz st Il to bed LC 60o, bed 60o 171.60-171.73 grey 1-2% pyrite.		44922	170.7	171.36	0.66	0.04				
				44923	171.36	171.86	0.5	0.66	0.47			
171.85	172.8	light to medium grey, qtz flood fragmental tuff, trace vfg py, LC slips 60o.		44924	171.86	172.8	0.94	0.15				

Property:			Hole No:			Sheet No.	5			
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
172.8	176.46	moderate pervasive sericitic altn, fragmental tuff qtz flooded, locally fuchsite associated with grey qtz stringer II to bed, bedding at 45o trace to <0.5% vfg py tuff fragmental.	44925	172.8	174	1.2	0			
			44926	174	175.25	1.25	0			
			44927	175.25	176.46	1.21	0			
176.46	176.75	dark grey to dark green, minor light grey laminated tuff LC 63o, small crenulations.	44928	176.46	177.87	1.41	0.01			
176.75	179.25	pervasive moderate sericitic altn, minor grey white qtz stringer up to 1cm from 176.90-177.30, 178.74 large scale crenulation, qtz stringer grey II to bed 178.88-179.25 LC 50o II to bed, few scattered fragments mostly tuff.	44929	177.87	179.25	1.38	0.06			
179.25	180.15	greyish tuff, py 1-2% local bed 53o, 179.56-179.62 ivory qtz, carbonate 55o	44930	179.25	180.15	0.9	0.02			
180.15	181	greyish with 1-3mm black lamination, scattered vfg fg py.	44931	180.15	181.18	1.03	0.01			
181	181.18	pervasive to patch sericitic altn II to bed <1% vfg py.								
181.18	182.76	massive fg ng lappli tuff, scattered up to 1% fg py void of stringers.	44932	181.18	182	0.82	0.03			
182.76	183.06	medium green grey to buff green, several white opaque qtz st II to bed 57-58 LC 57o	44933	182	183.06	1.06	0.01			
183.06	183.96	buff and black green lamination, well level bed, scattered 1% py, siliceous grey bed possible stringer II to bed 58o with associated vfg fg py with tuff near contact of stringer.	44934	183.06	183.96	0.9	0			
183.96	184.88	weak to moderate pervasive sericitic altn, void of stringers good bed few	44935	183.96	184.88	0.92	0			
184.88	188.53	laminated tuff to tuffaceous fragmental greyish to greyish green to black green rare stringers usually II to bed rare x-cut bedding, weakly to moderate	44936	184.88	185.73	0.85	0			
			44937	185.73	186.37	0.64	0.01			
184.88	188.53	carbonated, hard, 184.88-185 blackish grey, 185-185.73 light medium grey, 185.73-186.37 blackish grey to dark grey, scattered pyrite, 186.37-186.96 pale greenish buff to buff, 186.96-187.21 blackish grey to dark green, 187.21-187.42 vfg to aphanitic siliceous, weak sericitic altn, 187.42-188.53 light medium dark grey carbonated, scattered qtz st altn to light grey II to bed 56o scattered fg py, py streaks II to bedding.	44938	186.37	187.41	1.04	0.01			
			44939	187.41	188.53	1.12	0.01			
188.53	189.3	dark grey to black green laminated fg tuff, large crenulations at 189.20 LC sharp CA-51o.	44940	188.53	189.72	1.19	0.01			
189.3	189.72	1mm white carbonated x-cuts in chlorite black green laminated tuff.								
189.72	189.81	tuff dark green, carbonated no x-cuts.	44941	189.72	190.5	0.78	0			
189.81	190.5	greenish buff to medium grey green fg tuff carbonated, LC 65o.								
190.5	191.38	weak patch to moderate patch sericitic altn crenulations at 190.95-191.21, trace vfg py to 1% LC sharp 55o.	44942	190.5	191.38	0.88	0.11	0.11		
191.38	192.63	medium grey + light grey few tuff few fragments, py laminations 1mm 192.27 192.34 scattered vfg fg py 0.5-1%.	44943	191.38	192.63	1.25	0.02			
192.63	194.23	weak to moderate patchy sericitic altn with pale green white opaque and grey opaque qtz st II to bed buff, medium brown sections, siliceous, non carbonate	44944	192.63	193.45	0.82	0.01			
			44945	193.45	194.23	0.78	0.03			

Property: Hunter Mine		Hole No:	34A		Sheet No.	3				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
116.75	119.44	QFP, as above 116.75-117.13 grading from cg to mg, hard, 117.13-119.20 fg porphyritic, rare qtz stringer locally 1mm qtz ff, 119.20-119.38 scattered fg mg py 1-2%, 119.38 contact CA-60o, 119.38-119.44 massive dark green chlorite backed contact CA-80o moderately hard.								
119.44	128.25	tuff fragmental tuff, as above, 119.44-121.50 dark green to black green crenulated fragmental tuff, qtz flooded from 120.83-121.27, 120.14-120.22 white opaque qv 45+50o x-cut bed, nil to trace sulphides, 121.50-128.25 light grey to buff or pale greenish buff grey tuff, local crenulations, locally dark green talcose, 122.97-123.21, sections scattered chlorite II to 3mm slip planes, 123.50 1cm grey translucent qtz at CA-60o, 124.05-124.08 buff crenulated tuff, 124.67-124.70 in white q CA-70o irregular, 124.95-125.12 several qtz veinlets largest 5cm UC 50o, 125.12-126.50 medium grey green to dark grey green tuff crenulated 125.80-126.41 small S folds, 125.12-125.80 fragmental tuff scattered fg py <1%, 126.50 1cm grey white opaque + translucent q at CA-75o, 126.51-128.25 grading of core.								
128.25	129.2	FP, light brown altered to grey to greenish tuff fg with chlorite lineations, foliated moderate devel., tuff inclusions, void of stringers and pyrite, 128.25-128.76 grind of core, 129.20 contact 60o.	44978	128.25	129.2	0.95	0.31			
129.2	129.45	veined altn bleached to buff to tan, brecciated chlorite II healed, brecciated tuff to fragmental tuff, crispy chocolate brown ff tourmaline, diss. vfg 3-5% pyrite and 2-3% vfg fg pyrite in greyish irregular qtz stringer LC ground UC 60	44979	129.2	129.45	0.25	0.45			
129.45	167.86	Exhalitic Tuffs and Sedimentary Rocks								
129.45	131.75	tuff, 129.45-129.82 dark black green massive tuff, 129.82-130.58 light grey to buff patchy ivory tuff, scattered <0.5% vfg py, rare stringers qtz from 130.30-130.45 1-2mm with bleaching on 1-2% vfg fg pyrite, 130.58-130.99 light grey nil to weak patchy to pervasive sericitic altn, pale buff pale green altn of silicified section 130.80-130.88, 130.99-131.75 same as 129.82-130.58, light grey to buff, locally 2-3% vfg py LC 60o crispy brown tourmaline II to bed	44980	129.45	130.58	1.13	0.01			
			44981	130.58	131.75	1.17	1.03	0.79		
131.75	132	felsic dike, aphanitic, pale grey with faint purple tint, hard, veining qtz stringer	44982	131.75	132	0.25	1.86	1.95		
131.75	132	with fg-mg pyrite masses, wispy pyrite stringer near UC, LC 70-75o.								
132	132.39	fragmental tuff bed 55o.	44983	132	133.08	1.08	0.47			
132.39	132.43	very fuchsite siliceous tuff.								
132.43	132.64	dark grey tuff, LC 70o.								
132.64	136	silicified and veined qtz flood tuff, scattered vfg fg py locally contorted, 133.07-133.17 fuchsite silicified tuff scattered fg ng py 1% 55o LC, 133.25-133.53 qtz veined zone, grey opaque random with lumpy chocolate brown tourmaline, altn to medium brown, 133.53-134 large fold bedding II to CA at	44984	133.08	133.54	0.46	0.11			
			44985	133.54	135	1.46	0.01			
			44986	135	136	1	0			

Property: Hunter Mine		Hole No.	34A		Sheet No.	4				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		133.78, 134.0-136.0 crenulated with S folds qtz flooded to light medium brown, LC irregular 40o.								
136	137.43	dark black green fragmental tuff, contorted bedding rare 1cm qtz stringer trace ng py LC 80o.	44987	136	137.43	1.43	0.01			
137.43	149.53	light medium grey tuff massive uniform to light greenish medium grey, trace pyrite, 140.49-140.56 siliceous zone weak sericitic altn with grey opaque qtz veining 140.52-140.56, CA 80o scattered pyrite, 140.65-140.70 same as 140.49-140.56 pale green, 140.70 1cm grey to pale brown qtz st CA 55o with opposite direction, 141.13-141.20 greyish aphanitic felsic dike, trace to <0.5% py CA-80o brecciated qtz 1-2mm healed, 141.23-141.41 qtz st zone pale buff, grey opaque with chocolate brown tourmaline and white grey, LC II to bed CA 75o, 141.50-141.79 light grey tuff with white grey opaque qtz veinlets 1.5 and 2cm at contacts, 143-144 numerous chlorite II 1-3mm slip planes 25o-35o, 144.09-144.19 grey siliceous vein with tuff inclusions, trace to nil py, 45o and 70o sinuous contacts, 144.66-145.23 qtz flooded, light medium grey tuff, scattered to nil pyrite, 145.43 irregular grey translucent qtz st 1-1 1/2cm irregular, 145.88-146.53 silicified zone few irregular grey translucent qtz stringer with 146.17-146.34 white translucent qtz st II to CA on edge of core, 146.22-146.27 brecciated zone buff to light brown fragments healed greyish and blackish silicin, contacts 75o+70o, 146.53-147.38 grey green, tuff and veined tuff fragmental, 146.95-147.38 with scattered fg py 60o 80o contacts, 147.38-148.30 light brownish grey, fg massive poor devel. of bed tuff fragmental, 148.09-148.30 irregular S shaped UC to siliceous LC 50o qtz masses and stringer, 148.30-149.53 fragmental tuff light grey to light greenish grey.	44988 44989 44990 44991 44992 44993 44994 44995 44996 44997 44998 44999	137.43 138.93 140.43 141.13 141.8 143 144 145.43 145.88 146.53 146.53 147.38 148.03 148.03	138.93 140.43 141.13 141.8 143 144 145.43 145.88 146.53 147.38 148.03 149.53	1.5 1.5 0.7 0.67 1.2 1 1.43 0.45 0.65 0.85 0.65 1.5	0.02 0 0.01 0.01 0 0.03 0 0 0 0.09 0 0			0.07
149.53	150.73	siliceous q breccia zone, 149.53-149.70 buff brecciated fragments with grey to brownish grey siliceous matrix, 149.70-149.75 medium brown siliceous, 149.75-150.27 grey and white brecciated qv healed with unszp ff chocolate brown tourmaline x-cut by white 1cm qv CA 45o at 149.82-149.84, 149.92 1/2cm grey white qtz st CA33o, 150.27-150.30 medium brown siliceous, 150.30-150.73 brecciated buff fragments healed with chocolate brown siliceous 150.53 chlorite and white qtz stringer 1/2 to 1cm CA 30o terminates 1/2 white qtz stringer at 10o 150.53-150.66, 150.73 irregular CA 40-60o.	45000	149.53	150.73	1.2	0.01			
150.73	151.67	light grey tuff rare st 150.74 1cm bed 78o, 150.90 1cm grey opaque 75o.	43001	150.73	151.67	0.94	0.02			
151.67	152.19	pale green, scattered fuchsite tuff 151.76 1cm white opaque and grey translucent st q 80o II to bed, 151.84 3mm siliceous and fuchsite x-cut bed 75o, scattered mg py.	43002	151.67	152.19	0.52	0.02			

HunterMine - Diamond Drill Log HM-04-35

Property:	Hunter Mine	Hole Dip:	-75	Page No.:	1 of 11								
Location:		Hole Azimuth:	105o	Date Started:	July 15/04								
Claim No:	HR 1009	Hole Length:	248.87m	Date Finished:	July 19/04								
Elevation:	Porcupine Lake	Purpose:		Drill Co.:	Benoit								
UTM Coords.:	5370818.4N, E487097.3			Logged by:	K. Jensen								
Meterage		Description				Sample				Assays			
From	To					No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
0.0	75.0	Casing 0.00-54.0, water and overburden water to 11.5', 54-75 reamed.											
75.0	75.0	Ultramafic Talcose Rocks											
75.0	98.6	Ultramafic, same as usual, black to black green, moderately soft, talcose, non magnetic, massive to brecciated, 2-5 qtz stringers / meter, fg, nil to weak schist, 75.0-79.15 brecciated, contorted schist large angular fragments, trace pyrite occasional cg pyrite cubes, 79.15-84.85 massive, uniform, locally carbonated veining random overall with fragments, nil schistosity, 84.85-92.80 brecciated massive, intensely Bx with carbonate filling usually 1-2cm angular fragments, local devel. of schist at 50-55o at 92.5, void of distinct veining, 92.80-94.48 fragmental tuff to brecciated fragments, void of distinct veining, locally moderately devel. schist 25-30o 93.20-93.94, 93.94-94.48 contorted schistosity, LC broken											
94.48	133.8	Laminated Ultramafic Tuff											
94.48	95	crenulated tuff, well devel. bed & schist. 75o, 95.0-96.38 fragmental tuff contorted schist. locally, 96.38-96.89 foliated massive mg to cg porphyritic texture LC ground, 96.89-97.18 white opaque qtz with talcose inclusion nil sulphides, 97.18-97.33 silicified talcose fragmental, 97.33-97.91 massive weak to weak moderate level of foliation, scattered 1-2mm carbonated st CA 40-45o II to foliated, 97.91-98.21 brecciated tuffaceous, 98.21-98.55 massive low foliation LC 15o x-cut undertion tuff with bed CA-50o.											
98.55	99.5	fragmental tuff brecciated healed with random orientated qtz stringers.											
99.5	99.8	crenulated tuff											
99.8	102.05	qtz flood tuff, grey green to medium green random white opaque 1-4mm qtz st scattered fg py 102.05 LC irregular 50-55o.											
102.05	108.4	light to medium grey with greenish tint, tuff, moderately hard to moderately soft, with chlorite II slips planes, 1-4mm white opaque qtz st, with occasional white translucent qtz veinlets 1/2cm to 1cm, 103.55 CA 45o II to bed, 103.76 1.5cm CA-50o, 103.96 irregular 1cm + py II to crenulated tuff, 104.25-104.29 CA-35o, 104.89 II to bed 43o, 105.44 1cm CA-40o, 106.12-106.39 irregular qtz veining with contorted black green tuff CA-15+45o, 106.39-108.40 tuff to tuffaceous fragmental qtz flooded locally crenulated tuff, grey white translucent qtz st 1/2-1.5cm LC 20o				43022	106.34	107	0.66	0.05			
					43023	107	108.4	1.4	0				

Property		Hunter Mine			Hole No.	35		Sheet No.	2			
Meterage		Description			Sample			Assay				
From	To				No.	From	To	Width	Au (g/t)	Au Check	Au (2nd)	Au check
108.4	111	mafic metavolcanics, fg to aphanitic, light green at upper contact, to blackish green locally medium green chlorite, massive, brecciated, very hard silicified qtz filled breccia with grey translucent, greyish brown with vfg py, white opaque qtz st generally 1-2mm few up to 1/2cm are above 1cm, nil to very weak devel. of schistosity, non magnetic, non carbonate, 108.40-111.0 massive, 1-2% vfg fg overall, locally 3-5% 108.94-110.0.			43024	108.4	109	0.6	0.15	0.14		
					43025	109	110	1	0.04			
					43025	110	111	1	0.03			
111	113.03	tuff fragmental to fragmental flow top, scattered vfg py, 111.93-112.04 contact bleaching dark brown, 112.04-112.33 qtz veinlets brecciated with altn brown inclusions tuffaceous 1-2% vfg py.			43027	111	111.98	0.98	0.01			
					43028	111.98	112.33	0.35	0.01			
					43029	112.33	113.15	0.82	0.01			
113.03	114.9	silicified tuff, with rare grey brown qtz veinlet randomly overlaid x-cut locally by greyish white translucent qtz st, tuff 1-2% vfg py, brownish scattered to 1%, white 113.39-113.60 with scattered fg py splashes of chalcopyrite and vfg masses of pyrrhotite irregular CA-20+15o, 113.15-113.30 bleached light brown by irregular white opaque qtz st at 113.21-113.25, UC 35o II to bed, 114.12 1/2cm grey white translucent II to bed 25o, 114.58-114.85 light grey tuff with black green talcose LC 45o bed 30o, 114.75 2cm buff white opaque qtz v CA-47o, 114.40 LC contact CA-30o.			43030	113.15	114	0.85	0.02			
					43031	114	114.9	0.9	0.03			
114.9	237.47	Exhalitic Tuffs and Sedimentary Rocks										
114.9	115.45	fragmental tuff, moderately hard to hard to 115.04 grey green, qtz flood with pale brownish white opaque qtz, trace pyrite, LC 45 with qv 1 1/2.			43032	114.9	115.45	0.55	0			
115.45	116.93				tuff with occasional white fragments, grey green, brecciated tuff, siliceous, qtz st, grey to pale brownish grey 10-15m, locally crenulated with chlorite II slip planes, scattered pyrite fg, moderately hard st usually II to bed, follows crenulation, white translucent rare + x-cut bed, LC ground.			43033	115.45	116.93	1.48	0
116.93	118.1	fragmental tuff, fg altn light brown to 117.17 felsic tuff fragmental, sinuous CA 65-55o dark green light grey laminated fragmental tuff, rare qtz st, scattered pyrite LC 70o sineous.			43034	116.93	118.1	1.17	0			
118.1	118.93				white translucent qtz vein with pale brown inclusions and grey green altn chlorite nil sulphides LC ground.			43035	118.1	118.71	0.61	0.01
118.93	119.79	chlorite crenulated locally, tuff, thin laminations, 119.11 1/2cm grey opaque qtz st CA-65o 119.09-119.29 white translucent qtz st on side of core LC 75-80o 119.65 light brown felsic.			43036	118.71	119.79	1.08	0			
119.79	120.5				fg, light brownish altn, locally greyish brown with black to black green chlorite schist tuff inclusions, random orientated 1-2mm up to 1cm qtz white			43037	119.79	120.5	0.71	0.01
119.79	120.5	translucent stringer usually II to CA to 70-75o, scattered pyrite in felsic dike, trace to scattered upper part 119.79-120.0 with 2-3% fg locally LC 50-55o irr.										
120.5	121.5	black to black green crenulated tuff with 1m-4m chlorite II slip planes,			43038	120.5	121.5	1	0			

Property		Hunter Mine		Hole No.	35			Sheet No.	3		
Meterage		Description		Sample			Assay				
From	To			No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		scattered pyrite near contacts, 120.54 3-4cm white opaque qv CA 60o, void of stringers LC irregular 35o.									
121.5	122.75	same as 119.79-120.5, up to 123.06 trace pyrite, 123.06-123.37 scattered sections of fg pyrite overall 1%, 123.37-123.51 tuff inclusions chlorite irregular masses black, 123.46-122.75 hydro fracturing, qtz ff, random qtz st 3-5mm 122.75 irregular qtz veinlets 5cm LC 35o.		43039	121.5	122.75	1.25	0.01			
122.75	125.63	silicified medium grey felsic dike, nil to very poorly devel. foliation with minor inclusions of black green tuff, several white to white grey opaque qtz stringer usually at 60o, numerous white ll and brecciated filled qtz stringers, random irregular 1-3mm and irregular mass breccia vein 123.33-123.38 CA-55o small 1-3mm qtz stringer fg py CA-35o,25o,50o, level sections altn to pale brownish grey, scattered 1-2% fg py locally 3-5%, 125.40 altn contact 55o LC CA broken + ground.		43040	122.75	123.84	1.09	0.08	0.05		
				43041	123.84	124.67	0.83	0.04			
				43042	124.67	125.63	0.96	0.08			
125.63	127.2	very hard to hard, silicified brecciated tuff fragmental contorted bed to 126.62 qtz flood to 127.20 void of stringers, nil to trace py, LC ground.		43043	125.63	126.63	1	0			
				43044	126.63	127.2	0.57	0			
127.2	130.19	moderate hard, tuff to fragmental tuff, medium grey green to medium greenish grey locally crenulated tuff, distorted bedding, 128.0-128.09 light brownish to brownish grey felsic dikelet CA-40o, note 126-128 41cm ground lost, 128.86-129.0 greyish siliceous bands with 1-3% scattered fg py CA 55o curved, 129.0-129.27 light greyish light brown felsic dike with tuff inclusions 129.17-129.22 CA 73o irregular LC irregular 60-70o medium brown 129.22-129.27, 130.07-130.19 felsic dike light brown, siliceous with irregular white qtz and grey white qtz stringers, 130.31-130.40 white translucent qtz vein with chlorite inclusions CA irregular 60o, 130.42-130.55 same as 130.31-130.40.		43045	127.2	128	0.8	0			
				43046	128	129	1	0			
				43047	129	129.27	0.27	0.04			
				43048	129.27	130.07	0.8	0			
130.19	132.68	black green fg with grey siliceous brecciations occasional fragment, small S folds, local crenulations, local bed well level CA-65o trace to scattered py, void of stringers except 130.31-130.40 and 130.47-130.55, LC 70o.		43049	130.07	130.55	0.48	0			
				43050	130.55	131.65	1.1	0			
				43051	131.65	132.68	1.03	0.09			
132.68	133.4	black green very contorted bedding S folds.		43052	132.68	134.02	1.34	0			
133.4	134.02	black green with grey laminations minor crenulations and chlorite ll slip plane CA-30-45o, LC altn 75o.									
134.02	136.32	light medium green, tuffaceous to 134.90 more massive poorly devel. bed sch. to 136.32, weak pervasive sericitic altn, LC 73o, fg massive ivory qtz st CA-60o.		43053	134.0	135.0	0.98	0			
				43054	135.0	136.3	1.32	0.18			
136.32	140.2	fg, light medium green to medium green to grey green, well devel. bedding locally brecciated, chlorite ff slip planes 139.73-140.64, locally fragmental, grey white qtz stringer at 137.57 opalescence 1/2cm, ground core, 138.54-138.57 CA-85o x-cut bed CA-65o, 139.55-139.60 ll to contorted bedding 70o+35o, scattered fg py, few 2-3mm masses of py, weak to moderately weak patchy to locally		43055	136.3	137.8	1.48	0			
				43056	137.8	139.3	1.5	0			
				43057	139.3	140.4	1.1	0			

Property: Hunter Mine		Hole No.	35		Sheet No.	4					
Meterage		Description	Sample				Assays				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check	
140.2	142	greywacke siliceous laminations locally pale grey to medium grey green, hard well level 570-470.	43058	140.4	141.5	1.08	0.01				
			43059	141.5	142.8	1.29	0.01				
142	142.77	qtz flooded light grey green to medium green, fragmental tuff, trace pyrite, grey white qtz stringer mostly ll to contorted bedding, few x-cut bedding.									
142.77	144.64	light grey green, local buff fragments, fragmental tuff to tuff, 142.77-143.13 heavy veining and grey white qtz masses, 143.13-143.51 scattered irregular grey white qtz stringer contorted x-cut bedding, bedding tuff crenulated with 1mm chlorite ff slip, 143.51-143.69 1/2-1cm grey whit qtz string x-cut bed and ll to CA trace pyrite, 143.69-143.94 qtz white vein with medium green tuff inclusions CA-70-60 irregular, 143.94-144.64 medium green weak pervasive sericitic altn contorted crenulated tuff with 1-2mm scattered qtz st x-cut bedding, 144.14-144.16 irregular grey white translucent qtz st CA-80o 144.64 contact sharp CA 65o irreg	43060	142.8	144.0	1.23	0.01				
			43061	144.0	144.6	0.64	0.01	0			
144.64	145.5	qtz vein. Brown tourmaline, fg to aphanitic, light to pale brown, with 2-3mm white ghost phenocryst, aimilar to 121.5-122.75 and 119.79-120.5, minor chlorite ll, few grey white qtz stringers 2-4mm with chocolate ll tourmaline, scattered vfg fg py, 144.86-145.03 medium grey white 3-4mm white phenocryst ghosts, moderately devel. foliation CA-45o UC-50o LC Irregular, 145.03-145.07 white translucent qtz v CA-45o irregular, 145.50 irregular CA-60o.	43062	144.6	145.5	0.86	0				
145.5	147.83	light grey green to medium green patchy weak sericitic tuff, crenulated locally with 1m chlorite ll slips few grey contorted 3-5mm qtz stringers.	43063	145.5	146.3	0.8	0				
			43064	146.3	146.8	0.5	0				
147.83	148.15	fragmental tuff light grey green to medium green patchy sericitic altn LC 40o	43065	147.8	149.2	1.35	0.01				
148.15	148.24	light brown qtz vein with inclusions irregular trace pyrite.									
148.24	149.18	fg laminated crenulated tuff void of stringers weak pervasive sericite.									
149.18	150.63	light greenish to medium greyish green with qtz grey laminated tuff, locally crenulated weak patchy sericitic altn, 150.20-150.33 greyish felsic band with	43066	149.2	150.2	1.02	0.02				
			43067	150.2	150.6	0.43	0.02				
149.18	150.63	weak sericitic tuff inclusions, 150.34 1/2cm greyish white opalescence qtz st CA-50o, 150.34 1/2cm greyish white translucent qtz st CA-25o both x-cut bed 65o, 150.52-150.57 light brown to buff qtz vein scattered vfg py CA-80o 40o slips, 150.63 contact CA sharp 70o.									
150.63	150.8	felsic hydro fractured dike possible massive ash tuff, fg to aphanitic, medium grey, massive uniform with pale yellowish green ll hydro fractions with chalcopyrite void of laminations, schistosity etc, void of normal qtz st, hard to very hard, siliceous, non magnetic, non carbonated, matrix has vfg to fg py 3-5% overall LC CA-80o.	43068	150.63	150.8	0.17	0				
150.8	151.22	very similar to 150.63-150.80 both with fragments, well level bedding CA-75o chlorite filling between fragments and layers, scattered fg py and occasional 1mm bands, felsic fragmental tuff, overall 2-3% pyrite, 151.22 contact sharp	43069	150.8	151.22	0.42	0.27	0.27			

Property: Hunter Mine		Hole No.	35		Sheet No.	5				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		CA 55o-57o.								
151.22	151.42	greyish brown altn fragmental terminates in chloritic tuff UC at 70o, II to CA-30o LC gradual with qtz flooding.	43070	151.22	151.85	0.63	0			
151.42	151.72	medium green to light green locally weak moderate sericitia altn tuff lapilli tuff contrtd bedding.								
151.72	151.85	white translucent qtz veinlets irregular with inclusions black green patches of chlorite.								
151.85	151.93	white opague qtz vein CA irregular 20o.								
151.85	152.89	brecciated, light grey buff fragmental, minor to nil tuff, 152.35-152.48 1cm white opague with medium brown tourmaline, CA-60o, 152.42-152.47 curved 1/2-1cm white qtz st with brown tourmaline, 152.49 1cm white opague qtz veinlet CA-55o opposite to 152.35-152.48 stringers, 152.75 grey white qtz stringer translucent CA-35o, 152.80-152.87 irregular white opague on contact with grey white translucent in center, trace to <1% vfg py, 152.89 contact sharp CA 60-65o.	43071	151.85	152.89	1.04	0.02			
152.89	154.25	medium green to dark green fragmented tuff with randomly orientated 3-5mm white veined with grey center qtz st, trace to scattered vfg, moderate level bed, 154.25 sharp CA 40o.	43072	152.89	154.25	1.36	0			
154.25	156.22	light to medium green fragmental tuff, moderate to well devel. bedding CA 63o, 155.97-156.0 grey to buff bleaching, 156-156.07 irregular CA-35o, white opague qtz veinlet with brownish altn, fg py 1% in vein, 156.07-156.22	43073	154.25	155.3	1.05	0.03			
			43074	155.3	156	0.7	0			
154.25	156.22	blanked tuff, few fragments, buff to bluish hue green.								
156.22	156.47	bleached grey brown to med dark chocolate brown brecciated fragmental x-cut by 1-3mm grey white qtz st CA-15o, 80o, 50o, + large 2 white V shaped qtz opague stringer 156.32-156.47 CA-50o with scattered ng py, large 1cm by 4mm pyrrhotite, magnetic mass at lower contact CA-30-35o, 2mm x 1.2 pyrrhotite and chalcopyrite vg, 156.47 altn contact CA-75o.	43075	156	156.47	0.47	0.22			
156.47	158	light green to light brownish buff weak patchy sericitic altn, tuffaceous fragment minor small crenulations, scattered vfg py, good bed CA-70o, rare stringers 156.92 2m x-cuts bed CA-15o, 157.91-157.95 patches of grey white opalescant qtz st masses II to bed, 158.0 contact CA-60o.	43076	156.47	157.25	0.78	0			
			43077	157.25	158	0.75	0.01			
158	159.32	dark green with pale green buff laminations minor, scattered grey white opal qtz stringers masses, 158.47-158.49 2.5cm white grey qtz st CA-65o near II to bed, 158.70-158.80 pale grey buff altn with grey white siliceous, 2-3% fg ng pyrite CA-65o 158.85-159.32 very fine grained laminations with 1/2-1cm grey buff qtz carbonation laminations, 159.32 contact CA-75o.	43078	158	159.32	1.32	0			
Property: Hunter Mine			Hole No.	35		Sheet No.	6			

Property: Hunter Mine		Hole No.	35		Sheet No.	6				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
159.32	159.72	light green to light buff, pale grey, well laminated tuff, locally crenulations, scattered py, py 2mm band at 159.44 CA-80o bedding overall 75o.	43079	159.32	159.72	0.4	0			
159.72	163.24	moderate pervasive sericitic altn to locally patchy 161.66-163.24, scattered greyish 1/2-1cm bands, scattered vfg to fg pyrite locally 2-3%, moderate qtz flooding II to bed, CA-70o at 160, 60o at 161.5, rare grey translucent qtz stringer 1/2cm x-cut bed at 75o, bed at 163.05 70o.	43080	159.72	160.39	0.67	0.09			
			43081	160.39	160.95	0.56	0.1			
			43082	160.95	161.66	0.71	0.04			
			43083	161.66	162.45	0.79	0.01			
163.24	164.9	buff grey to light and medium grey pale green to weak patchy sericitic altn, tuff, few whitish qtz st usually II to bed occasional x-cut bedding and kinkle folded, scattered pyrite, nil to locally <0.5, weak carbonated, well bed.	43084	162.45	163.24	0.79	0.01			
			43085	163.24	164	0.76	0.01			
			43086	164	164.95	0.95	0.01			
164.9	165.74	weak moderate sericitic altn pervasive tuff.	43087	164.95	165.74	0.79	0.42			
165.74	166.28	weak patchy sericitic tuff.	43088	165.74	166.28	0.54	1.4	0.96		
166.28	166.96	weak moderate pervasive sericitic altn tuff, scattered py, bed 75o altn II-contact	43089	166.28	166.96	0.68	0.04			
166.96	168.97	nil to weak patch sericitic altn tuff, chlorite bands vfg pyrite locally <0.5-1%	43090	166.96	168	1.04	0.01			
		168.02-168.08 grey opaque qtz stringer 1/2cm and creamy white qtz CA 75-65	43091	168	168.97	0.97	0.01			
168.97	169.66	moderate pervasive sericitic altn, scattered pyrite.	43092	168.97	169.66	0.69	0			
169.66	171.2	grey green tuff nil to very weak sericitic patchy, 170.96-171.14 crenulations with chlorite ff stringers.	43093	169.66	171.2	1.54	0.01			
171.2	171.38	qtz flood tuff with scattered vfg fg py, bed CA-50o contact 35o with weak to moderate sericitic altn.	43094	171.2	171.38	0.18	0.12	0.14		
171.2	171.38	moderate sericitic altn.								
171.38	172.19	brecciated zone of tuff with occasional fragment, contorted bedding, healed with brown siliceous and occasional grey cherty q to 171.74, 171.74-172.19 healed grey opaque qtz and stringers, minor pyrite, scattered pyrrhotite in 171.38-171.74, LC 60o.	43095	171.38	172.19	0.81	0.01			
172.19	172.4	qtz flooded with weak sericitic altn LC 75o.	43096	172.19	172.4	0.21	0.01			
172.4	173.08	light green to grey green, weak patchy sericitic altn, rare stringers, trace py.	43097	172.4	173.08	0.68	0.01			
173.08	173.34	c	43098	173.08	174.03	0.95	0.01			
173.34	173.63	qtz vein white opaque with stringers and brecciated inclusions of medium green tuff, qtz vein at 173.36-173.44.								
173.63	174.03	same as 172.40-173.08.								
174.03	174.28	chesty greyish qtz vein x-cut by greyish opaque qtz stringers, chocolate brown ff, inclusions of weak sericitic altn tuff, contacts 50o-70o trace pyrite opposite direction, brecciated 174.03-174.09.	43099	174.03	174.28	0.25	0.03			
174.28	174.67	weak pervasive sericitic altn, tuff LC-70o.	43100	174.28	175.59	1.31	0.04			
174.67	174.81	irregular greyish white qtz stringer, tuff with minor fuchsite altn, 2-3% vfg to fg pyrite.								
174.81	175.59	weak patchy sericitic altn minor fragments 175.42 1/2cm grey qtz st CA 65o x-cuts bed 65o.								
175.59	176.23	qtz flooded usually II to bedding and x-cut bedding grey white translucent, with moderate pervasive sericitic altn.	43101	175.59	176.23	0.64	0			

Property: Hunter Mine		Hole No.	35		Sheet No.		7				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check	
176.23	178.64	weak pervasive sericitic altn, tuff grey white 1/2 opaque qtz stringer, x-cut bedding at 65o straight + 45o kinkle folded to 177.0, 177.73-178.64	43102	176.23	177.73	1.5	0.02				
		moderate qtz flooding all II to bed LC sharp 70o.	43103	177.73	178.64	0.91	0.01				
178.64	179.65	chloritic tuff, fg, dark grey to dark grey green, siliceous, non carbonated, locally contorted bedding near white translucent glassy vein 179.14-179.30 with grey brown to buff altn tuff, 179.23-179.29 with 2-3% fg py, chloritic tuff locally 2-3% fg py, greyish cherty bands.	43104	178.64	179.65	1.01	0.01				
179.65	181.04	weak pervasive sericitic altn, laminated tuff with band grey white translucent qtz stringer usually II to bed, few patches of fuchsite, LC bed 70o-75o.	43105	179.65	181.04	1.39	0.02				
181.04	182	light tan to light brown, locally dark to medium greyish tan, well level bedding and laminations, weak sericite.	43106	181.04	182	0.96	0.04	0.1			
182	182.91	dark brownish to medium grey brown, laminated with pale sections altn by white qtz stringer, trace pyrite.	43107	182	182.91	0.91	0.02				
182.91	185.25	similar to 181.04-182.0, weak sericitic altn scattered fg py occasional 1mm bands, 1-2m 2mm whitish qtz st II to bedding.	43108	182.91	184	1.09	0.01				
			43109	184	185.25	1.25	0.04				
185.25	188.15	weak to moderate pervasive sericitic altn scattered vfg fg py, occasional 1-2mm py band, light brown to tan, uniform, rare qtz st, scattered vfg fg py.	43110	185.25	186.75	1.5	0.02				
			43111	186.75	188.15	1.4	0.01				
188.15	190.48	weak pervasive to locally patchy sericitic altn, light brownish medium green to medium greyish medium green, massive uniform, rare stringers, void of folding, bed 75o.	43112	188.15	189	0.85	0.02				
			43113	189	190.48	1.48	0.03				
190.48	194.32	scattered weak patchy sericitic altn, tuff, scattered to trace pyrite fg vfg to 192.0, rare stringers, 192.0-194.32 fairly brecciated vfg to fg pyrite with occasional bands 1-2mm, local small crenulations with grey green units	43114	190.48	192	1.52	0.08				
		192.10-192.12, 192.69-192.98, overall py 2-3% ng to 3-5% LC sharp 72o, II to bed.	43115	192	193	1	0.14				
			43116	193	194	1	0.15				
			43117	194	194.32	0.32	0.63	0.59			
194.32	194.81	felsic dike aphanitic to fg, massive uniform, hard, siliceous, greenish grey to medium grey, very weakly foliated specially 194.45-194.81 nil sulphides, low angle grey white banded 4cm qtz stringer, 194.57-194.74 LC broken.	43118	194.32	194.81	0.49	0				
194.81	196.81	fg, black green to bark green, laminated chlorite with greyish siliceous 1-2mm bands, locally crenulations from 195.43-196.21 with 2-4mm chlorite II slip and 196.66-196.81, tuff LC 50o, rare grey 1cm qtz st CA folded.	43119	194.81	195.81	1	0				
			43120	195.81	196.81	1	0				
196.81	197.45	similar to 194.81-196.81 but fragmental tuff, chlorite.	43121	196.81	197.45	0.64	0.01				
197.45	197.74	qtz flooded grey green tuff fragmental irregular contacts chlorite.	43122	197.45	197.74	0.29	0				
197.74	200.43	light grey green to kinkle green, weak pervasive sericitic altn tp 198.66, tuff to fragmental tuff scattered to nil pyrite, 197.92 2cm qtz light grey translucent and carbonated irregular, 198.29 1/2cm light grey translucent with white carbonated CA 75o x-cut bed 75o, 198.29-198.66 crenulated tuff with black	43123	197.74	199	1.26	0				
			43124	199	200.43	1.43	0				

Property: Hunter Mine		Hole No.	35		Sheet No.	8				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		green chlorite II slip, 198.66-200.43 fragmental tuff, light opal qtz st CA 80-75o cherty, 200.43 LC 55o.								
200.43	200.73	pinkish to greyish pink qtz vein, glassy, translucent, 1-2mm x-cut greyish white qtz st, nil sulphides, LC sharp 75o-80o.	43125	200.43	200.73	0.3	0			
200.73	202.22	same as 198.66-200.43 fragmental buff grey green, very weak patchy sericitic altn, trace py LC 60o.	43126	200.73	202.22	1.49	0			
202.22	203.03	tuff same as 198.29-198.66 with crenulations void of stringers, dark green to dark greyish green, laminated with grey white silica, LC 65o x-cut bed 70o.	43127	202.22	203.03	0.81	0.01			
203.03	203.28	black green tuff as 202.22-203.03 with 3 greyish white qtz veinlets 2-4cm ash near to x-cut bedding, LC irregular 60-65o.	43128	203.03	203.28	0.25	0			
203.28	203.45	white opaque qtz vein nil sulphides CA-70o LC.	43129	203.28	203.45	0.17	0			
203.45	207.69	FP dike, aphanitic to fragmented light salmon to medium salmon pink in sections overall greyish white, with 1-2mm plagioclase phenocryst fine pin prick black phenocryst locally dark grey ff random 203.45-203.96, 204.4-204.60, white grey from 203.85-203.96 + 206.97-207.69, random grey translucent glassy 2-3mm stringers random with rare 2cm veinlet at 204.75 near II to CA scattered chalcopyrite at 204.69, 205.4, 205.73, 205.93, 206.28, 205.82 qtz st grey translucent CA-80o, random qtz st 206.10-206.89, 207.69 contact sharp 60o.	43130	203.45	204	0.55	0.09			
			43131	204	204.65	0.65	0			
			43132	204.65	205.25	0.6	0			
			43133	205.25	206	0.75	0.03			
			43134	206	206.97	0.97	0.01	0		
			43135	206.97	207.69	0.72	0			
207.69	208.72	chlorite tuff UM, fg black to black green, crenulated with chlorite ff slip planes 2-3mm siliceous, as above void of stringers, trace pyrite, silicified contorted bedding, sharp contact LC-35o.	43136	207.69	208.72	1.03	0			
208.72	209.17	grey feldspar porphyritic dike, fg light to medium grey, 1-2mm white plagioclase phenocryst, massive hard to very hard, random qtz st from 209-209.12 CA-10-15o x-cut to LC and terminated at lower contact 209.03 1cm grey translucent qtz st CA irregular 40-45o nil sulphides LC sharp CA-50o.	43137	208.72	209.17	0.45	0.02			
209.17	209.68	silicified tuff, fg, light medium grey, silicified tuff with random qtz st and qtz mass 209.31-209.52 irregular, hard, chlorite layers, trace LC 45o x-cuts bedding CA-80o foliation.	43138	209.17	209.68	0.51	0			
209.68	209.96	grey feldspar porphyritic same as 208.72-209.17 with more qtz st and irregular qtz veinlet 209.71-209.79, trace py LC grund broken CA-90o.	43139	209.68	209.96	0.28	0.03			
209.96	211.2	fragmental tuff, same as above, light grey green, white laminations, minor kinkle folding, bed good CA 70o, silicified trace py, rare contorted grey translucent qtz st 4-5mm LC graduation 60o.	43140	209.96	211.2	1.24	0.01			
211.2	211.59	qtz flooded random rare 1cm qtz st grey white CA-70o at 211.32 with pale yellow buff altn weak sericite.	43141	211.2	211.59	0.39	0			
211.59	211.96	moderate sericitic altn, siliceous tuff with large 1cm bands qtz inclusions and	43142	211.59	211.96	0.37	0			

Property: Hunter Mine		Hole No.	35		Sheet No.	9				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		firm laminations with fg mg pyrite 3-5% locally LC sharp CA-70o.								
211.96	215.01	tuff to fragmental tuff, fg, pale greenish grey to light brown to blackish green	43143	211.96	213	1.04	0			
		locally sections with white calcite x-cuts, porphyritic tint, moderately	43144	213	214	1	0			
		carbonated rare qtz at 212.47 1/2cm CA-65o, bedding CA-70-60o 213.15,	43145	214	214.75	0.75	0			
211.96	215.01	carbonated, moderately hard to hard, siliceous with local patches sections	43146	214.75	215.01	0.26	0.07	0.1		
		3-4cm of carbonated x-cuts ultramafic, rare st, vfg to fg tuff, occasional								
		fragments small, 214.85-214.95 scattered fg to vfg pyrite 1-2%, 214.95-215								
		semi massive py with qtz 85%, 213.08 ground core, LC 60o.								
215.01	216.97	argillite greywacke, fg, light to medium grey argillite to fine grained greywacke	43147	215.01	216	0.99	0			
		moderately carbonated, moderately hard, uniform, scattered randomly								
		orientated qc st 1-2mm from 215.01-215.36 80-70o in both direction few II to								
		weak devel. of bedding, at 70o, 215.65- 216.18 4mm -1cm qc II to bedding,								
		216.97 LC 73o.								
216.97	217.81	216.97-235.86 ultramafic, 216.97-217.81 black green, weak moderate level of								
		bedding tufferous, non magnetic.								
217.81	218.92	tuffaceous olive to light greenish buff, weak moderate level bedding, hard,	43148	217.81	218.92	1.11	0.01			
		siliceous to silicified, scattered qtz stringers usually II to bedding, scattered								
		vfg fg pyrite locally 1-2% occasional with randomly orientated grey opaque								
		siliceous, carbonated.								
218.92	219.42	black gren moderate hard tuff.	43149	218.92	219.56	0.64	0.01			
219.42	219.56	irregular qc veinlets with chloritic crenulated tuff inclusions, nil to trace pyrite,								
		CA 60o.								
219.56	232.43	mass, moderately soft to soft, carbonated talcose, massive with brecciated								
		healed with qtz st random, locally section II to schistosity, trace to scattered								
		pyrite, 221.58-221.68 porphyritic texture, 222.59-223.20 tuff with several folds								
		223.20-227.14 massive, brecciated qc healed, 227.14-227.68 tuffaceous with								
		qc II to bedding, local fragments, 227.68-230.60 massive brecciated black								
		230.52-230.60 massive talcose vein soft, 230.60-231.06 massive with few st								
		qc, soft, LC 50o, 231.06-232.43 moderately hard, light green, tuff to fragment								
		local crenulations 232.33-232.43.								
232.43	235.86	moderately soft to moderately hard, carbonated, light green to medium grey								
		green, massive, carbonated, scattered random qc st and veinlets.								
234.45	235.86	black green moderately hard massive large fragments within carbonated tuff								
		to fragmental tuff, qtz carbonated vein II to schistosity bedding CA-77-60o,								
		235.63-235.86 moderately hard, silicified intensely veined with grey white qc								
		stringer II to bed LC CA-70o.								
235.86	249	meta sediments, argillite to greywacke with graphitic argillite, light grey to	43150	235.86	236.72	0.86	0			

Property		Hunter Mine	Hole No.	36		Sheet No.	2			
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au Check	Au (2nd)	Au check
		103.26 crushed crumbly zone, 106.37-106.47 crushed crumbly core, 108-110.31 tuffaceous with fragments, less fragments as above.								
110.31	111	tuff bed CA-20o-60o.								
111	113.29	fragmental tuff moderately to strongly schistose, talcose.								
113.29	113.8	massive, black green, tuff, numerous 1mm calcite stringers CA-30-55o, broken contact, 1-2cm py blobs.								
113.86	114.27	massive, fg, void of stringers, soft, nil to weak magnetic LC sharp 44o.								
114.27	117.58	fragmental tuff varying degree of bedding.								
117.58	138.93	Laminated Ultramafic Tuffs								
117.58	118.95	more massive with well devel. schistosity numerous 1-3mm qc st II to schist ground contact.								
118.95	119.45	fragmental trace pyrite.	43156	118	119.45	1.45	0.01			
119.45	119.94	white opaque qv with grey green ultramafic inclusions usually with carbonate contacts 65o + 35o.	43157	119.45	119.94	0.49	0			
119.94	120.33	fragmental to locally brecciated, qtz stringers to 120.06, irregular contact.	43158	119.94	120.33	0.39	0			
120.33	122.11	black green fragmental tuff contorted bedding and x-cut by 1 1/2-3cm white qv with chlorite and talcose, moderately soft, non magnetic, qtz st.	43159	120.33	121	0.67	0			
			43160	121	122.11	1.11	0			
122.11	123.81	FP, fg medium to dark grey matrix with 1-2mm white plagioclase phenocryst, hard, massive uniform, siliceous, scattered to 3mm qtz st random CA-10, 20, 30, 40, 60, minor chlorite, white 1/2cm qtz veinlets CA- 87o at 122.67, 123.37-123.81 pale grey matrix due to altn from qtz st, qtz flooding and irregular chalk white qtz masses, 123.70-123.81 inclusions of chlorite, trace sulphides UC with 3cm of massive recrystallized chlorite, LC irregular CA sinuous 50o and massive recrystallized chlorite 123.81-123.85.	43161	122.11	123	0.89	0.01			
			43162	123	123.81	0.81	0			
123.81	124.64	UM, 123.81-124.28 hard silicified black green UM very chloritic with white qtz flooding, irregular stringers and masses, LC 60o, 124.28-124.64 fragmental to tuffaceous dark green moderately hard, qtz flooded, trace to scattered vfg pyrite LC broken ground.	43163	123.81	124.64	0.83	0			
124.64	124.88	felsic dike, aphanitic dark grey to blackish, altn due to qtz stringer to greenish brown, hard, siliceous, massive uniform, with chlorite fragments and inclusion chlorite ff trace vfg py, LC 85o.	43164	124.64	124.88	0.24	0			
124.88	127.96	fragmental tuff UM, fg, chloritic dark green matrix with light brown to grey buff fragments, overall light to medium grey, slightly broken from 124.88-125.63, locally crenulated bedding 126.33-127.0, few qtz st at 125.25 1cm CA 60-65 x-cut bed 60o 125.40 sinuous 1/2cm CA overall 60o x-cut bed, 125.55-125.74 grey white translucent with carbonated, 126.17 1/2-1cm grey translucent qtz st, CA sinuous deformed CA-30o to 60o LC.	43165	124.88	125.74	0.86	0.01			
124.88	127.96		43166	125.74	127.07	1.33	0.38	0.27		

Property: Hunter Mine		Hole No.	36		Sheet No.		4			
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
140.18	140.67	fg, tuff. Upper 10cm chocolate brown to medium brown, remainder light yellow green sericitic altn, patchy moderate to weak, trace to scattered py CA-50o.	43181	140.2	14.7	0.49	0.29			
140.67	141.41	grey green tuff fragmental light grey and black green lamination, minor kinkle scattered to <0.5% py, void of stringers, LC 50o.	43182	140.7	141.2	0.51	0.56	0.56		
141.41	141.48	grey translucent qtz veinlet with chlorite II and greyish carbonate LC-60o.	43183	141.2	142.4	1.18	0			
141.48	142.36	weak patchy sericitic altn with greyish crenulated tuff, scattered to trace pyrite, LC-70o, void of stringers.								
142.36	143.43	light grey fine grained to locally aphanitic contorted bedding, laminated locally occasional tuff, LC with qtz st irregular and x-cut bed LC 70o.	43184	142.4	143.4	1.07	0			
143.43	143.96	light grey green weakly sericitic altn folded with numerous 1-2 up to 1.5cm grey white opaque qtz stringers II to bedding, stringer at UC + LC irregular + x-cut bedding LC-70o.	43185	143.4	144.0	0.53	0			
143.96	144.35	light medium grey tuff, from 2-3mm qtz st II to bed.	43186	144.0	145.4	1.39	0.05			
144.35	145.35	light medium grey green weakly sericitic altn tuff 1cm white opaque qtz st at 144.57, 144.67, 144.92 all II to bedding.								
145.35	146.48	dark grey locally crenulated tuff, <0.5% vfg py, 145.56-145.71 qtz vein with dark green tuff inclusions and light brownish tan altn tuff, carbonated (ankirite) contact II to bed 60o-70o, 146.22-146.48 irregular qtz st 1-1 1/2cm x-cut bedding and stringers 3-5mm II to bedding, buff to tan altered inclusions, trace sulphides.	43187	145.4	146.5	1.13	0.01			
146.48	147	light grey green, weakly sericitic altn to weak moderate pervasive sericitic altn, local crenulations qtz st at 146.59 1cm 50-55o, x-cut bed, 146.85 1 1/4cm CA-75o near II to bed, 146.94-146.98 CA-60-70o irregular II and near II to bed with chocolate brown tourmaline ff, LC sharp CA-70o.	43188	146.5	147.0	0.52	0			
147	147.55	medium grey green, tuff few stringers II to bed.	43189	147.0	147.6	0.55	0.02			
147.55	150.42	bleached light grey due to white translucent qtz mass 147.65-147.71, veinlet	43190	147.6	148.0	0.45	0			
147.55	150.42	147.75-147.93 (curved UC 45o x-cut bed), grey to medium grey qtz veinlet	43191	148	148.57	0.57	0			
		148.32-148.57 contact 25-55o, grey white translucent st 148.80-148.97 II to 10o, qtz flooding 149.38-150.42.	43192	148.57	149.38	0.81	0.01			
			43193	149.38	150.42	1.04	0	0.01		
150.42	151.82	grey green tuff, crenulated, from 2-4mm grey qtz st II to bed, trace to <0.5% vfg py 151.44-151.57 light brown grey felsic dike, 2 stringers II to contact x-cut CA-40o, weak devel. of foliation contact slips CA 65o-75o LC irregular 60	43194	150.42	151.82	1.4	0			
151.82	153.3	qtz flooded, tuff to fragmental tuff, 151.82-153.30 patchy pervasive sericitic altn, hard to moderately hard, qtz carbonated stringers II to bed CA-60o x-cut 85o trace to <1% py vfg.	43195	151.82	153	1.18	0.01			
153.3	153.98	hard to very hard, intensely silicified numerous grey qtz veinlets stringers, pinkish brown, 153.54-153.56 and white 1cm qc veinlets II x-cut bedding at 55-60o, 153.93-153.98 grey opal qtz st 1cm CA-60o x-cut bedding cut by	43196	153	153.98	0.98	0			

Property: Hunter Mine		Hole No.	36		Sheet No.	5				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		1 1/2cm grey qtz stringer CA 65-70o which x-cuts bedding.								
153.98	154.47	qtz flooded light grey, 50% qtz trace to vfg scattered py LC-60o.	43197	153.98	154.41	0.43	0			
154.47	157.31	dark grey green to dark green, nil to weak patchy sericitic altn of crenulated with chlorite II slip planes CA-15-30o, scattered 2-5 1/2-1cm white grey	43198	154.41	155.27	0.86	0			
		opal qtz carbonated stringers II to and x-cut bedding CA-50o x-cut, 55 II to,	43199	155.27	155.91	0.64	0.01			
		75o x-cut with pyrite scattered mg py locally cg near vein 3 at 155.85, LC 53	43200	155.91	157.31	1.4	0			
157.31	157.64	numerous grey white translucent qtz stringer with white carbonated on margins, blobs masses and II to bedding.	35001	157.31	157.64	0.33	0			
157.64	158.38	patch to weak pervasive.	35002	157.64	158.38	0.74	0			
158.38	159.4	light grey crenulated well laminated tuff, trace pyrite, 159.28 1/2cm pyrite cube LC 56o.	35003	158.38	159.4	1.02	0.04	0.06		
159.4	160.53	weak pervasive sericitic altn, minor crenulation, few stringers II to bedding trace py LC 53o scattered pyrite.	35004	159.4	160.53	1.13	0			
160.53	161.01	siliceous with multiple grey siliceous bands chesty II to bed LC 50o sharp.	35005	160.53	161.01	0.48	0.01			
161.01	165	dark green to blackish green, fine laminations with light to medium grey, carbonated locally crenulations, to 162.0, dark green to 165.0, vfg py few py laminations, rare to void of qtz stringers, well level bedding CA 55o, 162.4	35006	161.01	162	0.99	0.02			
		65o at 164.3, 55o at 164.92.	35007	162	163	1	0.05			
			35008	163	164	1	0			
			35009	164	165	1	0			
165	165.36	weakly sericitic altn tuff carbonated.	35010	165	165.36	0.36	0			
165.36	166.79	medium to dark grey laminated rare qtz st LC bed CA 58o.	35011	165.36	166.79	1.43	0			
166.79	167.51	light to medium grey, tuff laminated 167.16-167.21 grey white qtz st with	35012	166.79	167.51	0.72	0.05			
166.79	167.51	1mm brecciated py band, 167.21-167.28 3-5% fg pyrite carbonated.								
167.51	168.26	siliceous, nil carbonated, weak patchy sericitic altn, crenulated tuff, scattered pyrite 1%, usually on slips forms of black green laminations, locally small crenulations.	35013	167.51	168.26	0.75	0.07			
168.26	170.1	light to medium grey, siliceous moderate pervasive sericitic altn, well laminated bed 60o rare stringers.	35014	168.26	169.2	0.94	0.01			
			35015	169.2	170.1	0.9	0			
170.1	171.25	patchy sericitic altn usually with few siliceous bands with medium grey to greenish medium grey sections, 170.43 1/2cm white opaque q st CA II to bed.	35016	170.1	171.25	1.15	0.03			
171.25	174.43	patchy moderate strong pervasive altn to locally pervasive sericitic altn, void of stringers, siliceous non carbonated, tuff, scattered pyrite, LC 55o 172.84	35017	171.25	172.25	1	0			
		formation CA-25o displacement 1.5cm left band, downhole, 172.97-173.47	35018	172.25	173.23	0.98	0.09			
		elongated greyish qtz discontinuous stringer II to bed, 173.24-173.47 1/2cm curved stringer x-cut bed.	35019	173.23	174.43	1.2	0.61	0.56		
174.43	176.11	patchy sericitic altn, 2 qtz stringer at 174.66 1.2cm II to bed, 175.18 boutinga II to bed 175.81-175.92 brecciated chloritic healed x-cut bed 35o, Bx-25o, 175	35020	174.43	175.57	1.14	0			
		bed 45o LC-25o x-cut bed by qtz vein.	35021	175.57	176.11	0.54	0.08			

Property: Hunter Mine		Hole No.	36		Sheet No.	6				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
176.11	176.15	white cloudy to opaque qtz vein CA-25o.	35022	176.11	176.61	0.5	0.3	0.33		
176.15	176.61	chloritic black green qtz rich laminated tuff scattered to <1% vfg py LC with qtz st 30-40o irregular.								
176.61	176.77	foliated to bedded, medium brown felsic, hard with irregular glossy grey white qtz mass, LC 35o 0.5-1% vfg py.	35023	176.61	177.05	0.44	0.19			
176.77	176.93	massive medium brown with small 1mm white grey plagioclase x-cut, some buff elongated angular fragment LC sharp CA 40o.								
176.93	177.05	same as 176.61-176.77 tuffaceous felsic sharp LC 35o with altn chloritic laminations 1mm.								
177.05	181.82	light grey to grey green weakly pervasive sericitic altn, rare greyish qtz stringer masses fragmental tuff to tuff, 178.70-178.74 whitish irregular mass, 179.28-179.78 scattered greyish white 3-5mm stringer usually x-cut bedding, 179.78-pinkish orange qc veinlet with pink calcite, trace sulphides, contacts 40-35o irregular, 179.98-180.43 qtz flooded fragmental tuff, scattered py, 180.43-181.82 grey green laminated minor crenulated tuff bed 40o, 180.55 2cm qtz veinlet CA-45o sinuous, 180.86-181.70 crenulated with 1-2mm chlorite ll slips, 181.59 3/4cm qtz st grey translucent CA-75 x-cut bed, 181.71 1cm greyish	35024	177.05	178.25	1.2	0.13			
			35025	178.25	179.29	1.04	0.22			
			35026	179.29	179.78	0.49	0			
			35027	179.78	179.98	0.2	0.07			
			35028	179.98	180.43	0.45	0.03			
			35029	180.43	181.82	1.39	0			
177.05	181.82	white translucent CA-70o LC 35o lost 10-12cm hard, baked.								
181.82	182.4	feldspar porphyritic, fg, medium grey matrix with 1mm white to whitish grey plagioclase phenocryst, massive uniform, hard siliceous, vfg pyrite scattered, chlorite py ll, qtz st ll, 181.84 1-2mm grey white ll qtz st deformed 182.10 3-4mm grey white translucent straight q st 65o LC sharp 50o.	35030	181.82	182.4	0.58	0.02			
182.4	187.54	tuffaceous fragmental, grey green, hard, siliceous, small pale buff fragments, scattered 3mm grey white qtz st to 1cm and 3cm qtz mass, 182.97-183.0 cloudy white qtz mass irregular x-cut bed, 183.44 1cm grey white translucent qtz st CA-35-40o irregular x-cut bedding 50-55o, 183.56-183.63 irregular grey white qtz stringer and mass, 184 increasing tuff and less fragments, 184.70-186.0 local crenulations and 2-3 chlorite ll slips planes, 186.03-186.06 light brown to medium brown felsic laminated fine, tuff, CA-40o, 186.06-187.54 tuff fragmental, light grey and black green, laminations, trace pyrite, moderately hard, siliceous, 187.0-187.05 white qtz st 1cm discontinuous 1cm straight CA 80o near ll to bed, 187.23-187.52 irregular 2-4mm grey qtz st near ll to CA overall 60o x-cut bed, 187.45-187.52 whitish grey qtz veinlet with inclusions CA 50o, 187.52-187.54 massive chlorite contorted altn, baked contact, LC-50o.	35031	182.4	183	0.6	0			
			35032	183	184	1	0.04			
			35033	184	185	1	0.03			
			35034	185	186	1	0.04			
			35035	186	187	1	0.07			
			35036	187	187.54	0.54	0.03			
187.54	189.27	felsic dike, aphanitic to fg, light grey to medium grey, hairlike qtz stringer hydrofracturing, monor chlorite ff, massive, uniform, irregular siliceous, non	35.37	187.54	188.4	0.86	0.01			
			35.38	188.4	189.27	0.87	0			

Property: Hunter Mine		Hole No:	36		Sheet No.	7				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		magnetic, few scattered 2-3mm grey opaque translucent qtz st, usually at 25-40o, few discont. white qc st vfg to fg py trace to scattered <0.5% LC 55o with 1cm black green chlorite baked contact.								
189.27	193.16	fragmental tuff, black green, moderately hard, few irregular 1-1.5cm white qtz st ll to bedding, vein few qtz stringers CA-45o, well develop bedding 190 at 43o, 191 at 47o, 192.6 at 70o, 192.84-193.16 folded kinkled bedding, medium light grey and black chlorite laminations, trace to nil sulphides, LC 50o contact baked chlorite band.	35039	189.27	190.75	1.48	0			
			35040	190.75	192.25	1.5	0			
			35041	192.25	193.16	0.91	0			
193.16	194.02	felsic dike foliated, aphanitic to fine grained, massive light brownish grey 193.16-193.25, foliated medium to dark grey foliated 50o 193.25-193.83, light to medium brown 193.83-194.02, colour due to chlorite qtz carbonated irregular stringers, hard, siliceous rare sulphides, numerous 2-5mm qtz grey opaque and grey white translucent stringers ll to foliation or near ll to foliation	35042	193.16	194.02	0.86	0.04			
193.16	194.02	CA 45o, chlorite ll with 1/2cm qtz stringers 30-33o x-cut foliation, 193.47-193.50 grey opaque qtz veinlet with scattered pyrite, CA 75o, 193.82 1cm grey white with pyrite in altn sections CA-30-35o LC CA-45o.								
194.02	197.4	fragmental tuff, same as 189.27-193.16, few stringers, light grey green to medium grey, few grey white stringers 195.76-196.80 overall trace to scattered py, 196.90-197.40 qtz flooding with scattered vfg py LC 40o.	35043	194.02	195.5	1.48	0			
			35044	195.5	196.9	1.4	0.01			
			35045	196.9	197.4	0.5	0			
197.4	203.82	felsic tuffaceous fragmental, aphanitic to fine grained, weakly to moderately fine laminations silicified to 197.81, carbonated weakly to moderately 197.81-203.37, rare distinct greyish cloudy white qtz stringers at 198.0 CA-40o ll to bed CA-36o at 198.05, 199.0 CA-20o 2-3mm 203.41 CA-55o, 203.47 CA-45o both 4mm, numerous hairlike to 1mm calcite carbonate ll stringers ll to bed, locally laminations fg ophanitia difficult to determine tops, local crenulations from 201.83-202.43 more tuffaceous 202.43-203.82 fragmental tuff with irregular crispy qtz carbonated stringers hairlike, 202.73-203.82 scattered 1-2% overall pyrite with scattered pyrrhotite 203.06-203.37 as crenulation masses up to 2-3% associated with greyish qtz carbonated between fragments LC-45o, 203.37-203.62 silicified, hard, few stringers, silicified grey brecciated zone 3-5% pyrrhotite with up to 1% pyrite associated with brecciated fragmental, trace to <0.5% with greyish opaque cloudy siliceous and rare pyrite in stringers, LC 70o, 203.62-203.82 buff to buff green fragment moderately hard to hard, rare to scattered fg pyrite, LC 70o gradational colour change lust 2cm to buff blackish green, conformable contact.	35046	197.4	197.77	0.37	0.06			
			35047	197.77	199.27	1.5	0			
			35048	199.27	200.27	1	0			
			35049	200.27	201.4	1.13	0			
			35050	201.4	201.83	0.43	0			
			35051	201.83	202.43	0.6	0.01			
			35052	202.43	202.73	0.3	0			
			35053	202.73	203.37	0.64	0.11			
			35054	203.37	203.62	0.25	0.1	0.07		
			35055	203.62	203.82	0.2	0.03			
203.82	207.2	UM fragmental tuff silicified, fg, black green, hairlike qtz flooding stringer ff locally well laminated tuff sections within fragmental tuff, hard, silicified, qtz	35056	203.82	205.1	1.28	0			
			35057	205.1	206	0.9	0.01			

Property: Hunter Mine		Hole No.	36		Sheet No.	8					
Meterage		Description	Sample			Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check	
		flooding 204-205.08, scattered to trace pyrite associated with qtz scattered to trace, few distinct qtz str, LC gradational, moderately soft, 205.08-205.34 same as above but weakly to moderately carbonated to 205.34, locally siliceous due to grey opaque qtz masses and veinlets 205.72-206.0 weakly carbonated 206-209.85, 203.82-205.10 fragmental tuff, 205.10-207.20 qc flood weak carbonate flooding moderately carbonated fragmental tuff.	35058	206	207.2	1.2	0				
207.2	209.07	brecciated and qtz flooded, massive, tuff, fragmental, 206.91-206.97 qc vein CA-80o, 207.04-207.21 tuffaceous, 207.21-209.07 massive, with porphyritic texture carbonated x-cuts.	35059	207.2	208	0.8	0				
			35060	208	209.07	209.07	0.07				
207.2	209.07	brecciated and qtz flooded, massive, tuff, fragmental, 206.91-206.97 qc vein CA-80o, 207.04-207.21 tuffaceous, 207.21-209.07 massive, with porphyritic texture carbonated x-cuts.	35059	207.2	208	0.8	0				
			35060	208	209.07	209.07	0.07				
209.07	209.85	tuffaceous fragmental qtz flooded weakly carbonated moderately soft.	35061	209.07	209.85	0.78	0				
209.85	211.6	very strongly carbonated porphyritic texture massive intensely qc veining usually CA-55o II to foliation schistosity, qtz flood brecciated from 210.83-211.12, qc vein 211.12-211.21 CA-50o x-cut schistosity, 211.21-211.60 very carbonated tuff to tuff fragmental.									
211.6	212.86	weakly carbonated, carbonate stringers, tuff to tuff fragmental intensely vein, LC 50-55o.									
212.86	219.38	1mm white grey carbonate phenocryst massive with locally veining to brecciation angular fragments, 213.27-215.76, 216.58-218.40 non porphyritic locally porphyritic.									
219.38	220.3	carbonated flooded non porphyritic.									
220.3	221.12	tuff fragmental.									
221.12	221.34	crenulated tuff.									
221.34	221.41	massive tuff, conformable contact CA 65o.									
221.41	234	Argillite Graphitic and Grey Arenites									
221.41	223.67	graphitic argillite, aphanitic to fg, black, fine laminations, well devel. bedding 50o, massive, rare hairlike II stringers usually II to bed to 35o x-cut bed at 221.57-222.20 pink calcite vein 221.61 2 1/2cm CA028-30o x-cut bedding.									
223.67	226.98	banded argillite with minor graphitic argillite dark grey to black laminations, 224.0-224.41 glassy translucent qc veinlet greenish inclusions CA-50o II to bed and broken. 224.41 1 1/2cm qc veinlet II to bed, 224.56-225.10 irregular qc veining x-cut bed, 225 bed 38o-40o, 225.10-225.36 kinkle folding to crenulations, 225.36-225.92 crenulation bedding change to vein II to CA then to 50o numerous slip planes CA 25-40o, 225.92-226.16 crenulated graphitic argillite, 226.16-226.43 qtz vein with graphitic argillite siliceous, CA-30o	35062	224	224.8	0.8	0				
			35063	226	226.4	0.4	0.01				

Hunter Mine - Diamond Drill Log HM-04-37

Property:	Hunter Mine	Hole Dip:	-66	Page No.:	1 of 13		HM-04-37			
Location:		Hole Azimuth:	105	Date Started:	July 20,04					
Claim No:	HR 1009	Hole Length:	248.81m	Date Finished:	July 28/04					
Elevation:	Porcupine Lake	Purpose:		Drill Co.:	Benoit					
UTM Coords.:	5370768.3N, E487021.0			Logged by:	K. Jensen					
Meterage		Description	Sample				Assays			
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
0.0	63.0	Casing								
63	70.5	Ultramafic Talcose Rocks								
		ultramafic, 63.0-65.33 fg schistose, local shearing, schist CA-30o, tuffaceous, carbonated stringers, soft, talcose, 65.33-68.78 more massive, locally schistose carbonated qtz stringers 10-20o to II to CA, moderately hard, 68.78-70.5 crumbly brecciated sheared zone, void of stringers, only small brecciated fragments of qtz, 69.0-72.0 only 1.33m lost core 1.67m.								
70.5	77.82	fragmental tuff, scattered qtz carbonated fragments of stringers, minor stringers boudinage and near II to CA same as bedding schistose, local sections with bed at 15-20o, moderately soft to soft, talcose, scattered fg mg py, LC 28o at 1.3cm qtz c veinlet CA-28o.								
77.82	79.45	massive, porphyritic texture, nil to very weak devel. of schistose, rare qc stringers 1cm CA-35o, 78.44-78.60 irregular mass of carbonate + talcose LC-50o irregular								
79.45	82.31	massive with local brecciated sections and appearance of polysuturing, randomly orientated carbonate healling stringers with brecciated sections, trace to scattered pyrite LC irregular.								
82.31	105	brecciated healed with creamy pale greenish carbonate, 85.94-86.60 intense shearing 40o with small section of breccia, 87-93 2.99m lost core sections of brecciated and intensely sheared and crushed zone, 93.50- 94.22 crushed zone, 94.22-95.31 schistose, probably tuffaceous fragmental bedding CA-40o, 95.31-98.84 brecciated tuff fragmental with intense shearing and crushed zone, local slick on side, 98.84-103.0 brecciates carbonate healed, intense random veining, 103-103.47 massive weak level of schistose CA-55o trace pyrite, 103.47-105.0 brecciated scattered mg cg py in veining.								
105		Laminated Ultramafic Tuffs								
105	105.8	tuff fragmental stringer carbonated veining 2-5mm all II well level of schistose bed CA-40o LC 40o scattered 1-3mm pyrite masses in veining.								
105.8	108	brecciated fragmental.								
108	110.51	tuff to tuffaceous fragmental, local kwinkle folding 108.0-108.18, local intense st carbonated probably due to laminations 108.23-110.51, contorted schist bedding								
108	110.51	carbonate fragmental stringers, 109.64-109.68 intense searing CA-60o.								

Property		Hunter Mine		Hole No.	37		Sheet N	2				
Meterage		Description		Sample			Assay					
From	To			No.	From	To	Width	Au (g/t)	Au Check	Au (2nd)	Au check	
110.51	111.48	massive non magnetic, few 2-3 white carbonated stringers usually CA-70-80o moderately soft, trace sulphides, LC 50o with qc veining 1.5cm with 2-3% mg cg py minor chalcopyrite and traces of pyrrhotite.										
111.48	113.29	fg, weak to moderately level of bedding, black to black green, tuff, random orientated qc st irregular, contorted, few S folds, no sections to HCC trace sulphides.										
113.29	113.43	qc vein with inclusions UC grade LC irregular.										
113.43	113.9	more massive weak devel. of bed random orientated 2-5mm qc st kinkle fold.										
113.9	118.37	vfg to fg, tuff local small crenulations, scattered small fragments, random stringers, 113.90-115.41, afterwards majority 1-2mm ll to bedding and apperance of fg py to small 2mm blobs, locally 115.75-116.15 3-5% vfg py, 116.84-117.0 3-5% mg, 117.0-118.37 1-2% vfg py, stretching to have small fragments.		35065	115.4	116.15	0.75	0.04				
				35066	116.15	116.84	0.69	0				
				35067	116.84	118.37	1.53	0.04				
118.37	120	tuff to tuffaceous fragmental medium grey to greenish tint, locally crenulations 119.30-119.64, 119.64-120.0 bedding near ll to CA contorted, occasional stringers near ll to bed 60o.		35068	118.37	119.64	1.27	0.03				
				35069	119.64	120.8	1.16	0				
120	122.04	10-15o qtz st white opaque, 1-2mm up to 1-1 1/2cm random orientated from near ll to bedding to 40o-60o x-cut bedding.		35070	120.8	122.04	1.24	0				
122.04	123.07	high siliceous contact, fragmental void of stringers, brecciated with black green chlorite ff, trace pyrite LC sharp 50o.		35071	122.04	123.07	1.03	0				
123.07	123.22	fg grey green tuff, 1/2cm qc stringer CA-40o at 123.13.		35072	123.07	124.23	1.16	0.03				
123.22	124.23	similar to 122.04-123.07, fragmental LC-65o x-cut by 1cm grey qtz st CA-50o 124.04-124.10 qtz vein V shaped with inclusions CA-55o.										
124.23	126	fg tuff, light greenish light medium grey well devel. bedding small crenulations, locally contorted with scattered 1/2-1cm grey translucent qtz st, CA-20-25o x-cut bed CA-55o at 124.5, 65o at 125.3 80oat 125.80.		35073	124.23	125.07	0.84	0.01				
				35074	125.07	126	0.93	0				
126	126.24	qtz flood vein with inclusion, silicified scattered 1-4mm pyrite masses to blocks contact sharp 65 + 70-75o.		35075	126	126.23	0.23	0.02				
126.24	126.88	tuff as 124.23-126.0.		35076	126.23	126.88	0.65	0.01				
126.88	127.13	white qtz vein with talcose medium grey green inclusion trace py CA-60-70o.		35077	126.88	127.13	0.25	0.09				
127.13	132.24	light grey to medium grey green, laminated with grey siliceous and altered chloritic tuff fragmental with minor crenulated tuff 127.82-127.92, 2-4 qtz grey stringer per meter, from ll to bed to x-cut bedding locally contorted 127.33		35078	127.13	128.55	1.42	0.12	0.17			
				35079	128.55	129.95	1.4	0				
127.13	132.24	1/4cm CA-15o, 127.47 3cm mass, 127.50 ll to CA to 10o, 127.93 CA 70o, 128.12-128.16 qtz mass, 128.61 20o 1cm, 129.04-129.09 qtz mass, 129.23 1cm CA 55o, 129.26 1 1/2 CA 30o, 129.97-130.13 buff white qtz vein irregular 50o with inclusions V shaped x-cut and near ll to 50o, 130.85-131.0		35080	129.95	131	1.05	0.01				
				35081	131	132.24	1.24	0				

Property		Hunter Mine	Hole No.	37	Sheet N	3				
Meterage		Description	Sample		Assay					
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		crenulated tuff, 131.31 1/2cm grey qtz st CA-20o x-cut bedding 70o, 131.69 1cm tops to 1/2cm grey qtz st CA 30o-20o x-cut bedding.								
132.24	133	massive, nil to very weak schistosity, weakly porphyritic with fg chlorite x-cuts, x-cuts by random orientated contorted grey white opaque qtz stringer from 1-4mm, trace to nil pyrite LC 75o angular UC 50o irregular.	35082	132.24	133	0.76	0			
133	134.17	qtz flooded fragmental fragments, nil to weak schistosity contorted qtz st, grading from fg tuff 133.0-133.20 to fragmental scattered to <0.5% fg pyrite, LC 20-30o contacted.	35083	133	134.45	1.45	0			
134.17	134.45	fg, kacki green brecciated fragments with chlorite healed, 1% vfg fg pyrite.								
134.45	135.4	massive to brecciated massive flow, 134.45-134.67 massive, few 2-3 qtz st random, 134.67-135.30 massive brecciated large fragments filled with qtz, 135.30-135.40 massive brecciated small fragments qtz filled, LC 65o.	35084	134.45	135.4	0.95	0	0.01		
135.4	138.76	fragmental to fragmental tuff, 135.40-135.47 appears to be rubble zone small qtz fragments 4-7mm contorted schist, 135.47-138.76 qtz flooded, brecciated tuff fragmented contorted schistosity bedding, grey green to olive green, locally more intense qtz flooding, 137.90-138.76 larger fragments possible Bx massive flow.	35085	135.4	136.77	1.37	0			
			35086	136.77	138.2	1.43	0			
			35087	138.2	139.53	1.33	0			
138.76	139.38	qtz flooding decreasing tuff fragmental with crenulated bedding with chlorite ff slip planes near II to CA.								
139.38	139.53	buff to light tan tuff and felsic dike internal fractured and filled with black green chlorite, interlayered into chlorite, fragmental tuff, LC sharp 47o UC 50o.								
139.53	140.24	felsic dike, aphanitic to light tan at contacts to medium brown, heavy hydrofracturing chlorite II massive uniform, very hard siliceous, scattered 1% to disseminated 1% py in dike, minor py associated with chlorite II, rare white hairlike qtz II, random, more intensely chlorite II st lower contact to bottom 1/2, LC 75o.	35088	139.53	140.28	0.75	0.02			
140.24	140.28	chlorite tuff and 1cm qc st CA-75o.								
140.28	140.73	foliated felsic fragmental tuff, fg, buff with chlorite matrix massive uniform	35089	140.28	140.73	0.45	0.02			
140.28	140.73	weak to weak moderately devel. foliation CA-40o, LC 15-32o, UC 60o, 6 3mm grey qtz st from 140.48-140.73 CA-70, 60, 90o, trace sulphides, hard, siliceous.								
140.73	141.21	chlorite tuff fragmental, 140.73-141.01 fragmental very chlorite, 141.01-141.21 tuff crenulated, chlorite II slips II to CA contacts 50o irregular LC 40o overall.	35090	140.7	141.2	0.48	0			
141.21	141.38	foliated felsic fragmental tuff same as 140.28-140.73, elongated stretched buff fragments near UC, few hairlike chlorite ff with py, UC kinkled 3mm qc st x-cut contact, 141.30 3mm grey qtz st CA-65o x-cut bed CA-45o better bedding than above, LC 43o II to chlorite tuff fragmental.	35091	141.2	141.4	0.17	0			
141.38	147.32	tuff fragmental, same as above, grey green to locally buff grey green, 141.38-	35092	141.4	142.5	1.12	0			

Property: Hunter Mine		Hole No.	37		Sheet N	5				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		154.17-154.21 grey translucent qtz and ankerite CA-50o-60o, 154.48-154.50 grey translucent qtz and askinete CA-80o, 154.74 LC sharp CA-40o.								
154.74	157.83	tuff to tuffaceous fragmental, as above grey medium to grey light greenish grey, locally pale brownish tint, 154.74-155.27 tuff, locally crenulated, 155.27-156.26 fragmental tuff well bedded CA-40-45o with 1/2cm grey qtz st II to bedding, minor crenulations with qtz st II to crenulations, rare x-cut qtz st, 155.72 1cm CA-80o, trace py, 156.26-157.40 tuff fragmental with contorted qtz st II and x-cut bedding numerous stringers with medium grey, 157.40-157.83 qtz flood medium grey fragmental tuff.	35108	154.74	155.27	0.53	0			
			35109	155.27	156.26	0.99	0			
			35110	156.26	157.4	1.14	0			
			35111	157.4	157.83	0.43	0			
157.83	160.55	medium grey tuff to fragmental, rare qtz stringers, trace to scattered vfg py st II to bedding CA-40o at 158.3, 45o at 159.7 weak moderate devel. of bedding, 158-158.86 contorted and qtz grey flooding layer fragments, 158.86-160.30 fragmental tuff, 160.30-160.55 well bedded laminated tuff, tuff fragmental crenulated with 2mm chlorite II slip planes, bed 45-50o, 160.55 LC with qtz veinlet CA-80o x-cut bed.	35112	157.83	158.5	0.67	0			
			35113	158.5	158.86	0.36	0			
			35114	158.86	159.7	0.84	0			
			35115	159.7	160.55	0.85	0			
160.55	161.33	qtz flooded fragmental tuff with distinct qtz ankerite veinlets at 160.55-160.59 CA-80o 160.84 1/2 CA-75o, 161.12-161.15 grey with whitish x-cuts II to bedding CA-40o, 161.33 contact sharp CA-30o.	35116	160.55	161.33	0.78	0			
161.33	162.8	fg well bedded laminated pale grey green weakly sericitic pervasive altn, with numerous 1mm grey ff st to 161.90 x-cut bed CA 50-60o 161.90-162.80	35117	161.33	162.8	1.47	0	0		
161.33	162.8	few greyish qtz st 1/2cm x-cut bed ranges from 65o at 161.95, 40o at 162.56 40o discont. at 162.66 all translucent, grey white qtz ank at 162.50 1cm CA 80o, 162.78-162.80 baked contact CA-50o sharp.								
162.8	167.08	QFP, aphanitic light brownish matrix 2-4mm milky white plagioclase and grey opaque qtz phenocrysts, massive uniform, very hard, siliceous, fragmental and filled with chocolate brown siliceous tourmaline, locally rare intense, locally matrix medium grey unaltered 163.65-163.87, 164.87-165.37, inclusions of grey green tuff fragmental from 163.62-163.66 CA-60o qtz stringers 163.20-163.62 usually 1/2 grey translucent CA-25o in both direction to 20o with intensely brown tourmaline breccia healed, scattered to disseminated vfg 1-2% pyrite, 164.62-164.85 intensely fractured, 165.66-166.07 intensely fractured, 164.58 1/2 grey qtz st x-cut intensely fractured CA-45o, 166.60-167.08 intensely fractured, 164.59-167.08 void of large qtz stringers except ff, 1mm CA-25o, 167.08 sharp contact CA-35o.	35118	162.8	163.2	0.4	0.04			
			35119	163.2	163.62	0.42	0.03			
			35120	163.2	164.5	1.3	0.02			
			35121	164.5	165.64	1.14	0.01			
			35122	165.64	166.6	0.96	0			
			35123	166.6	167.08	0.48	0.02			
167.08	245.42	Exhalitic Tuff and Sedimentary Rocks								
167.08	168.35	tuff to fragmental tuff, fg, pale green massive tuff x-cut by several siliceous medium brown felsic dikelets 167.27-167.30 45o, 167.39-167.48 brecciated	35124	167.76	0.68	0.11				
			35125	168.35	0.59	0.1				

Property: Hunter Mine		Hole No.	37		Sheet N	6				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		tuff irregular and 47o, 167.62-167.76 V shaped with tuff fragment inclusions CA-40-60o, 167.76-168.35 pale grey green tuff, few stringers II to bedding, very weak to weakly sericitic altn, 168.28-168.35 1cm qtz stringer CA 60o with Bx fragmental tuff altn, medium brown 168.39-168.35 LC 55o.								
168.35	169.02	QV, aphanitic light grey matrix intensely fractured filled with chocolate brown siliceous tourmaline, x-cut by grey white randomly orientated 3-4mm stringers qtz, scattered buff white ankerite x-cuts, scattered fg pyrite 0.5-1% LC at 169.02 CA-45o.	35126	168.35	169.02	0.67	0.01			
169.02	169.29	QFL Bx tuff, fg, laminated tuff yellow brown to buff green altn, qtz flooded and brecciated x-cut by white q schist irregular stringer, trace py LC 40o curved.	35127	169.02	169.29	0.27	0.76			
169.29	170.02	carbonated fragmental tuff, fg, buff yellowish green to pale green, small and inclusions size fragments, massive tuff, to pale grey in qtz flood silicified sections, weakly carbonated, hard, trace to <0.5% LC 45o sharp.	35128	169.29	170.02	0.73	0.02			
170.02	171.31	weakly pervasive sericitic altn and qtz flooded fragmental tuff pale green, minor fuchsite, pale yellow green rare distinct qtz stringer or fragments CA-55o original matrix of fragmental lapilli tuff with small blackish grains, scattered mg pyrite <0.5-1% locally LC gradural.	35129	170.02	171.31	1.29	0.05			
171.31	171.74	weakly pervasive sericitic altn tuff, moderately level of bedding, numerous layers with 2-3% fg pyrite, void of stringers except 171.56-171.74 1/2 greyish with scattered pyrite, near II to CA, deformed kinkle folded LC 55o sharp.	35130	171.31	171.74	0.43	0.84			
171.74	172.28	fg, grey green to pale green 2-7mm laminations, minor grey silicic bands, scattered vfg py <0.5%, 2 1/2cm white opaque qtz st at 172.17, CA-50o II to bed 55o at 171.85, 50o at LC at 172.28.	35131	171.74	172.28	0.54	1.18	1.13		
172.28	172.78	weakly pervasive sericitic altn, well laminated with grey qtz banding, few grey kwinkle qtz st x-cut bed, scattered vfg py, py band 172.65 3-5% pale grey green to light green, tuff, siliceous and silicified, LC 50-52o.	35132	172.28	172.78	0.5	0.72			
172.78	173.43	weakly patchy sericitic altn, light grey pale green, medium green, grey white silicin banding, greyish medium brown kinkle qtz stringer near II to CA, irregular greyish qtz mass at 173.02-173.05, several disseminated pyrite bands 0.5-1.5cm 2-3 to 7-10% fg py associated with pale green bands, LC sharp 52o.	35133	172.78	173.43	0.65	0.89	1.16		
173.43	173.91	patchy moderately sericitic altn to pale buff to buff light yellowish green, trace to locally 1% fg py grey white translucent qtz st at 173.52 1.5cm CA-55o near II to bedding LC sharp CA-35o.	35134	173.43	173.91	0.48	0.07			
173.91	174.81	same as 173.43-173.91, qtz flooded minor fragmental, majority II to bedding few grey white translucent massive 174.0-174.07 qtz veinlet with inclusions	35135	173.91	174.81	0.9	0.03			

Property: Hunter Mine		Hole No:	37		Sheet N	7				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		and minor fuchsite trace py, locally 174.61 mg py in band LC 55o.								
174.81	175.3	medium grained fragmental tuff, massive weak moderate devel. of bedding, trace py, grey green to medium green, LC broken.	35136	174.81	175.3	0.49	0			
175.3	176.51	qtz flooded, moderate patchy sericitic altn fragmental tuff, with few pinkish tan felsic fragments at 176.29-176.41, increasing chlorite downhole as brecciation distinct qtz veinlet at 175.30-175.54 grey trace white opaque ank CA irregular at 55o irregular, trace py ng.	35137	175.3	176.51	1.21	0.03			
176.51	177.98	same as above, but moderate pervasive sericitic altn, qtz flooded fragmental tuff with few pale pinkish buff felsic layers 176.65-177.04, q flood usually ll to bedding, scattered black 1mm q eyes with tuff grade contact CA-40o sharp.	35138	176.51	177.98	1.47	0.04			
177.98	179.38	moderate patchy to pervasive sericitic altn, numerous greyish translucent qtz stringers 1/2-1cm ll to bedding few with fuchsite, ankerite common with st usually CA-40o, rare greyish qtz st x-cut bed CA-65o, LC sharp CA-52o scattered py fg mg locally <0.5%.	35139	177.98	179.38	1.4	0.02			
179.38	180.62	aphanitic to fine grained pale pinkish grey to pale greenish light grey massive sections up to 30cm with very fine laminations with few stretches of fuchsite grey to grey green tuff sections from 179.76-180.02 and 180.32-180.62, bedding CA-42o at 179.55 pale pinkish green to 52o at 180.45, trace to nil py void of stringers.	35140	179.38	180.62	1.24	0.08			
180.62	180.91	aphanitic blackish grey siliceous matrix with chlorite, brecciated fragments grey green to buff lower portion 180.77-180.85 siliceous brownish matrix 180.85 contact CA-75o sinuous, 180.85-180.91 tuffaceous fragmental breccia moderately sericitic, LC 75o.	35141	180.62	180.91	0.29	0.08			
180.91	182.07	fg, pale buff to light tan tuff, trace to scattered vfg py, good level bedding CA 47o, scattered qtz stringers 181.07-181.18 white opaque qtz veinlets CA-60-65o with inclusions, 181.34 1/2 cm x-cut bed, 75o, 181.62 1cm with minor fuchsite CA 50-60o, 181.62 1cm white qtz st CA 80o x-cut and termination 1cm pale greenish white grey qtz st CA-32o 182.01-182.07 greyish white opaque with faint green tint CA-60o-45o LC-45o.	35142	180.91	182.07	1.16	0.01	0.01		
182.07	183.63	patchy weak sericitic altn, laminated buff and pale brown tuff to grey green, scattered pyrite void of stringers bed CA-50o LC 53o sharp.	35143	182.07	182.93	0.86	0			
			35144	182.93	183.63	0.7	0			
183.63	185.11	patchy weak moderate to moderate sericitic altn laminated buff, pale brown and medium green, minor kinkle folding at 184.07, scattered 1-2% pyrite with buff grey band at 184.24 1cm, scattered fg py, qtz stringer at 183.76 1 1/2cm ll to bed CA-60o, 184.03 1cm qtz st CA-80o 184.39-184.56 greyish white opal with milky white qtz and inclusions with vfg py CA-30o and 60-70o irregular, 184.91-184.97 irregular grey white qtz st CA 45-55o py fg to mg at	35145	183.63	184.39	0.76	0			
			35146	184.39	185.11	0.72	0			

Property: Hunter Mine		Hole No.	37		Sheet N	8				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		lower contact.								
185.11	188.82	moderate patchy sericitic altn, tuff, fragmental tuff to 185.51, 2cm very qtz st cloudy with sericitia CA-60-55o, local sections qtz flooded, 185.51-186.0	35147	185.11	185.51	0.4	0.12	0.16		
		fragmental numerous qtz II to bed boudinage, scattered py, 185.84 3cm	35148	185.51	186	0.49	0			
		blackish qtz st x-cut bed with vfg py CA irregular 60o-70o, 185.86-186.0	35149	186	186.71	0.71	0.01			
		approximate 2-3% fg mg pyrite, 186.25-186.27 grey irregular qtz flooding with	35150	186.71	187.1	0.39	0			
		2-3% fg mg py, 186.34 chlorite altn to fuchsite 2mm band with pyrite, 186.78	35151	187.1	187.46	0.36	0.01			
		188.09 kink fold axis at 10o CA, 187.29-187.33 3/4-1cm band with 5-7% fg	35152	187.46	188.03	0.57	0.02			
		pyrite, 187.37-187.46 irregular sinuous qtz veinlet grey white with altn	35153	188.03	189.2	1.17	0			
		inclusions 0.5-1% vfg py CA-35-60o, 187.93-188.03 weak altn with greyish								
185.11	188.82	siliceous bands 5-7% fg py, CA-55o 188.82.								
188.82	189.2	weak to nil sericitic altn.								
189.2	192.94	weak patchy sericitic altn felsic tuff, tuffaceous fragmental. Local sections	35154	189.2	190.33	1.13	0.02	0.03		
		1-2% mg py, with scattered 10-20cm sections of chlorite tuff grey green	35155	190.33	191.04	0.71	0.09			
		medium green laminations 189.38-189.48, 43o, 190.73-191.04, 68o, 191.04,	35156	191.04	192	0.96	0.05			
		2-3% py, 189.29-189.33 V shape grey white qtz st 40o-65o II to bed, 190.35-	35157	192	192.94	0.94	0.43	0.42		
		190.42 grey white and sericitia and trace vfg py CA 60-80o, 191.19 grey								
		blackish qtz st CA 70o x-cut bed 2cm 1% py, 192.37-192.44 chlorite buff tuff								
		fragments 3-5% fg py CA-32-30o, 192.86 1 1/2cm white grey qtz st CA 50o								
		II to bed.								
192.94	194.42	fg laminated tuff, weak patchy sericitic altn, bedding good devel. CA 193.5 at 50o 194.3 at 50o.	35158	192.94	194.42	1.48	0.11			
194.42	194.81	light grey to pale brownish tint tuff.	35159	194.42	195.36	0.94	0.03			
194.81	196.53	nil to weak patchy sericitic altn, fg mg, uniform, moderate devel. of bedding, less distinct lamination, cleavage good CA-55o, scattered pyrite 195.36 with siliceous band, 195.82, 195.89, 196.23 1cm pink qtz CA-55o 196.53 LC-55o.	35160	195.36	196.53	1.17	0			
196.53	198.16	medium grey green laminated tuff, scattered py bands at 196.73, 197.22-	35161	196.53	197.16	0.63	0.02			
		197.25, 197.60-197.71 2-3% py, 197.18-197.62 scattered pinkish grey qtz	35162	197.16	198.16	1	0.06			
		pink carbonated st II to bedding 2-5mm, 197.53 5mm, 197.57 1cm, 197.73								
		1/2-1cm kinked grey white q st near II to CA with vfg pyrite on contact.								
198.16	198.97	light grey to light grey green, siliceous scattered to finely disseminated vfg py overall <1% locally 1-2% with band, LC 48o.	35163	198.16	198.97	0.81	0.09	0.1		
198.97	199.85	patchy weak sericitic altn tuff, pale greenish qtz st 199.04 1.2cm blackish qtz st CA-73-80o x-cut bed 55o scattered pyrite, 3-5% 199.03-199.04.	35164	198.97	199.85	0.88	0.07			
199.85	200.88	weak pervasive sericitic altn, tuff minor kwinkle fold, pale green q st II to bed.	35165	199.85	200.88	1.03	0.08			
200.88	201.47	medium grey to blackish green, tuff, locally qtz flooded 201.0-201.18, qtz st 2-3% py CA-65o irregular at 201.35.	35166	200.88	201.47	0.59	0.21			

Property: Hunter Mine		Hole No.	37		Sheet N	9				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
201.47	203.8	moderate sericitic altn to locally moderate patch sericitia, tuff to locally tuff fragmental, scattered vfg py, blackish qtz st at 201.77 1/2cm CA-70o x-cut bed 45o, qtz breccia vein 202.09-202.13 with 2-3% pyrite, 202.73 1cm grey Bx st CA 55o, x-cut fragmental bed 70o, 202.90 1/2cm grey brown qtz st 70o x-cut bedding.	35167	201.47	202.6	1.13	0.21			
			35168	202.6	203.8	1.2	0.31			
203.8	206.86	light grey to pale grey pale green tuff local kinkled folds, scattered vfg pyrite	35169	203.8	205.3	1.5	0.3			
203.8	206.86	<0.5% good bedding CA-57o, small fragments, 205.52-205.70 parts of pale grey opaque qtz st, appears ll to bed, 205.71 1.5cm grey opaque qtz st CA 60o, 205.93-206.07 irregular white opaque qtz masses, and grey opaque masses, 206.20-206.55 qtz flooded and white opaque, grey opaque qtz veinlets overall 65% veining irregular contacts and inclusions of grey tuff fragmental CA-40o 70o trace to rare sulphides, 206.55-206.86 fragmental tuff large fragments UM moderate level bedding CA-55-60o LC 50o.	35170	205.3	206.2	0.9	0			
			35171	206.2	206.55	0.35	0			
			35172	206.55	206.86	0.31	0			
206.86	208.9	qtz flooding with distinct grey opaque with whitish carbonate from 1-1.5cm usually ll to bed and white opaque qtz veinlets 5-9cm usually x-cut bedding in fragmental tuff, trace to <0.5% vfg py, 40-50% veining LC 45o with 2cm grey opaque qtz st.	35173	206.86	207.77	0.91	0			
			35174	207.77	208.45	0.68	0			
			35175	208.45	208.9	0.45	0.01	0.03		
208.9	209.73	grey green to dark green laminated tuff fragmental, nil altn, few 1/2 qtz ank st CA-60o ll to bedding x-cut by grey opaque 3mm st qtz CA-45o.	35176	208.9	209.73	0.83	0.03			
209.73	210.26	irregular grey white opaque qtz stringers and masses, 209.91-210.26 tuff fragmental olive green and medium green fragmental.	35177	209.73	210.26	0.53	0			
210.26	210.75	qtz flooded and white qtz veinlets and opaque grey stringers and masses, white at 210.38-210.46, grey 210.60-210.63 white 210.70-210.75, host fg laminated tuff.	35178	210.26	210.75	0.49	0			
210.75	215.33	medium to dark green, laminated tuff fragmental with grey and white silicic and carbonated ll to bed moderate to well level CA-55-58o moderately hard, less siliceous, 212.85-212.98 dark green tuff, 212.58-212.85 fg medium grey tuff lapilli massive poor moderate level bedding CA-60o, small elongated fragments locally 2-3% fg py.	35179	210.75	211.7	0.95	0			
			35180	211.7	212.58	0.88	0			
			35181	212.58	212.98	0.4	0			
			35182	212.98	214.4	1.42	0			
			35183	214.4	215.33	0.93	0			
215.33	216.06	foliated felsic dike, fg buff to light brown at contacts for approximate 10cm changing to fg, moderately foliated CA-60-65o, hard, non magnetic non carbonated, massive, uniform, contacts sharp with massive chlorite and minor amount of qtz carbonated stringers, qtz ff stringers random with bleaching altn, few grey opaque qtz stringers 60o x-cut foliation, trace to scattered fg py, LC sharp CA-60o x-cut qtz ankinte chlorite schistose tuff fragmental at low angular oblique.	35184	215.33	216.06	0.73	0.09	0.13		
216.06	218.92	fragmental tuff UM, same as 210.75-212.58 and 212.85-215.33, medium	35185	216.06	217.5	1.44	0			

Hunter Mine - Diamond Drill Log HM-04-38												
Property:	Hunter Mine			Hole Dip:	-80	Page No.:	1 of 11					
Location:				Hole Azimuth:	105	Date Started:	July 28, 04					
Claim No:	HR 1009			Hole Length:	218.63m	Date Finished:	Aug 3, 04					
Elevation:	Porcupine Lake			Purpose:		Drill Co.:	Benoit					
UTM Coords.:	5370768.3N, E487021.0					Logged by:	K.Jensen					
Meterage		Description			Sample			Assays				
From	To				No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
0.0	59.2	Casing										
59.2		Ultramafic Talcose Rocks										
59.2	66.2	UM same as before, 59.23-59.34 white opaque qtz vein LC 40o, 59.34-60.0 light black green, very talcose mass schistose near II to CA, 60.0-61.09 brecciated carbonated healed large fragments, 61.09-61.13 intensely sheared schistose CA 55-60o and near II to CA LC sharp CA-65o, 61.13-62.27 white qtz vein nil sulphide or carbonate LC sharp 62o, 62.27-62.37 brecciated UM, 62.37-62.73 qtz vein with large angular inclusions, talcose on contacts, nil sulphides, white irregular contacts, 62.73-63.88 massive, schistose near II to CA 12-15o LC 20o, 63.88-64.80 brecciated qtz carbonate fragments, nil to poor level of schistosity gradation contact, 64.80-65.13 massive poor level schistose small 2-3mm qtz eyes, 65.13-66.17 brecciated qtz carbonated healed.			35201	62.13	62.73	0.6	0			
66.17	79.11	qtz carbonated brecciated UM of possible fragmental UM, crushed zone sections from 66.32-66.44, 66.65-68.09, 69.85-70.30, 71.95-72.0, 74.08-74.33 75o, scattered mg to cg pyrite, 75.26-75.56 irregular masses of white opaque qtz irregular with inclusions with breccia.			35202	75	75.56	0.56	0			
79.11	81.79	massive, black green, porphyritic from 79.74-80.55 brecciated 80.55-80.84 on vein near II to CA, 80.84-81.66 few scattered irregular qc veinlets slips, 81.66-81.79 porphyritic qtz and plagioclase LC sharp CA-70o.										
81.79	97.27	brecciated, calcite and qtz healed, rare distinct stringers or veins except at 82.40 3cm qtz calcite CA-30o, 83.05-83.11 white to creamy calcite CA-35o, 84.40-86.56 schistosity II to CA, 94.15-95.35 scattered small sections of sheared schistose UM with elongated black hexagonal x-cut, 95.56 1cm band of pyrite fg with tuffaceous material or schistose CA-30o.										
97.27	143.5	Laminated Ultramafic Tuffs										
97.27	105.03	tuff to tuffaceous fragmental, fg, black to black green grading to medium green with veining siliceous bands at 104.43 locally crenulations, minor kinkle folding			35203	99.75	100.45	0.7	0.02			
		qtz carbonate usually II to bed well level CA-48o at 98.0, 60o at 100.4, scattered			35204	100.45	100.62	0.17	0			
		fg, mg py, 102.0-102.11 2-4mm pyrite stringers II to bed tuff, 99.0-99.23 intense			35205	100.62	101.56	0.94	0.02			
		veining kinked S shaped, 100.45-100.62 grey qtz vein minor pyrite fg CA-50-62o			35206	101.56	102.2	0.64	0			
					35207	102.2	103	0.8	0			
97.27	105.03	100.62-101.56 qtz carbonated st 1/2 II to bedding and fragments all mg			35208	103	104.02	1.02	0.03	0		

Property		Hunter Mine	Hole No.	38		Sheet No.	2			
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au Check	Au (2nd)	Au check
		bedding intense veining, 101.43-101.49 qtz veinlet irregular CA-65o, 101.56-104.02 more tuffaceous, devel. crenulations, scattered py, qtz st II to bedding	35209	104.02	104.4	0.38	0			
		usually less vfg-fg pyrite, several near II to CA and fractured LC-65o, 104.02-104.40 irregular qtz ankerite stringers, veinlets with minor inclusions, trace to scattered pyrite, 104.40-105.03 scattered grey opaque qtz veining fragments and irregular stringers.	35210	104.4	105.03	0.63	0			
105.03	109.56	tuffaceous with isolated small fragments locally crenulated, scattered qtz carbonated stringers random orientated 106.95 1 1/2 cm qc CA-20o x-cut bed, 107.11-107.30 1 1/2cm near II to CA irregular with fragment, LC 60-70o.								
109.56	110.4	brecciated qtz carbonated healed, contorted bedding x-cut by chlorite filled slip planes CA-15o.								
110.4	110.93	laminated black green and q S fold and kwinkled folded laminated tuff LC-50o								
110.93	112.12	fragmental with minor tuff, contorted bedding, qtz ankerite stringer at 111.06 V shaped and pyrite 4mm to 1.5cm CA-40o II to bed, 111.91 1 1/2cm irregular CA-50o x-cut bed low angle, 112.05-112.12 qtz ankerite and pyrite and chlorite CA-75o sinuous x-cut bed.								
112.12	112.63	brecciated small fragments chlorite healed, no stringer, LC-65o.								
112.63	113.41	tuffaceous fragmental scattered mg py, local contorted bedding, LC sinuous CA-50o.								
113.41	115.68	massive with 2-3mm greenish white phenocrysts, porphyritic texture, x-cut by irregular orientated qc stringers 5-8mm, scattered mg cg py x-cuts LC ground CA-45o.								
115.68	116.32	tuffaceous fragmental contorted and tight S folding with scattered cg up to 1/2cm py cubes contorted LC.								
116.32	117.24	contorted tuff, upper portion, non deformed bedding CA-80o to 116.67, scattered fg py LC irregular S shaped.	35211	116	117.24	1.24	0			
117.24	118.11	intense qtz veining, white opaque, 4-1 1/2 to 3cm CA-50-70o x-cut bedding, 117.74-118.11 large v with talcose tuff inclusions, LC ground 80o UC CA-45o.	35212	117.24	118.11	0.87	0			
118.11	121.4	tuff with fine fragments lapilli, grey green local laminated and bedding to 118.50 CA-50o, 118.50-121.29 fragments are all porphyritic with 1-2mm plagioclase phenocrysts showing no schistosity or bedding, possibly massive filled intensely veined in all same direction CA-65-55o, 119.13-119.93 with	35213	118.11	119	0.89	0			
		1cm qtz opaque st CA-42o at 119.93, intensely stringers 120.56-121.0 all same direction with felsic aphanitic dikelet from 120.64-120.70 CA-55o and scattered fg py, 121.29-121.40 laminated with grey silic LC at 121.40 broken.	35214	119	120	1	0			
			35215	120	121.4	1.4	0.05			
118.11	121.4	1cm qtz opaque st CA-42o at 119.93, intensely stringers 120.56-121.0 all same direction with felsic aphanitic dikelet from 120.64-120.70 CA-55o and scattered fg py, 121.29-121.40 laminated with grey silic LC at 121.40 broken.								
121.4	122.47	light grey to light medium grey, very hard, siliceous, medium grained grading to fine grained downhole, scattered irregular crispy qtz calcite stringers from	35216	121.4	122.47	1.07	0.33			

Property		Hunter Mine		Hole No.	38		Sheet No.	3			
Meterage		Description		Sample			Assay				
From	To			No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		121.96-122.47, vfg to fg pyrite scattered to disseminated 1% from 121.63-122.36, 121.40-121.63 greyish white development of plagioclase x-cuts with mg pyrite, LC CA-60o weak moderate level bed CA-50o.									
122.47	122.59	tuff fragmental UM light grey green to dark grey green, LC with 1/2cm qtz st CA-60o.		35217	122.47	122.79	0.32	0.04	0.07		
122.59	122.79	porphyritic felsic 2mm plagioclase x-cuts grading to fragmental, grading mg to fg py overall 2-3% LC sharp CA-45o.									
122.79	127.39	tuff-tuff fragmental, fg, black green, tuff to tuffaceous fragmental, moderate well development of bedding locally poor devel. low to massive tuff sections, local sections intensely veined with 2-4mm qtz stringer overall 50-60%, locally kinkle folding, with chlorite ff slip planes, more crenulations, nil to trace to locally 1% fg pyrite, 122.79-127.39 qtz flooded, small stringers, generally all in same direction ll to or near bedding 60o, 123.56-123.63 felsic dikelet CA-70-50o, 124.39-124.47 felsic dikelet CA-50o 2-3% mg py, 124.47-124.99 kinkled folds, 124.99-125.05 white opaque qtz vein CA 40-52o, 125.52-126.41 1-2mm grey white x-cuts, porphyritic texture, massive, nil schistose or bedding, stringer veins same direction as tuff fragmental above less density LC 55o, 126.41-127.39 qtz flooding LC-50-55o.		35218	122.79	124	1.21	0.02			
127.39	128.58	light to medium grey green to olive, decreasing orientation of stringers usually ll to bed 50o with few scattered stringers x-cut bedding mostly discont. due to kinkle folding, lapilli tuff, trace sulphides LC-50o.									
128.58	129.05	fragmental unit with buff to light greenish buff fragments elongated with medium to dark green tuff matrix LC sharp CA-60o.									
129.05	130.94	tuff with small sections of tuff fragmental light grey green, few stringers usually ll to bedding scattered stringers x-cut bedding and folded stringers x-cut bedding and folded stringers x-cut bed, 130.92-130.94 pinkish light greenish yellow sericitic altn felsic dikelets near ll to bed CA-55o.									
130.94	131.17	fragmental tuff.									
131.17	133.85	medium green grading to dark green tuff with small isolated sections of fragment crenulations with chlorite ll slip planes, scattered qtz stringers 2-5m.									
133.85	134.55	qtz flooded 1/2-4cm grey opaque qtz with ankerite generally CA-55o, few grey opal qtz stringer x-cut by above CA-40o, contacts UC 50o LC-45o.		35219	133.9	134.6	0.7	0			
134.55	138.72	tuff fragments as above, qtz ankerite, x-cut bed scattered, CA-75,50o, 135.28 3mm qtz ankinte stringer with pyrite CA-45o ll to bed, 136.0-137.05 intense qtz weak stringer x-cut bed 2-5mm CA-60o, 75o, 85o, 30o discont., 137.35 1cm qtz ankerite stringers CA-75o, 137.53-138.72 qtz flood, brecciated fragmental tuff with scattered pyrite.		35220	137.5	138.7	1.19	0.51	0.54		

Property: Hunter Mine		Hole No.	38		Sheet No.	4				
Meterage		Description	Sample			Assays				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
138.72	138.79	felsic pale pinkish light tan, fragments tuff, siliceous, void of stringers CA-70-60o	35221	138.7	138.9	0.15	0			
138.79	138.87	siliceous fragmental breccia LC 40o.								
138.87	139.62	mafic medium green fragmental tuff with irregular greyish qtz stringer fragments scattered to 0.5% vfg to fg pyrite LC sharp 53o.	35222	138.9	139.6	0.75	0			
139.62	139.87	light tan to pale buff siliceous minor amount of chlorite felsic fragmental grading contact with veining chlorite.	35223	139.6	139.9	0.25	0			
139.87	141.41	chlorite matrix with greyish fragmental locally brecciated, x-cut by greyish opague and grey qtz opague and ankerite stringers up to 1cm CA-40, 55, 70o and II to CA at 141.07-141.24 side of core, weak moderate devel. of bedding CA-40o LC-35o, scattered fg py in stringers, nil to trace pyrite in fragmental tuff.	35224	139.9	140.6	0.73	0			
			35225	140.6	141.4	0.81	0			
141.41	143.5	light grey to light greyish light green, to medium grey, tuff, stringers II to bed 25o same x-cut at 50o, 65o, scattered vfg pyrite, 142.68-142.79 intense stringers II to bedding with minor fuchsite CA-30o, 143.30-143.33 light brownish siliceous CA-55o dikelet felsic.	35226	141.4	142.3	0.89	0			
			35227	142.3	142.8	0.49	0.01			
			35228	142.8	143.5	0.71	0			
143.5	213.25	Exhalitic Tuffs and Sedimentary Rocks								
143.5	143.97	felsic breccia, fg, light grey to medium grey, brecciated, healed laminated by chlorite, with pyrite to 143.73, x-cut by white qtz stringer with pyrite CA-80o at 143.58 and 50o at 143.64, from 143.58-143.64 healed with chocolate brown silica LC sharp CA-35o	35229	143.5	144.0	0.47	0			
143.97	145.1	felsic foliated dike, fg, medium grey massive, uniform, weakly foliated with greyish silica qtz fragmental, silicified, siliceous, nil to very weak development of foliation except near contacts, scattered pyrite vfg to fg, rare distinct stringers 143.97-144.16 weak level of foliation CA-25o, 144.16-144.78 massive, scattered py, 144.78-144.91 chloritic matrix with buff to tan fragments possible inclusion, 144.91-145.10 weakly foliated scattered pyrite 1% fg.	35230	144.0	144.8	0.81	0.05	0.09		
			35231	144.8	145.1	0.32	0.01			
145.1	145.71	qtz vein brecciated, felsic to approx, 145.28, 145.28-145.71 altn tuff fragment inclusions, silicified with dark brown silica tourmaline II white opague qtz ankerite veining and stockwack, 145.52-145.71 1/2cm of grey qtz stringers CA-40o, white opague qtz ankerite with scattered py and minor pyrrhotite, 145.71 LC CA-45o sharp.	35232	145.1	145.71	0.61	0			
145.71	147.76	tuff to tuff fragmental, 145.71-146.35 fg tuff, patchy moderate sericitic altn, minor contorted bedding, CA-40o changing to 60o at LC-60o, 146.35-146.71 crispy chocolate brown matrix to 146.57 changing to medium green chloritic fragmental tuff, LC-45o, 146.71-146.87 medium green tuff with small 2-3mm grey white silica bands, bands, LC-40o, 146.87-147.29 light grey fragments in dark green tuff matrix poorly devel. bedding, trace sulphides LC-45o, 147.29-147.37 light tan to buff felsic dikelet CA-60o, 147.37-147.41 chloritic tuff,	35233	145.71	146.35	0.64	0.07			
			35234	146.35	146.87	0.52	0.93	0.9		
			35235	146.87	147.76	0.89	1.13	1.05		

Property: Hunter Mine		Hole No.	38		Sheet No.	5				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		147.41-147.71 qtz flooded greyish brown silica stringers 1-1 1/2cm CA-60o x-cutting weakly bedded Bx fragments, LC-55o, 147.71-147.76 medium green laminated tuff LC-50o x-cut bedding.								
147.76	148.33	brecciated zone, fg, buff to tan, sub angular fragments within altn matrix to yellow greenish tan to chocolate brown siliceous matrix, small grey white opaque qtz masses or fragments of veins, trace to scattered vfg fg py, void of distinct stringers, LC sharp CA-50o.	35236	147.76	148.33	0.57	0			
148.33	152.12	tuff fragmental, grey green, light grey to medium grey, 148.77-148.89 irregular grey white opaque stringers CA-70o II to bed, 149.08-149.35 tuff fragmental with white opal 1/2cm qtz st CA-70, 48, 65o opposite and irregular mass and vfg py CA-55, 60o LC II to bed, 149.35-150.05 tuff dark grey to grey green, 150.05-152.12 weak patchy sericitic altn, tuff fragmental pale green and medium green, 150.15-150.68 brecciated, qtz flood light to medium grey qtz stringers and folding vfg py, 150.68-152.12 patch weak to weak moderate sericitic tuff fragmental, with white opaque and grey white opaque qtz stringer 1/2 to 2.5cm CA-50o II to bedding, 151.39-151.45 1/2cm pinkish qtz carbonated stringer CA-30o II bed, 152.04-152.12 qtz veinlet with minor fuchsite, irregular CA-50o near II to bed.	35237	148.33	149.08	0.75	0.01			
			35238	149.08	149.35	0.27	0.19			
			35239	149.35	150.05	0.7	0.44			
			35240	150.05	150.68	0.63	0.05			
			35241	150.68	152.12	1.44	0			
152.12	152.68	patchy moderate sericitic altn fragmental tuff, trace to scattered vfg py, LC-50o	35242	152.12	152.68	0.56	0			
152.68	153.55	q flooded, moderate to strongly sericitic altn qtz II to bedding of fragmental	35243	152.68	153.55	0.87	0.03			
152.68	153.55	tuff, LC 45o II to bed.								
153.55	154.09	buff to dark brown locally minor sericitic altn fg tuff with small fragments lapilli tuff LC 45o, pink carbonated qtz st at 153.59 1cm, 153.64 1cm, 153.75 1/2cm CA-45o.	35244	153.55	154.09	0.54	0			
154.09	154.68	qtz flooded locally brecciated moderate to strongly sericitic altn, minor fuchsite altn stringers in BX 154.09-154.28 pale greenish tint 154.28 qtz flooding II to bed, trace to scattered fg py LC 35-37o bedding.	35245	154.09	154.68	0.59	0			
154.68	155.01	tuff, pervasive moderate sericitic altn good laminations CA-42o.	35246	154.64	155.01	0.33	0			
155.01	156.15	grey green to black green, scattered milky white qtz carbonated to wispy pink carbonated qtz stringer II to bedding CA-40o nil to trace pyrite LC-50o.	35247	155.01	156.15	1.14	0			
156.15	156.68	weak to locally moderate patch sericitic altn, weak rare pervasive, tuff laminated CA-55o, 156.39-156.51, grey white qtz ankerite veinlet CA-50o x-cut by 2 - 1/2cm grey to brownish grey and black chlorite stringer CA-70+65o, 156.68 LC CA-50o.	35248	156.15	156.68	0.53	0.02			
156.68	157.04	qtz tuff flooded laminated tuff, brecciated by qtz, grey white opaque and medium grey opaque qtz, moderate sericitic altn of tuff, trace to scattered	35249	156.68	157.04	0.36	0			

Property: Hunter Mine		Hole No.	38		Sheet No.	6				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		pyrite LC 60o.								
157.04	157.35	brecciated tuff healed with weakly sericitic altn chlorite, qtz grey to medium grey, light grey to buff tuff, scattered to disseminated vfg pyrite 1-2%, LC-55o.	35250	157.04	157.37	0.33	0.08			
157.35	158.02	weak pervasive with locally moderate sericitic altn silicified by minor qtz flooding, scattered 1% overall locally 157.35-157.47 vfg 2-3% py LC-65o II bed	35251	157.37	158.02	0.65	0.15	0.16		
158.02	158.59	fg light buff to light brownish, hard, weakly carbonated well laminated tuff to argillite CA-55o 158.05 1/2cm qtz st x-cut bed CA-60o, 158.02-158.20 more brecciated vfg pyrite 3-5%, grading to 1-2% vfg, LC sharp CA-60o.	35252	158.02	158.59	0.57	0.38	0.39		
158.59	160.02	fg to mg massive uniform, weak devel. bed, grading to fg buff to medium brown carbonated weakly greywacke to argillite 159.12, well bedded 159.42-160.02 alternating black green, buff, tan laminations, odd 2-3mm qtz apperance of fragmental tuff, vfg py <0.5%.	35253	158.59	159.12	0.53	0			
			35254	159.12	160.02	0.9	0.01			
160.02	160.9	qtz flooded, weak patchy sericitic altn, fragmental tuff grading to fine laminated tuff, qtz masses, II to bed and x-cut bed, scattered vfg pyrite LC-50o.	35255	160.02	160.9	0.88	0.01			
160.9	161.94	well laminated tuff, weakly pervasive sericitic altn, rare qtz stringer usually x-cut bed, scattered to desseminted vfg to fg pyrite 1-2%, LC-50o.	35256	160.9	161.94	1.04	0.02			
161.94	162.49	blackish, light to medium grey, buff laminations, minor greyish qtz stringers,	35257	161.94	162.49	0.55	0.02			
161.94	162.49	3 pink qtz carbonated stringer II to bedding 2-5 mm CA-50o sharp, 162.11 grey to blackish qtz stringer CA-65o x-cut bedding with dragging of bedding on both sides, 162.39-162.49 greyish qtz stringers 1-3mm II to bed 162.49 contact CA 55o sharp.								
162.49	163.53	fg to mg, medium grey, massive uniform, weak to weak moderate level of bedding disseminated white 1/2mm grains, possible greywacke, non carbonated, bed 50o, locally mg pyrite up to 2mm x-cuts, no baked contacts hard, siliceous, scattered hairlike to 2mm qtz stringers II to bed, possible crystal tuff, LC sharp CA-45o.	35258	162.49	163.53	1.04	0			
163.53	163.9	fg, laminated, medium brown to light brown and buff laminations grading to greyish white and buff, bed excellent CA-55o, trace sulphides LC-43o sharp.	35259	163.53	163.9	0.37	0			
163.9	164.3	qtz stringers II to bed, irregular near II to CA and grey white to creamy qtz ank 164.27-164.30 in tuff fragmental LC-40o near II to bed CA-42-45o sinuous	35260	163.9	164.3	0.4	0.08	0.08		
164.3	165.32	nil to weak patchy sericitia altn with scattered 1cm to 2cm straight grey qtz stringers II to bed and irregular grey white qtz II and x-cut bedding, fragmental tuff LC-55o, scattered pyrite with tuff and grey qtz stringer.	35261	164.3	165.32	1.02	0			
165.32	165.86	fg, weak pervasive sericitic altn, rare stringers minor kinkle folding, grey black crenulated 1/2cm qtz stringer CA 70o x-cut bed at 165.37, 165.65 1/2 cm grey black qtz stringer II to bedding with pyrite, scattered fg pyrite LC-55o	35262	165.32	165.86	0.54	0.02			
165.86	166.68	patchy moderate sericitic altn with black to black green siliceous, scattered	35263	165.86	166.68	0.82	0			

Property: Hunter Mine		Hole No:	38		Sheet No.	7				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		pyrite with sericitic altn sections, more pyrite with unaltered sections, few 1/2 pink carbonated qtz stringers II to bed, buff to light brown to dark brown, tuff bands with black green fragmental tuff LC-60o.								
166.68	166.96	pervasive weak to weak moderate sericitic altn tuff, trace sulphides LC-40o trace pyrite.	35264	166.68	166.96	0.28	0			
166.96	167.12	qtz vein medium grey to blackish grey qtz vein with sericitic tuff inclusions UC straight sharp LC sinuous irregular CA-50-60o x-cut bedding.	35265	166.96	167.12	0.16	0.1			
167.12	167.9	weak patchy sericitic altn, tuff, scattered pyrite bands 1-2mm, scattered 1% to locally 1-2% disseminated pyrite, few 1-2cm grey opaque qtz ankerite stringers near II to bedding, LC-50o.	35266	167.12	167.9	0.78	0.05			
167.9	169.48	weak to moderate pervasive sericitic altn well laminated tuff, local small bands of crenulated tuff, bed 55-56o, scattered py locally 1-2% vfg-fg LC-55o.	35267	167.9	168.7	0.8	0.05			
169.48	170.01	black green and buff laminated tuff, few fragments, few blackish to black grey	35268	168.7	169.48	0.78	0.14			
169.48	170.01	siliceous II to bed, occasionally pyrite 1-2mm band, void of distinct st LC-55o	35269	169.48	170.01	0.53	0.09			
170.01	171.6	moderate pervasive sericitic altn, laminated tuff with few fragments, void of stringers, scattered pyrite <0.5% well level bed CA-57-60o LC-45o.	35270	170.01	170.9	0.89	0.21			
171.6	171.9	laminated tuff fragmental, grey to buff and black green, LC 1/2cm grey white qtz ankerite st CA-65o x-cut bed.	35271	170.9	171.6	0.7	0.25	0.18		
171.9	172.08	laminated tuff fragmental, grey to buff and black green, LC 1/2cm grey white qtz ankerite st CA-65o x-cut bed.	35272	171.6	172.2	0.6	0.13			
171.9	172.08	brecciated tuff fragmental healed with grey qtz and chlorite, minor to scattered py vfg fg, LC sharp 50o.								
172.08	172.2	altn to buff and medium from fragmental tuff, silicified 2 grey opal qtz st in opposite direction forming x-cut CA-35o+28o near II to bed, scattered vfg fg py, splash of chalcopyrite, LC with qtz vein 43o.								
172.2	172.9	light tan tuff fragmental of small elongated qtz altn bleached by qtz veinlet at UC 3cm, 1cm grey qtz at 172.27 90o, and qtz vein with inclusions 172.35-172.90, fragmental to 172.39, tuff 172.39-172.90 with few qtz eyes LC-50o overall irregular.	35273	172.2	172.9	0.7	0			
172.9	175.87	fg, light grey to medium grey green to light olive, small numerous fragments, occasional qtz stringer usually II to moderate devel. of bedding CA-50-55o, local kinkle folding with ff chlorite slip planes, scattered to trace vfg py qtz ar 173.0-173.03 irregular x-cut bed, 173.70-173.82 with inclusions near II to bed CA 60o irregular, 174.05 1cm grey II to bed, 174.13-174.16 II to bed grey 174.22 grey white translucent sinuous near II to bed, 174.82 grey qtz ankerite 1cm CA-55o II to bed, 175.04-175.55 several grey st, qtz st 1/2-1cm II to bed 175.70-175.76 irregular qtz veinlet 50o II to bed then x-cut bed, 175.76-175.87 fg tuff, LC 48-50o.	35274	172.9	173.68	0.78	0.03			
			35275	173.68	173.87	0.19	0			
			35276	173.87	174.25	0.38	0			
			35277	174.25	175.04	0.79	0			
			35278	175.04	175.87	0.83	0			
175.87	176.66	light to medium grey fragmental tuff, trace pyrite, massive uniform, laminated	35279	175.87	176.66	0.79	0			

Property		Hunter Mine	Hole No.	38		Sheet No.	8			
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		bedding 2-3mm 175.92 4cm whitish qtz st opaque ll to bed with grey opaque qtz stringer branching x-cut bedding, 176.66 increasing silica grey bands and darker colour gradational contact over 10cm.								
176.66	178.48	layers of grey silica and black green and dark grey layers, fragmental tuff, uniform, good bed 50-55o slightly talcose, chloritic, hard, trace pyrite, void of distinct stringers LC-45o.	35280	176.66	177.52	0.86	0.02			
			35281	177.52	178.48	0.96	0	0		
178.48	179.59	same as 176.66-178.48 lght qtz stringer and qtz flooded, grey green to black green, uniform, LC 60o.	35282	178.48	179.59	1.11	0			
179.59	180	same as 176.66-178.48 LC approximate 60o.	35283	179.59	180	0.41	0			
180	182.6	mafic dike, fg, medium to dark grey, altered to medium brown by qtz white cloudy stringers from 180.0-180.48 and 181.77-182.60, massive uniform, hard to very hard siliceous, weak foliation of black green chlorite from 181.14-182.35, scattered white grey qtz stringers with pyrite to 180.60, large low angular qtz stringer x-cut by grey opaque qtz vein with minor chalcopyrite, from 180.60-180.84, scattered po in stringers and dike matrix to 181.44, 181.44-181.48 white opaque qtz st CA 40o irregular with pyrite patchy fracture CA-45o, 181.77-182.13 pyrrhotite associated with ff qtz stringers, 182.13 1cm white translucent qtz st CA-60o, 182.14-182.60 chlorite ll and qtz hydrofracturing, light brownish tan to medium brown altn, scattered pyrrhotite associated with qtz ff, 182.60 LC gradational with altn tuff fragment CA-40o.	35284	180	180.6	0.6	0			
			35285	180.6	181.49	0.89	0.05			
			35286	181.49	182.13	0.64	0.03			
			35287	182.13	182.6	0.47	0.02			
182.6	184.74	fragmental tuff, fg grey green very weak pervasive sericitic altn, fragmental tuff high silica contact with grey qtz, rare grey opaque qtz stringer, 184.05-184.07 grey opaque qtz stringer CA-65o x-cut bed, bed average 50o, trace to nil sulphides LC-40-45o, 184.54-184.74 x-cut 1cm grey opaque to opal qtz st some straight and some contorted S folded.	35288	182.6	183.65	1.05	0.01			
			35289	183.65	184.74	1.09	0			
184.74	185.23	very qtz rich, cloudy grey with black green chlorite and few fragments of olive green tuff fragmental, LC with grey opal qtz stringers CA-50o.	35290	184.74	185.41	0.67	0			
185.23	185.41	fg, olive green, weak sericitic altn tuff fragmental with grey opal bands or stringers ll to bed.								
185.41	186.43	grey green laminated tuff fragmental, trace pyrite, qtz st at 185.41-185.44 and 185.67-185.69 ll to bed LC ground.	35291	185.41	186.41	1	0			
186.43	186.57	mafic dike to intermediate, fg, medium grey to light grey bleaching around qtz ankerite stringers, massive, uniform, moderate development of foliation CA-50 with black chlorite alignment to form foliation, siliceous scattered qtz ankerite stringers 2-3mm up to 2-3cm, trace to scattered vfg pyrite locally mg py up to 1-2%, 186.41-186.57 aphanitic to fg contact siliceous qtz st fragments,	35292	186.41	186.71	0.3	0			

Property: Hunter Mine		Hole No.	38		Sheet No.	9				
Meterage		Description	Sample			Assay				
From	To		No.	From	To	Width	Au (g/t)	Au check	Au (2nd)	Au check
		displaced qtz stringer by chlorite ff CA-70o, medium to dark grey, 186.57- contact sharp 80o sinuous.								
186.57	186.71	light brownish grey, poor to weakly foliated deformed 1cm qtz stringers which are laminated at both contacts fractures CA-60-65o, stringers CA-55o qtz ankerite with scattered pyrite, possible inclusion.								
186.71	186.99	aphanitic to fine grained medium to dark grey massive uniform, hard, siliceous, weak devel. foliation few chlorite ff stringers hairlike, pale grey white ghost phenocrysts, 186.73 4mm qtz grey st CA-70o, 186.98 3mm qtz grey st CA-80o with pyrite blobs, scattered to disseminated vfg to fg pyrite 1-2%.	35293	186.71	186.99	0.28	0.04			
186.99	187.58	medium grey weak moderate devel. foliation several qtz ankinte stringers x-cut foliation CA-50o, 187.00 2mm qtz stringer with 2mm alteration on both sides 40o, 187.03 2mm qtz st with 2mm alteration on both sides 40o, both x-cut by chlorite qtz ff CA-27o, 187.16 1-1.2cm qtz ankerite veinlet with chlorite, pyrrhotite CA-40o crenulated 20o x-cut foliated at low angle, 187.22-187.24 grey qtz opaque veinlet with 3-5% pyrrhotite and splashes of chalcopyrite ll to foliation, CA-45o, 187.25-187.29 qtz carbonated veinlet with pyrite and pyrrhotite LC-50o, 187.39-187.43 grey qtz vein with grey inclusions and pyrrhotite 1% CA-70+50o, 187.43-187.58 scattered qtz ankerite stringers and grey opaque qtz stringers near ll to foliation.	35294	186.99	187.58	0.59	0.06			
187.58	188.61	moderate to well foliated, medium to dark grey, fg, with alignment of chlorite scattered qtz stringers ll to and near ll to bed x-cut foliation, 187.74 1cm grey white qtz stringer CA-40o, 187.89 3/4cm grey opaque qtz stringer CA-65o, 187.91-187.97 x-cut foliation qtz white creamy stringers with pyrrhotite CA-45-85o 2-4mm, 188.06-188.14 1.2cm grey creamy qtz stringer ll to foliation CA-45o with po associated with chlorite on contact with branch ivory stringer 188.23 chlorite and py ff st CA-40o 2-4mm, 188.61 contact sharp CA-50o.	35295	187.58	188.5	0.92	1.85	1.87		
188.61	188.66	brownish grey felsic dikelet 2-3% vfg pyrite, few chlorite ff hairlike, massive uniform, qtz and plagioclase feldspar, minor mafic LC sharp at 55o.								
188.66	189.3	medium grey well foliated, 188.89-189.0 4 chlorite ll stringers with bleaching of dike, 2-3mm gash CA-55-35o x-cut foliation at low angle, overall 1% vfg to fg py, 189.0-189.30 x-cut by numerous 2-3 up to 1cm qtz grey translucent stringers near ll to foliation and 1 x-cut foliation at high angle CA-70o with pyrrhotite, 189.17-189.30 5 1-2mm white qtz st with pyrite CA-70-75o and 65o in opposite direction, 189.30 baked contact CA-60o.	35296	188.5	189.3	0.8	0.1			
189.3	191.42	tuff fragmental, fg, matrix light to medium brown, with small greyish and creamy fragments well laminated with grey opaque, grey translucent and greyish white qtz carbonated layers, non carbonated calcite, rare distinct st	35297	189.3	190.4	1.1	0			
			35298	190.4	191.42	1.02	0			

Hunter Mine Project HM-04-39

Property:	Hunter Mine	Hole No.:	HM-02-39	Page No: 1	1 of 1
Location:		Hole Angle:	-75	Date Started:	30-Jul-04
Claim Number:	HR 1009	Hole Azimuth:	105	Date Finished:	03-Aug-04
Elevation:	Porcupine Lake	Hole Length:	309	Drilling Company:	Benoit
UTM Coordinates	5370982.0N, E486941.0	Avg. Core Recovery:	+99%	Logged By:	D. McBride

Meterage		Description	Sample				Assays	
From	To		No.	From	To	Width	Au ppb	Cu(ppm)
0.00	49.00	Casing						
49.00	275.80	Talcose Ultramafic Fragmental Tuffs and Agglomerates	35313	142.60	143.28	0.68	no assay	
		Soft, dark green matrix with harder fragments to 10 cm and lapilli bands	35314	143.28	143.90	0.62	no assay	
		So 60o CA, S1 50o CA	35315	143.90	144.63	0.73	no assay	
		very heterogeneous sequence with fine-grained more massive sections from 149.7 to 159.3 and 169.2 to 190	35316	144.63	145.40	0.77	no assay	
			35317	145.40	146.20	0.80	no assay	
275.80	301.20	Laminated Ultramafic Tuffs						
		Fine-grained laminated rock, diagnostic by dark grey to black laminations, S0 75o CA						
		after 289m, lighter grey and less laminated than most S0 60o CA						
		lacks brownish colour of exhalitic sedimentary rocks and seems slightly coarser grained, S0 50o CA- lower contact						
301.20	305.00	Talcose Ultramafic Fragmental Tuffs						
		similar to above, but with carbonate bands						
305.00	309.00	Clastic Sedimentary rocks						
		Fine-grained, medium grey to dark grey sericitic sedimentary rocks						
		S0 10-20o CA, S2 05o CA & 30oCA to S1, S3 70o & 80o CA S1						
309.00		EOH						

APPENDIX II
Assay Certificates



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1444-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-16-04

We hereby certify the following Assay of 45 Core samples submitted JUL-08-04 by .

Sample Number	Au g/tonne	Au Check g/tonne	Au 2nd g/tonne	Au 2ndCk g/tonne
44501	0.02	-	-	-
44502	0.03	0.04	-	-
44503	Nil	-	-	-
44504	Nil	-	-	-
44505	Nil	-	-	-
44506	Nil	-	-	-
44507	Nil	Nil	-	-
44508	0.01	-	-	-
44509	0.02	-	-	-
44510	0.03	-	-	-
44511	0.16	-	-	-
44512	32.88	37.85	37.03	34.49
44513	0.22	0.21	-	-
44514	0.16	-	-	-
44515	0.01	-	-	-
44516	0.10	-	-	-
44517	0.24	-	-	-
44518	0.03	-	-	-
44519	0.03	-	-	-
44520	0.05	-	-	-
44521	0.10	-	-	-
44522	0.84	0.81	-	-
44523	0.23	-	-	-
44524	0.43	-	-	-
44525	0.02	-	-	-
44526	0.07	-	-	-
44527	0.02	-	-	-
44528	0.05	0.06	-	-
44529	Nil	-	-	-
44530	0.03	-	-	-

Certified by Judy Russo



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate


4W-1444-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **JUL-16-04**

We hereby certify the following Assay of 45 Core samples submitted JUL-08-04 by .

Sample Number	Au g/tonne	Au Check g/tonne	Au 2nd g/tonne	Au 2ndCk g/tonne
44531	Nil	-	-	-
44532	0.04	-	-	-
44533	Nil	-	-	-
44534	0.10	-	-	-
44535	0.04	-	-	-
44536	0.26	-	-	-
44537	0.68	-	-	-
44538	0.21	-	-	-
44539	0.01	-	-	-
44540	0.01	-	-	-
44541	0.02	-	-	-
44542	0.42	0.41	-	-
44543	Nil	-	-	-
44544	Nil	-	-	-
44545	Nil	-	-	-
Blank	Nil	-	-	-
STD OxK18	3.39	-	-	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1445-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **JUL-16-04**

We hereby certify the following Assay of 45 Core samples submitted JUL-08-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44546	Nil	-
44547	Nil	-
44548	0.02	0.01
44549	0.05	-
44550	Nil	-
44551	Nil	-
44552	0.03	-
44553	0.11	-
44554	Nil	-
44555	0.10	0.19
44556	0.03	-
44557	0.02	-
44558	Nil	-
44559	0.01	-
44560	0.01	-
44561	Nil	-
44562	Nil	-
44563	Nil	-
44564	0.03	-
44565	Nil	-
44566	0.04	0.03
44567	0.01	-
44568	Nil	-
44569	Nil	-
44570	Nil	-
44571	Nil	-
44572	Nil	-
44573	0.17	-
44574	0.03	-
44575	Nil	-

Certified by  _____



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate

4W-1445-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **JUL-16-04**

We hereby certify the following Assay of 45 Core samples submitted JUL-08-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44576	Nil	-
44577	Nil	-
44578	0.01	0.01
44579	Nil	-
44580	Nil	-
44581	0.01	-
44582	0.03	-
44583	Nil	-
44584	Nil	-
44585	Nil	-
44586	Nil	-
44587	0.01	-
44588	0.07	-
44589	Nil	-
44590	0.29	0.22
Blank	Nil	-
STD OXK8	3.50	-

Certified by *Julie Revo*



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1479-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **JUL-22-04**

We hereby certify the following Assay of 57 Core samples submitted JUL-11-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44591	0.02	-
44592	0.05	-
44593	0.05	0.04
44594	0.01	-
44595	0.03	-
44596	0.03	-
44597	0.01	-
44598	Nil	-
44599	0.06	-
44600	0.01	-
44601	0.04	0.04
44602	0.02	-
44603	Nil	-
44604	Nil	-
44605	Nil	-
44606	0.01	-
44607	1.51	1.49
44608	Nil	-
44609	Nil	-
44610	0.01	-
44611	0.10	-
44612	0.24	-
44613	0.33	-
44614	0.07	-
44615	0.22	0.23
44616	0.05	-
44617	0.58	0.67
44618	0.05	-
44619	0.20	-
44620	Nil	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate

4W-1479-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-22-04

We hereby certify the following Assay of 57 Core samples submitted JUL-11-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44621	0.10	-
44622	0.04	-
44623	0.06	-
44624	Nil	-
44625	0.04	-
44626	0.05	0.04
44627	0.05	-
44628	Nil	-
44629	0.08	-
44630	0.01	-
44631	0.04	-
44632	0.07	-
44633	Nil	-
44634	0.01	-
44635	0.02	-
44636	Nil	-
44637	Nil	-
44638	0.07	-
44639	Nil	-
44640	Nil	-
44641	0.10	0.06
44642	0.02	-
44643	0.03	-
44644	Nil	-
44645	Nil	-
44646	0.08	-
44647	0.09	-
Blank	Nil	-
STD OxK18	3.64	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1480-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-22-04

We hereby certify the following Assay of 33 Core samples submitted JUL-15-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44648	0.02	-
44649	0.05	0.08
44650	0.02	-
44651	0.03	-
44652	0.03	-
44653	0.41	-
44654	2.61	1.89
44655	0.29	-
44656	2.06	-
44657	36.48	36.82
44658	0.05	-
44659	Nil	-
44660	0.04	-
44661	0.01	0.02
44662	0.02	-
44663	0.01	-
44664	0.05	-
44665	0.02	-
44666	0.01	-
44667	Nil	-
44668	0.10	-
44669	1.17	0.90
44670	0.54	-
44671	0.03	-
44672	0.01	-
44673	0.01	-
44674	0.02	-
44675	0.02	-
44676	0.03	-
44677	0.01	-

Certified by Judy Perra



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate


4W-1480-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-22-04

We hereby certify the following Assay of 33 Core samples submitted JUL-15-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44678	0.01	-
44679	0.08	0.09
44680	0.01	-
Blank	Nil	-
STD OxK18	3.45	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate


4W-1496-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-20-04

We hereby certify the following Assay of 50 Core samples submitted JUL-15-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44681	Nil	-
44682	0.15	0.14
44683	0.11	-
44684	0.17	-
44685	0.01	-
44686	0.06	-
44687	0.02	-
44688	0.02	-
44689	0.04	-
44690	0.06	-
44691	0.01	-
44692	0.01	-
44693	0.01	-
44694	Nil	-
44695	Nil	-
44696	0.01	-
44697	0.01	-
44698	0.01	-
44699	0.01	0.01
44700	Nil	-
44701	Nil	-
44702	0.02	-
44703	0.02	-
44704	0.02	-
44705	0.01	-
44706	0.03	-
44707	0.01	-
44708	Nil	-
44709	Nil	-
44710	Nil	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate


4W-1496-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: JUL-20-04

We hereby certify the following Assay of 50 Core samples submitted JUL-15-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44711	Nil	-
44712	0.05	-
44713	Nil	-
44714	Nil	0.02
44715	Nil	-
44716	0.07	-
44717	0.07	-
44718	0.06	-
44719	0.02	-
44720	0.01	-
44721	Nil	-
44722	0.02	-
44723	Nil	-
44724	0.03	0.04
44725	0.02	-
44726	Nil	-
44727	0.02	-
44728	0.01	-
44729	0.02	-
44730	Nil	-
Blank	Nil	-
STD OxK18	3.55	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1497-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-22-04

We hereby certify the following Assay of 50 Core samples submitted JUL-15-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44731	0.03	0.03
44732	0.02	-
44733	Nil	-
44734	0.01	-
44735	Nil	-
44736	0.04	0.06
44737	0.13	0.10
44738	0.09	-
44739	Nil	-
44740	Nil	-
44741	Nil	-
44742	Nil	-
44743	Nil	-
44744	0.01	-
44745	Nil	-
44746	Nil	-
44747	Nil	-
44748	Nil	Nil
44749	Nil	-
44750	Nil	-
44751	Nil	-
44752	0.02	-
44753	Nil	-
44754	Nil	-
44755	Nil	-
44756	Nil	-
44757	Nil	-
44758	0.01	0.01
44759	Nil	-
44760	Nil	-

Certified by *Judy Pervo*



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate

4W-1497-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-22-04

We hereby certify the following Assay of 50 Core samples submitted JUL-15-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44761	0.01	-
44762	Nil	-
44763	Nil	-
44764	Nil	-
44765	Nil	-
44766	0.08	0.10
44767	Nil	-
44768	0.01	-
44769	Nil	-
44770	Nil	-
44771	Nil	-
44772	Nil	-
44773	Nil	-
44774	Nil	-
44775	Nil	-
44776	Nil	-
44777	0.01	-
44778	0.05	-
44779	Nil	-
44780	0.02	-
Blank	Nil	-
STD OxK18	3.51	-

Certified by Judy Revu



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 3

Assay Certificate


4W-1566-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: J. Jensen

Date: JUL-30-04

We hereby certify the following Assay of 75 Core samples submitted JUL-21-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43201	0.01	-
43202	0.02	-
43203	0.11	-
43204	0.01	-
43205	0.05	-
43206	0.11	-
43207	0.07	-
43208	0.04	-
43209	0.02	-
43210	0.07	-
43211	0.01	-
43212	0.11	0.08
43213	0.01	-
43214	0.04	-
43215	0.03	-
43216	0.02	-
43217	Nil	-
43218	0.18	-
43219	0.03	-
43220	Nil	-
43221	0.01	-
43222	0.01	-
43223	Nil	-
43224	Nil	-
43225	0.01	-
43226	Nil	-
43227	0.04	0.06
43228	0.01	-
43229	0.02	-
43230	0.01	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate

4W-1566-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **J. Jensen**

Date: **JUL-30-04**

We hereby certify the following Assay of 75 Core samples submitted JUL-21-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43231	0.05	-
43232	0.03	-
43233	0.01	-
43234	Nil	0.01
43235	Nil	-
43236	Nil	-
43237	Nil	-
43238	Nil	-
43239	Nil	-
43240	0.02	-
43241	Nil	-
43242	Nil	-
43243	Nil	-
43244	Nil	Nil
43245	Nil	-
43246	Nil	-
43247	0.01	-
43248	Nil	-
44781	Nil	-
44782	0.01	-
44783	0.03	-
44784	Nil	-
44785	Nil	-
44786	Nil	-
44787	Nil	-
44788	Nil	-
44789	Nil	-
44790	2.32	2.59
44791	0.20	-
44792	0.51	0.46

Certified by  _____



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate


4W-1566-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **J. Jensen**

Date: JUL-30-04

We hereby certify the following Assay of 75 Core samples submitted JUL-21-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44793	Nil	-
44794	Nil	-
44795	0.13	-
44796	Nil	-
44797	Nil	-
44798	Nil	-
44799	Nil	-
44800	0.12	0.12
44801	0.19	-
44802	Nil	-
44803	Nil	-
44804	Nil	-
44805	Nil	-
44806	Nil	-
44807	Nil	-
Blank	Nil	-
STD OxK18	3.28	-

Certified by  _____



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate


4W-1567-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-30-04

We hereby certify the following Assay of 37 Core samples submitted JUL-21-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44808	Nil	-
44809	1.06	0.98
44810	0.52	0.54
44811	0.02	-
44812	Nil	-
44813	0.02	-
44814	0.01	-
44815	0.01	-
44816	0.01	-
44817	Nil	-
44818	0.17	0.18
44819	0.01	-
44820	Nil	-
44821	Nil	-
44822	0.02	-
44823	Nil	-
44824	0.01	-
44825	0.05	-
44826	0.29	0.26
44827	0.01	-
44828	Nil	-
44829	Nil	-
44830	Nil	-
44831	Nil	-
44832	Nil	-
44833	0.04	-
44834	Nil	-
44835	Nil	Nil
44836	Nil	-
44837	0.06	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate


4W-1567-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: JUL-30-04

We hereby certify the following Assay of 37 Core samples submitted JUL-21-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44838	Nil	-
44839	Nil	-
44840	Nil	-
44841	Nil	-
44842	0.05	0.04
44843	0.02	-
44844	Nil	-
Blank	Nil	-
STD OxK18	3.40	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1607-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mines**
Attn: **K. Jensen**

Date: **AUG-03-04**

We hereby certify the following Assay of 56 Core samples submitted JUL-24-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44845	Nil	-
44846	0.01	-
44847	0.01	-
44848	Nil	-
44849	Nil	-
44850	0.01	-
44851	0.01	-
44852	0.02	-
44853	0.02	0.02
44854	0.01	-
44855	Nil	-
44856	Nil	-
44857	Nil	-
44858	Nil	-
44859	Nil	-
44860	Nil	-
44861	Nil	-
44862	Nil	-
44863	0.02	0.04
44864	0.01	-
44865	0.01	-
44866	Nil	-
44867	Nil	-
44868	0.02	-
44869	Nil	-
44870	0.01	-
44871	Nil	-
44872	Nil	-
44873	Nil	-
44874	Nil	-

Certified by Judy Puro



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate


4W-1607-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mines
Attn: K. Jensen

Date: AUG-03-04

We hereby certify the following Assay of 56 Core samples submitted JUL-24-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44875	Nil	-
44876	Nil	-
44877	Nil	-
44878	Nil	-
44879	Nil	Nil
44880	0.01	-
44881	0.03	-
44882	Nil	-
44883	0.01	-
44884	Nil	-
44885	Nil	-
44886	0.03	-
44887	0.02	-
44888	Nil	-
44889	Nil	-
44890	0.21	0.20
44891	Nil	-
44892	Nil	-
44893	0.04	-
44894	0.03	-
44895	Nil	-
44896	0.09	0.15
44897	0.02	-
44898	Nil	-
44899	Nil	-
44900	0.01	-
Blank	Nil	-
STD OxK18	3.37	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1608-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: AUG-03-04

We hereby certify the following Assay of 32 Core samples submitted JUL-24-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44901	0.02	-
44902	Nil	-
44903	Nil	-
44904	0.01	-
44905	0.03	0.04
44906	0.02	-
44907	0.01	-
44908	Nil	-
44909	Nil	-
44910	Nil	-
44911	0.02	-
44912	Nil	-
44913	Nil	-
44914	Nil	-
44915	0.03	-
44916	Nil	Nil
44917	0.01	-
44918	0.01	-
44919	0.01	-
44920	Nil	-
44921	Nil	-
44922	0.04	-
44923	0.66	0.47
44924	0.15	-
44925	Nil	-
44926	Nil	-
44927	Nil	-
44928	0.01	-
44929	0.06	-
44930	0.02	-

Certified by Judy Resco



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate

4W-1608-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **AUG-03-04**

We hereby certify the following Assay of 32 Core samples submitted JUL-24-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44931	0.01	-
44932	0.03	-
Blank	Nil	-
STD OxK18	3.37	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 3

Assay Certificate

4W-1700-RA1

Company: **VALGOLD RESOURCES INC.**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **AUG-23-04**

We hereby certify the following Assay of 68 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44933	0.01	-
44934	Nil	-
44935	Nil	-
44936	Nil	-
44937	0.01	-
44938	0.01	-
44939	0.01	-
44940	0.01	-
44941	Nil	-
44942	0.11	0.11
44943	0.02	-
44944	0.01	-
44945	0.03	-
44946	0.10	0.13
44947	0.06	-
44948	Nil	-
44949	Nil	-
44950	Nil	-
44951	Nil	-
44952	Nil	-
44953	0.01	-
44954	0.07	0.07
44955	0.02	-
44956	0.04	-
44957	0.03	-
44958	0.04	-
44959	0.02	-
44960	0.01	-
44961	Nil	-
44962	0.01	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate

4W-1700-RA1

Company: **VALGOLD RESOURCES INC.**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: AUG-23-04

We hereby certify the following Assay of 68 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44963	Nil	-
44964	0.01	-
44965	0.03	-
44966	Nil	-
44967	Nil	-
44968	0.01	-
44969	0.01	-
44970	Nil	-
44971	Nil	-
44972	0.06	-
44973	0.09	0.10
44974	0.04	-
44975	0.01	-
44976	Nil	-
44977	0.06	-
44978	0.31	-
44979	0.45	-
44980	0.01	-
44981	1.03	0.79
44982	1.86	1.95
44983	0.47	-
44984	0.11	-
44985	0.01	-
44986	Nil	-
44987	0.01	-
44988	0.02	-
44989	Nil	-
44990	0.01	-
44991	0.01	-
44992	Nil	-

Certified by Judy Perro



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate

4W-1700-RA1

Company: **VALGOLD RESOURCES INC.**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **AUG-23-04**

We hereby certify the following Assay of 68 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
44993	0.03	-
44994	Nil	-
44995	Nil	-
44996	Nil	-
44997	0.09	0.07
44998	Nil	-
44999	Nil	-
45000	0.01	-
Blank	Nil	-
STD OXK18	3.42	-

Certified by  _____



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 2

Assay Certificate

4W-1701-RA1

Company: **VALGOLD RESOURCES INC.**
Project: Hunter Mine
Attn: K. Jensen

Date: AUG-23-04

We hereby certify the following Assay of 50 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43001	0.02	-
43002	0.02	-
43003	0.01	-
43004	0.02	-
43005	0.50	-
43006	0.01	-
43007	0.55	0.52
43008	0.03	-
43009	0.02	-
43010	Nil	-
43011	0.02	-
43012	0.19	-
43013	0.62	0.65
43014	0.37	-
43015	0.13	-
43016	0.12	-
43017	0.04	-
43018	0.01	-
43019	0.01	-
43020	Nil	-
43021	Nil	-
43022	0.05	-
43023	Nil	-
43024	0.15	0.14
43025	0.04	-
43026	0.03	-
43027	0.01	-
43028	0.01	-
43029	0.01	-
43030	0.02	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 2

Assay Certificate

4W-1701-RA1

Company: **VALGOLD RESOURCES INC.**
Project: Hunter Mine
Attn: K. Jensen

Date: AUG-23-04

We hereby certify the following Assay of 50 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43031	0.03	-
43032	Nil	-
43033	Nil	-
43034	Nil	-
43035	0.01	-
43036	Nil	-
43037	0.01	-
43038	Nil	-
43039	0.01	-
43040	0.08	0.05
43041	0.04	-
43042	0.08	-
43043	Nil	-
43044	Nil	-
43045	Nil	-
43046	Nil	-
43047	0.04	-
43048	Nil	-
43049	Nil	-
43050	Nil	-
Blank	Nil	-
STD OxK18	3.31	-

Certified by  _____



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 3

Assay Certificate

4W-1696-RA1

Company: **VALGOLD RESOUARES INC.**
Project: Hunter Mine
Attn: K. Jensen

Date: AUG-23-04

We hereby certify the following Assay of 80 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43051	0.09	-
43052	Nil	-
43053	Nil	-
43054	0.18	-
43055	Nil	-
43056	Nil	-
43057	Nil	-
43058	0.01	-
43059	0.01	-
43060	0.01	-
43061	0.01	Nil
43062	Nil	-
43063	Nil	-
43064	Nil	-
43065	0.01	-
43066	0.02	-
43067	0.02	-
43068	Nil	-
43069	0.27	0.27
43070	Nil	-
43071	0.02	-
43072	Nil	-
43073	0.03	-
43074	Nil	-
43075	0.22	-
43076	Nil	-
43077	0.01	-
43078	Nil	-
43079	Nil	-
43080	0.09	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate


4W-1696-RA1

Company: **VALGOLD RESOUCRES INC.**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **AUG-23-04**

We hereby certify the following Assay of 80 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43081	0.10	-
43082	0.04	-
43083	0.01	-
43084	0.01	-
43085	0.01	-
43086	0.01	-
43087	0.42	-
43088	1.40	0.96
43089	0.04	-
43090	0.01	-
43091	0.01	-
43092	Nil	-
43093	0.01	-
43094	0.12	0.14
43095	0.01	-
43096	0.01	-
43097	0.01	-
43098	0.01	-
43099	0.03	-
43100	0.04	-
43101	Nil	-
43102	0.02	-
43103	0.01	-
43104	0.01	-
43105	0.02	-
43106	0.04	0.10
43107	0.02	-
43108	0.01	-
43109	0.04	-
43110	0.02	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate

4W-1696-RA1

Company: **VALGOLD RESOUCRES INC.**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **AUG-23-04**

We hereby certify the following Assay of 80 Core samples submitted AUG-05-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43111	0.01	-
43112	0.02	-
43113	0.03	-
43114	0.08	-
43115	0.14	-
43116	0.15	-
43117	0.63	0.59
43118	Nil	-
43119	Nil	-
43120	Nil	-
43121	0.01	-
43122	Nil	-
43123	Nil	-
43124	Nil	-
43125	Nil	-
43126	Nil	-
43127	0.01	-
43128	Nil	-
43129	Nil	-
43130	0.09	-
Blank	Nil	-
STD OXK18	3.57	-

Certified by Judy Ruser



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 3

Assay Certificate


4W-1859-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: SEP-01-04

We hereby certify the following Assay of 70 Core samples submitted AUG-12-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43131	Nil	-
43132	Nil	-
43133	0.03	-
43134	0.01	Nil
43135	Nil	-
43136	Nil	-
43137	0.02	-
43138	Nil	-
43139	0.03	-
43140	0.01	-
43141	Nil	-
43142	Nil	-
43143	Nil	-
43144	Nil	-
43145	Nil	-
43146	0.07	0.10
43147	Nil	-
43148	0.01	-
43149	0.01	-
43150	Nil	-
43151	0.04	-
43152	0.03	-
43153	0.01	-
43154	0.04	0.08
43155	Nil	-
43156	0.01	-
43157	Nil	-
43158	Nil	-
43159	Nil	-
43160	Nil	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate

4W-1859-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: SEP-01-04

We hereby certify the following Assay of 70 Core samples submitted AUG-12-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43161	0.01	-
43162	Nil	-
43163	Nil	-
43164	Nil	-
43165	0.01	-
43166	0.38	0.27
43167	0.02	-
43168	Nil	-
43169	Nil	-
43170	Nil	-
43171	Nil	-
43172	Nil	-
43173	0.02	-
43174	0.09	-
43175	Nil	-
43176	0.08	0.09
43177	0.02	-
43178	0.05	-
43179	0.03	-
43180	Nil	-
43181	0.29	-
43182	0.56	0.56
43183	Nil	-
43184	Nil	-
43185	Nil	-
43186	0.05	-
43187	0.01	-
43188	Nil	-
43189	0.02	-
43190	Nil	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate


4W-1859-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **SEP-01-04**

We hereby certify the following Assay of 70 Core samples submitted AUG-12-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
43191	Nil	-
43192	0.01	-
43193	Nil	0.01
43194	Nil	-
43195	0.01	-
43196	Nil	-
43197	Nil	-
43198	Nil	-
43199	0.01	-
43200	Nil	-
Blank	0.01	-
STD OxK18	3.55	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate

4W-1860-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **SEP-01-04**

We hereby certify the following Assay of 80 Core samples submitted AUG-12-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35031	Nil	-
35032	0.04	-
35033	0.03	-
35034	0.04	-
35035	0.07	-
35036	0.03	-
35037	0.01	-
35038	Nil	-
35039	Nil	-
35040	Nil	-
35041	Nil	-
35042	0.04	-
35043	Nil	-
35044	0.01	-
35045	Nil	-
35046	0.06	-
35047	Nil	-
35048	Nil	-
35049	Nil	-
35050	Nil	-
35051	0.01	-
35052	Nil	-
35053	0.11	-
35054	0.10	0.07
35055	0.03	-
35056	Nil	-
35057	0.01	-
35058	Nil	-
35059	Nil	-
35060	0.07	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate


4W-1860-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: SEP-01-04

We hereby certify the following Assay of 80 Core samples submitted AUG-12-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35061	Nil	-
35062	Nil	-
35063	0.01	-
35064	0.03	-
35065	0.04	-
35066	Nil	-
35067	0.04	-
35068	0.03	-
35069	Nil	-
35070	Nil	-
35071	Nil	-
35072	0.03	-
35073	0.01	-
35074	Nil	-
35075	0.02	-
35076	0.01	-
35077	0.09	-
35078	0.12	0.17
35079	Nil	-
35080	0.01	-
Blank	Nil	-
STD OxK18	3.39	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 3

Assay Certificate

4W-1861-RA1

Company: **VALGOLD RESOURCES**
Project: Hunter Mine
Attn: K. Jensen

Date: SEP-01-04

We hereby certify the following Assay of 70 Core samples submitted AUG-17-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35081	Nil	-
35082	Nil	-
35083	Nil	-
35084	Nil	0.01
35085	Nil	-
35086	Nil	-
35087	Nil	-
35088	0.02	-
35089	0.02	-
35090	Nil	-
35091	Nil	-
35092	Nil	-
35093	Nil	-
35094	Nil	-
35095	Nil	-
35096	Nil	-
35097	0.02	-
35098	Nil	-
35099	Nil	-
35100	Nil	-
35101	0.02	-
35102	0.03	0.05
35103	Nil	-
35104	Nil	-
35105	Nil	-
35106	Nil	-
35107	Nil	-
35108	Nil	-
35109	Nil	-
35110	Nil	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate

4W-1861-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **SEP-01-04**

We hereby certify the following Assay of 70 Core samples submitted AUG-17-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35111	Nil	-
35112	Nil	-
35113	Nil	-
35114	Nil	-
35115	Nil	-
35116	Nil	-
35117	Nil	Nil
35118	0.04	-
35119	0.03	-
35120	0.02	-
35121	0.01	-
35122	Nil	-
35123	0.02	-
35124	0.11	-
35125	0.10	-
35126	0.01	-
35127	0.76	-
35128	0.02	-
35129	0.05	-
35130	0.84	-
35131	1.18	1.13
35132	0.72	-
35133	0.89	1.16
35134	0.07	-
35135	0.03	-
35136	Nil	-
35137	0.03	-
35138	0.04	-
35139	0.02	-
35140	0.08	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate

4W-1861-RA1

Company: **VALGOLD RESOURCES**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **SEP-01-04**

We hereby certify the following Assay of 70 Core samples submitted AUG-17-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35141	0.08	-
35142	0.01	0.01
35143	Nil	-
35144	Nil	-
35145	Nil	-
35146	Nil	-
35147	0.12	0.16
35148	Nil	-
35149	0.01	-
35150	Nil	-
Blank	Nil	-
STD OxK18	3.52	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 1 of 3

Assay Certificate


4W-1965-RA1

Company: **VALGOLD RESOURCES LTD**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: SEP-09-04

We hereby certify the following Assay of 75 Core samples submitted AUG-27-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35151	0.01	-
35152	0.02	-
35153	Nil	-
35154	0.02	0.03
35155	0.09	-
35156	0.05	-
35157	0.43	0.42
35158	0.11	-
35159	0.03	-
35160	Nil	-
35161	0.02	-
35162	0.06	-
35163	0.09	0.10
35164	0.07	-
35165	0.08	-
35166	0.21	-
35167	0.21	-
35168	0.31	-
35169	0.30	-
35170	Nil	-
35171	Nil	-
35172	Nil	-
35173	Nil	-
35174	Nil	-
35175	0.01	0.03
35176	0.03	-
35177	Nil	-
35178	Nil	-
35179	Nil	-
35180	Nil	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 2 of 3

Assay Certificate


4W-1966-RA1

Company: **VALGOLD RESOURCES LTD**
Project: Hunter Mine
Attn: K. Jensen

Date: SEP-13-04

We hereby certify the following Assay of 85 Core samples submitted AUG-27-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35256	0.02	-
35257	0.02	-
35258	Nil	-
35259	Nil	-
35260	0.08	0.08
35261	Nil	-
35262	0.02	-
35263	Nil	-
35264	Nil	-
35265	0.10	-
35266	0.05	-
35267	0.05	-
35268	0.14	-
35269	0.09	-
35270	0.21	-
35271	0.25	0.18
35272	0.13	-
35273	Nil	-
35274	0.03	-
35275	Nil	-
35276	Nil	-
35277	Nil	-
35278	Nil	-
35279	Nil	-
35280	0.02	-
35281	Nil	Nil
35282	Nil	-
35283	Nil	-
35284	Nil	-
35285	0.05	-

Certified by 



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Page 3 of 3

Assay Certificate

4W-1966-RA1

Company: **VALGOLD RESOURCES LTD**
Project: Hunter Mine
Attn: K. Jensen

Date: SEP-13-04

We hereby certify the following Assay of 85 Core samples submitted AUG-27-04 by .

Sample Number	Au g/tonne	Au Check g/tonne
35286	0.03	-
35287	0.02	-
35288	0.01	-
35289	Nil	-
35290	Nil	-
35291	Nil	-
35292	Nil	-
35293	0.04	-
35294	0.06	-
35295	1.85	1.87
35296	0.10	-
35297	Nil	-
35298	Nil	-
35299	0.04	-
35300	Nil	-
35301	Nil	-
35302	Nil	-
35303	Nil	-
35304	Nil	-
35305	Nil	-
35306	Nil	-
35307	Nil	Nil
35308	Nil	-
35309	Nil	-
35310	Nil	-
Blank	Nil	-
STD OxK18	3.51	-

Certified by Judy Lewis



Established 1928

Swastika Laboratories Ltd

Assaying - Consulting - Representation

Metallic Assay Certificate

4W-2423-RM1

Company: **VALGOLD RESOURCES INC.**
Project: **Hunter Mine**
Attn: **K. Jensen**

Date: **NOV-04-04**

We hereby certify the following Metallic Assay of 21 Core samples submitted OCT-22-04 by .

Sample Number	Total		Assay Value Au		Total Weight Au		Metallic Au		Net Au	
	Wt (g)	+100 M Wt (g)	+100 (g/t)	-100 (g/t)	+100 (mg)	-100 (mg)	(oz/ton)	(g/t)	(oz/ton)	(g/t)
35401	1220.62	5.14	4.41	0.59	0.023	0.717	0.001	0.02	0.018	0.61
35402	1505.30	20.55	1.21	1.17	0.025	1.737	0.000	0.02	0.034	1.17
35403	681.64	21.75	0.01	0.01	0.000	0.007	0.000	0.00	0.000	0.01
35404	934.87	11.66	0.99	0.49	0.012	0.452	0.000	0.01	0.014	0.50
35405	1675.93	14.26	0.18	0.06	0.003	0.100	0.000	0.00	0.002	0.06
35406	1587.88	15.62	1.13	2.16	0.018	3.396	0.000	0.01	0.063	2.15
35407	693.20	19.89	0.09	0.52	0.002	0.350	0.000	0.00	0.015	0.51
35408	991.52	8.13	1.32	1.18	0.011	1.160	0.000	0.01	0.034	1.18
35409	1056.76	12.72	173.27	35.87	2.204	37.450	0.061	2.09	1.094	37.52
35410	1353.96	18.21	0.18	0.23	0.003	0.307	0.000	0.00	0.007	0.23
35411	2027.80	15.26	0.66	2.06	0.010	4.146	0.000	0.00	0.060	2.05
35412	1208.42	4.54	21.37	2.61	0.097	3.142	0.002	0.08	0.078	2.68
35413	1811.60	13.86	5.05	2.26	0.070	4.063	0.001	0.04	0.067	2.28
35414	939.21	19.45	136.24	29.59	2.650	27.216	0.082	2.82	0.927	31.80
35415	1741.18	16.41	0.01	0.05	0.000	0.086	0.000	0.00	0.001	0.05
35416	1153.35	16.87	1.50	1.13	0.025	1.284	0.001	0.02	0.033	1.14
35417	1181.83	20.54	3.16	3.33	0.065	3.867	0.002	0.05	0.097	3.33
35418	560.94	15.18	0.57	0.59	0.009	0.322	0.000	0.02	0.017	0.59
35419	1773.74	15.53	1.02	0.81	0.016	1.424	0.000	0.01	0.024	0.81
35420	400.20	5.24	0.61	1.72	0.003	0.679	0.000	0.01	0.050	1.71
35421	2348.75	18.41	0.64	0.25	0.012	0.533	0.000	0.01	0.007	0.25

Certified by 

ValGold Resources Ltd.
Hunter Mine Property
 Whitney Township, Porcupine Mining District
 Ontario
Surface Geology and Drilling

- HM04-39**
- 2004 Drill Hole
 - S-9 1986 Drill Hole
 - H-5 1985 Drill Hole
 - △ Sample Location

