

Report of Diamond Drilling

On the


Mine Centre Gold Properties

Northwestern Ontario

*Mining Claim K-3000815
& K-3000814*

Held by

Q-Gold (Ontario) Limited



Prepared by
Northwest Mineral Development Services

2 • 36631

Kenora, Ontario
December 5, 2007

Richard Beard, P.Eng

Report of Diamond Drilling on Mine Centre Gold Properties of Q-Gold Resources, Limited

Summary

During the months of March through May 2007, Q-Gold (Ontario) Limited drilled a number of diamond drill holes on the Company's Mine Centre gold properties located in the Kenora Mining Division. One hole, Q-07-03, collared on mining claim K-3000815, is reported herein. This hole tested for gold mineralization related to a geophysical anomaly over a major deformation zone in gabbroic rocks. Total depth of this hole was 391.00 metres. The core (NQ2 (50.8mm)) is stored on the property.

Location and Access

Q-Gold's Mine Centre properties are situated in unorganized territory in Northwestern Ontario, approximately 65 kilometres east of Fort Frances, Ontario (Key Map).

The village of Mine Centre is located near the northern edge of the claim groups. All claims lie within NTS 52-C/10, C / 15 and C/16 map sheets. Highway 11 passes through the property, and the drill site and the numerous known mineral deposits and showings on the property can be easily accessed by the Shoal Lake Road, which takes off from highway 11 one kilometre east of the village of Mine Centre, and runs south the full length of the claim groups.

Property

Q-Gold's property consists of a large group of un-patented mining claims, leases and patents. This property is described in a previous Company report, "Report on the Northwestern Ontario Gold & Base Metal Properties, Mine Centre Area, Rainy River District, Held by Hexagon Gold (Ontario) Ltd." by Northwest Mineral Development Services, March 20, 2003.

The property holdings of Q-Gold mentioned above include claims K-3000815 and K-3000814. The work reported herein consists of one drill hole, Q-07-03, that was collared on claim K- 3000815 and extends onto claim K-300814.

General Geology

Rocks of the Seine Bay - Bad Vermilion anorthosite complex underly most of claims K-3000815 and K-3000814. This anorthosite intrusion is in contact to the east with the Bad Vermilion tonalite/trondhjemite intrusive that is the host rock for most of the gold bearing veins on the property. Contact relationships between the anorthosite and the tonalite/trondhjemite are unclear, and the relative ages of the two bodies are uncertain. Some copper occurrences have been noted previously in the Bad Vermilion anorthosite intrusion.

Exploration Summary

Considerable work was carried out in the past by Nipigon Gold Ltd., on the McKenzie-Grey Property immediately to the southeast of claim K-3000815. The adjacent Nipigon Gold property was acquired by Q-Gold and, in 2006, Q-Gold carried out an airborne geophysical survey that included claim K-3000815 and 300814. This was followed up by ground geophysical surveys. The diamond drill hole reported herein, and several other drill holes that will be reported on at a later date, were drilled to test geophysical anomalies revealed by these surveys. It was believed that the anomalies might reflect a major deformation zone, the Finger Lake Fault, that may have served as the feeder for the MacKenzie-Grey gold deposits.

Diamond Drill Hole Q-07-03

Between March 14 and March 25, 2007, Q-Gold Resources Limited drilled diamond drill hole Q-07-03 that was collared on mining claim K-3000815. Most of the length of the hole (93.2%) was on claim K-3000814 but, at the bottom of the hole, extended (6.8%) onto claim K-3000814. Plan and section views of these holes as well as logs are part of this report.

Results

Hole Q-07-03 intersected anorthosite and other related mafic intrusive rocks throughout its length. Several shear zones and zones of schistose material were encountered, ranging from 3 to 20 metres in width.

Pyrite, usually less than 3%, was not uncommon, both in the mafic intrusive rocks and the shear zones. Several sections of quartz veining, some with tourmaline, were also noted. The highest gold value reported in the assaying was 0.03 grams per tonne.

Summary of Costs

Diamond Drilling George Downing Estate Drilling Ltd.	\$ 41,216
Assaying SGS Canada Inc.	\$ 1,157
Report Preparation Northwest Mineral Development Services	\$ <u>750</u>
Total:	\$ 43,123

Distribution of costs

K-3000815	93.2%	\$40,191
K-3000814	6.8%	\$ 2,932

References

- Beard, R.C. and Garratt, G.L. 1976. Gold Deposits of the Kenora - Fort Frances area, Districts of Kenora and Rainy River; Ontario Geological Survey, Mineral Deposits Circular 16, 46p.
- Blackburn, C.E., Johns, G.W., Ayer, J. and Davis, D.W. 1991. Wabigoon Subprovince; *in* Geology of Ontario, Ontario Geological Survey, Special Volume 4, Part 1, p. 303-381.
- Neilson, James N. & Bray, R.C.E., 1981. Feasibility of Small Scale Mining in Northwestern Ontario (parts of the Districts of Kenora, Rainy River and Thunder Bay; Ontario Geological Survey. Open File Report 5332 (Volumes 1 and 2).
- Poulsen, K.H. 1984. The Geological Setting of Mineralization in the Mine Centre - Fort Frances area, District of Rainy River; Ontario Geological Survey. Open File Report 5512, 126p.
- Poulsen, K.H. 2000. Precambrian Geology and Mineral Occurrences, Mine Centre - Fort Frances area; Ontario Geological Survey, Map 2525, scale 1:50,000.
- Poulsen, K.H. 2000. Archean Metallogeny of the Mine Centre - Fort Frances area; Ontario Geological Survey, Report 266, 121p.
- Stone, D., Halle, J. and Murphy, R. 1997a. Precambrian Geology, Mine Centre area; Ontario Geological Survey, Preliminary Map P.3372, scale 1:50 000.
- Tanton, T.L. 1936. Mine Centre Area, Rainy River District; Geological Survey of Canada, Map 334A, scale 1:31 680.
- Williams, H.R. 1991. Quetico Subprovince; *in* Geology of Ontario, Ontario Geological Survey, Special Volume 4, Part 1, p. 383-403.
- Wood, J., Dekker, J., Jansen, J.G., Keay, J.P. and Panagapko, D. 1980a. Mine Centre Area (west half), District of Rainy River; Ontario Geological Survey, Map P.2201, scale 1:15 840.
- Wood, J., Dekker, J., Jansen, J.G., Keay, J.P. and Panagapko, D. 1980b. Mine Centre Area (east half), District of Rainy River; Ontario Geological Survey, Map P.2202, scale 1:15 840.

Author of Report

Richard Beard

Northwest Mineral Development Services

Summer: Site 148, Comp. 9, RR #1

Kenora, ON, P9N 3W7

Winter: 4065 E. University Dr., #96

Mesa, AZ, USA, 85205

Supervisor of Work

Jack Bolen

521 Mowat Ave.

Fort Frances, ON


P9A 3M5

Submitted by

Richard Beard, P.Eng.

December 6, 2007

Q-GOLD (ONTARIO) LTD.: MINE CENTRE PROJECT AREA
Ontario, Canada



Drillhole Log

Q-Gold (Ontario) Ltd

Units Meters

Province/State	UTM East	Datum	Local Grid E	Azimuth Grid (°)	Length	Core Size	Date Started
Ontario	522731	NAD 83	800.00	125.00	391.00	NQ2	14/03/2007
District	UTM North	UTM Zone	Local Grid N	Azimuth Astro. (°)	Collar Survey Method		Date Completed
Kenora	5392503	15	1700.00				25/03/2007
Grid/Property	UTM Elevation	Drill Contractor		Dip (°)	Logged By		
Foley/Mine Centre	350.00	George Downing Estate		-50.00	Jack M. Bolen, B.Sc.		
Claim No.	<input type="checkbox"/> Pulsed	Geophysics Contractor		<input type="checkbox"/> Casing Pulled	Casing	<input type="checkbox"/> Plugged	Plug Depth
K-3000815					9.30		
Purpose				Core Storage			
Results				Comments			

Survey Tests

Lithology		Assays			Au	Ag	Cu
FROM	TO	SAMPLE #	FROM	TO	ppm	ppm	ppm
0.00	- 9.30	OVB	Overburden				
			Overburden, casing.				
9.30	- 33.50	8b, sch	Leucogabbro, anorthosite. Sch				
			Anorthositic. Massive, fine to medium grained. Moderately sheared, chloritic, strong calcite alteration as fracture fillings and interstitial. Plagioclase crystals often indistinct due to alteration. Occasional pyrite crystal. Brecciated on a decimetre scale, cemented with 1-3 mm white calcite veinlets.				
					22303	29.06	29.66
						0.00	
			<i>Mineralization:</i>				
9.30	- 33.50		Pyrite Trace				
			Massive, brecciated, plagioclase crystals often indistinct due to alteration				
29.06	- 29.66		Pyrite Disseminated 2.00%				
			Disseminated pyrite as fracture fills				
			<i>Alteration:</i>				
9.30	- 33.50		Chloritization, Calcareous Interstitial Strong				
			Calcite fracture filling, brecciated and cemented with 1-3 m white calcite veinlets				
29.06	- 29.66		Silicification Weak				
			<i>Structure:</i>				
9.30	- 33.50		Brecciated 0° to C/A				
29.06	- 29.66		Fracture 0° to C/A				
33.50	- 42.82	7, 8a	Coarse-Grained Mafic Intrusive Rocks. Gabbro, melagabbro				
			Clinopyroxene Diorite/Gabbro. Equigranular, fine to medium grained, massive, dark green. 60%, 1-5 mm clinopyroxene grains with a matrix of white plagioclase. Contacts sharp. Weakly fractured with calcite and pyrite fracture filling. Upper contact chilled, aphanitic for 0.5 m. Lower contact chilled fine grained to aphanitic for 0.6 m. Pyrite as disseminated grains, 1 cm blebs and fracture fillings, 3-4 %. Unit is magnetic throughout, no visible magnetite.				
					22304	33.50	34.50
						0.00	
					22305	34.50	35.50
						0.00	
					22306	35.50	36.50
						0.00	
					22307	36.50	37.50
						0.00	
					22308	37.50	38.50
						0.00	
					22309	38.50	39.50
						0.00	
					22310	39.50	40.50
						0.02	
					22311	40.50	41.50
						0.00	
					22312	41.50	42.10
						0.00	
					22313	42.10	42.82
						0.00	

<i>Lithology</i>		<i>Assays</i>					
<i>FROM</i>	<i>TO</i>	<i>SAMPLE #</i>	<i>FROM</i>	<i>TO</i>	<i>Au</i> <i>ppm</i>	<i>Ag</i> <i>ppm</i>	<i>Cu</i> <i>ppm</i>
<i>Mineralization:</i>							
33.50	- 42.82	Pyrite Disseminated 4.00% Pyrite as disseminated grains, 1 cm blebs and fracture fillings					
33.51	- 34.50	Pyrite Disseminated 2.00% Disseminated pyrite within fractures					
34.50	- 40.50	Pyrite Fracture Planes 4.00% Fine grained, massive					
35.50	- 36.50	Pyrite 3.00%					
36.50	- 37.50	Pyrite 2.00%					
37.50	- 38.50	Pyrite Disseminated 1.00%					
40.50	- 41.50	Pyrite Trace					
41.50	- 42.10	Pyrite Trace Very fine grained					
42.10	- 42.82	Pyrite Trace					
<i>Alteration:</i>							
33.50	- 42.82	Calcareous Fracture controlled					
33.51	- 34.50	Calcareous Fracture controlled					
34.50	- 40.50	Calcareous Fracture controlled					
42.10	- 42.82	Calcareous Cemented , Chloritization Weak					
<i>Structure:</i>							
33.50	- 42.82	Contact 53° to C/A Sharp					
33.51	- 34.50	Fracture 0° to C/A Moderate, calcite fracture filling					
33.52	- 34.50	Contact 0° to C/A Chilled margin					
34.50	- 40.50	Fracture 0° to C/A Moderate					
42.10	- 42.82	Fracture 0° to C/A					
42.82	- 47.75	8b	<u>Leucogabbro, anorthosite</u> Anorthositic Gabbro. 60% plagioclase, 40% chloritic clinopyroxene. Fractured with calcite fracture filling. Weakly sheared and altered. Trace pyrite.				
<i>Mineralization:</i>							
42.82	- 47.75	Pyrite Trace					
<i>Alteration:</i>							
42.82	- 47.75	Chloritization Weak, Calcareous Fracture controlled Weak					
<i>Structure:</i>							
42.82	- 47.75						

Lithology		Assays			Au	Ag	Cu
FROM	TO	SAMPLE #	FROM	TO	ppm	ppm	ppm
	Shearing 0° to C/A Weak						
47.75	- 51.70	2c, sch	<u>Quartz-chlorite schist, quartz-amphibole schist. Sch</u>				
			Shear Zone. Anorthositic Gabbro. Highly sheared. Highly calcareous. Mottled red, pink and green. Foliated. Tourmaline as crystals on foliation planes. 1-2% extremely fine pyrite.				
47.75	- 48.75	CCS	<u>Chlorite Carbonate Schist</u>				
		22314	47.75	48.75	0.00		
		22315	48.75	49.25	0.01		
		22316	49.25	50.25	0.02		
		22317	50.25	51.00	0.03		
		22318	51.00		0.02		
<i>Mineralization:</i>							
47.75	- 51.70	Pyrite 2.00%, Tourmaline Foliated					
		Fine pyrite					
47.76	- 48.75	Pyrite Disseminated 1.00%					
		Pyrite as disseminated grains and laminae on foliation planes					
48.75	- 49.25	Pyrite Grains 2.00%					
		Fine pyrite					
49.25	- 51.00	Pyrite Disseminated 1.00%					
51.00	- 51.70	Pyrite 2.00%					
		Fine pyrite					
<i>Alteration:</i>							
47.75	- 51.70	Calcareous Strong					
		Mottled red, pink, green					
47.76	- 48.75	Silicification Patchy					
		Red					
48.75	- 49.25	Chloritization Moderate, Calcareous Strong, Silicification Patchy Moderate					
		Less chloritic					
49.25	- 51.00	Chloritization Moderate, Sericitization Moderate, Carbonatization Moderate					
		20% chlorite, 20% quartz, 9% carbonate, 50% sericite					
51.00	- 51.70	Calcareous Weak, Silicification Bands					
		Pink quartz cherty					
<i>Structure:</i>							
47.75	- 51.70	Shearing 0° to C/A Highly					
47.76	- 51.70	Foliation 37° to C/A Strong					
51.70	- 133.40	8b	<u>Leucogabbro, anorthosite</u>				
			Anorthositic Gabbro. Medium grained, massive. 70% plagioclase, 30% clinopyroxene partially altered to chlorite. Trace pyrite.				

<i>Lithology</i>		<i>Assays</i>			<i>Au</i>	<i>Ag</i>	<i>Cu</i>
<i>FROM</i>	<i>TO</i>	<i>SAMPLE #</i>	<i>FROM</i>	<i>TO</i>	<i>ppm</i>	<i>ppm</i>	<i>ppm</i>
116.00	- 116.08	QV	<u>Quartz Vein</u>	22319	67.10	68.10	0.00
			No sulphides				
116.87	- 116.95	QV	<u>Quartz Vein</u>				
			White, 90 to CA, no sulphides				
117.21	- 117.41	QV	<u>Quartz Vein</u>				
			White, no sulphides				
123.30	- 125.35	QV	<u>Quartz Vein</u>				
			White, chloritic shear				
<i>Mineralization:</i>							
51.70	- 133.40		Pyrite Trace				
			70% plagioclase, 30% clinopyroxene, medium grained, massive				
67.10	- 68.10		Pyrite Disseminated 3.00%				
			Fine pyrite				
123.30	- 125.35		Pyrite Blebs 8.00%				
<i>Alteration:</i>							
51.70	- 133.40		Chloritization				
67.10	- 68.10		Silicification				
			Silicified zone, 90% quartz				
78.70	- 78.71		Chloritization Sheared				
81.80	- 81.81		Chloritization Sheared				
82.45	- 82.46		Chloritization Sheared				
100.50	- 100.68		Silicification Sheared				
123.30	- 125.35		Chloritization Sheared				
<i>Structure:</i>							
67.10	- 68.10		Contact 35° to C/A Sheared contacts				
78.70	- 78.71		Shearing 20° to C/A Chlorite filled shear				
81.80	- 81.81		Shearing 20° to C/A Chlorite filled shear				
82.45	- 82.46		Shearing 60° to C/A Chlorite filled shear				
100.50	- 100.68		Shearing 35° to C/A Silicified shear				
133.40	- 145.76	7	<u>Coarse-Grained Mafic Intrusive Rocks</u>				
			Pyroxenite. 80% dark green clinopyroxene, matrix of white to gray plagioclase. Contacts sharp. Contacts sheared, chilled for 2 m with strong chlorite alteration, weakly fractured with calcite cementing. Centre of unit is massive, fine grained. 2-4% pyrite disseminated throughout as well as fracture filling. Weakly magnetic, magnetite not visible.				

<i>Lithology</i>		<i>Assays</i>			<i>Au</i>	<i>Ag</i>	<i>Cu</i>
<i>FROM</i>	<i>TO</i>	<i>SAMPLE #</i>	<i>FROM</i>	<i>TO</i>	<i>ppm</i>	<i>ppm</i>	<i>ppm</i>
		22320	133.40	134.40	0.00		
		22321	134.40	135.40	0.00		
		22322	135.40	136.40	0.00		
		22323	136.40	137.40	0.00		
		22324	137.40	138.40	0.01		
		22325	138.40	139.40			
		22326	139.40	140.40	0.00		
		22327	140.40	141.40	0.00		
		22328	141.40	142.40	0.00		
		22329	142.40	143.40	0.01		
		22330	143.40	144.40	0.02		
		22331	144.40	145.20	0.00		
		22332	145.20	145.76	0.00		
<i>Mineralization:</i>							
133.40	- 145.76	Pyrite Disseminated 4.00%					
		Disseminated throughout and fracture filling, weakly magnetic, no magnetite visible					
133.41	- 134.40	Pyrite 2.00%					
134.40	- 135.40	Pyrite 1.00%					
135.40	- 139.40	Pyrite Trace					
		Massive pyroxenite					
139.40	- 140.40	Pyrite Trace					
		Massive pyroxenite					
140.40	- 143.40	Pyrite 1.00%					
		Massive pyroxenite					
143.40	- 144.40	Pyrite Disseminated 1.00%					
		Fine pyrite					
144.40	- 145.20	Pyrite 1.00%					
145.20	- 145.76	Pyrite 1.00%					
<i>Alteration:</i>							
133.40	- 145.76	Chloritization , Calcareous Cemented					
		Calcite cementing fractures					
133.41	- 134.40	Chloritization Fracture controlled , Calcareous Cemented , Silicification Cemented					
		Sheared contact					
134.40	- 135.40	Chloritization , Calcareous In Veins					
		Less chlortic, more massive 4-5% calcite veinlets					
143.30	- 144.40	Chloritization					
		Fine grained					

<i>Lithology</i>		<i>Assays</i>			<i>Au</i>	<i>Ag</i>	<i>Cu</i>	
<i>FROM</i>	<i>TO</i>	<i>SAMPLE #</i>	<i>FROM</i>	<i>TO</i>	<i>ppm</i>	<i>ppm</i>	<i>ppm</i>	
144.40	- 145.20	Chloritization Strong						
145.20	- 145.76	Chloritization , Calcareous Cemented						
<i>Structure:</i>								
133.40	- 145.76	Contact 35° to C/A Sharp, sheared, chilled for 2 m with strong chlorite alteration						
133.41	- 134.40	Contact 0° to C/A Sheared, fractured, chloritic, cemented with calcite an quartz						
145.20	- 145.76	Shearing 0° to C/A						
145.21	- 145.76	Fracture 0° to C/A Calcite cementing						
145.76	- 219.00	8b	<u>Leucogabbro, anorthosite</u>					
Anorthositic Gabbro. Medium grained, massive. 50%-60% gray plagioclase, 40-50% dark green clinopyroxene. Locally weakly sheared and chloritic, moderately calcareous. Occasional pyrite cube.								
168.30	- 168.31	QV	<u>Quartz Vein</u>					
			2 cm					
		22333	145.76	146.76			0.00	
		22334	199.85	201.00			0.00	
168.50	- 168.51	QV	<u>Quartz Vein</u>					
			White, 4 cm					
		22335	203.80	204.70			0.00	
		22336	204.70	205.30			0.00	
		22337	205.30	206.25			0.00	
171.60	- 171.61	QV	<u>Quartz Vein</u>					
			White, 4 cm					
172.70	- 172.71	QV	<u>Quartz Vein</u>					
			White, 10 cm					
194.50	- 194.69	MD	<u>Mafic Dike</u>					
			Contacts sharp, chloritic					
<i>Mineralization:</i>								
145.76	- 219.00	Pyrite Trace						
145.77	- 146.76	Pyrite Disseminated 2.00%						
171.60	- 171.61	Pyrite Trace						
172.70	- 172.71	Pyrite 4.00%						
199.85	- 201.00	Pyrite 1.00%						
203.80	- 204.70	Pyrite 1.00%						
204.70	- 205.30	Pyrite 4.00%						
205.30	- 206.25	Pyrite Trace						
<i>Alteration:</i>								
145.76	- 219.00	Chloritization , Calcareous Moderate						
145.77	- 146.76	Silicification						

Lithology		Assays			Au	Ag	Cu
FROM	TO	SAMPLE #	FROM	TO	ppm	ppm	ppm
194.50	- 194.69						
	Chloritization						
199.85	- 201.00						
	Silicification Patchy Moderate						
203.80	- 204.70						
	Chloritization , Calcareous						
204.70	- 205.30						
	Silicification Strong 80% quartz						
<i>Structure:</i>							
145.76	- 219.00						
	Shearing 0° to C/A						Locally weakly sheared
145.77	- 146.76						Weak
194.50	- 194.69						Contact 40° to C/A Sharp
199.85	- 201.00						Shearing 0° to C/A Weak
203.80	- 204.70						Shearing 0° to C/A Weak
219.00	- 232.60						SZ <u>Shear Zone</u>
	Sheared, highly calcareous, plagioclase become indistinct, foliated, trace pyrite, strong chlorite alteration.						
<i>Mineralization:</i>							
219.00	- 232.60						Pyrite Trace
228.05	- 228.22						Pyrite 1.00%
232.00	- 232.60						Pyrite 2.00%
<i>Alteration:</i>							
219.00	- 232.60						Chloritization Strong, Calcareous Strong
228.05	- 228.22						Sericitization , Silicification Weak
232.00	- 232.60						Silicification
<i>Structure:</i>							
219.00	- 232.60						Foliation 44° to C/A
219.01	- 232.60						Shearing 0° to C/A
228.05	- 228.22						Shearing 0° to C/A
232.60	- 261.65						8b <u>Leucogabbro, anorthosite</u>
	Anorthositic Gabbro. Massive, medium grained. 60% white to gray plagioclase, 40% dark green clinopyroxene. Occasional pyrite crystal. Minor fractures.						
<i>Mineralization:</i>							
		22338	228.05	228.22	0.00		
		22339	232.00	232.60	0.00		
		22340	261.10	262.00	0.00		

Lithology		Assays			Au	Ag	Cu			
FROM	TO	SAMPLE #	FROM	TO	ppm	ppm	ppm			
300.50	- 302.50	Chloritization , Calcareous Strong Schistose								
301.00	- 302.00	Chloritization Strong, Calcareous Strong								
304.00	- 326.20	Chloritization Spots Fine grained structures up to 10 cm, may be digested clasts								
<i>Structure:</i>										
277.00	- 280.00	Shearing 0° to C/A Weak								
282.00	- 284.00	Shearing 0° to C/A Weak								
289.00	- 291.00	Shearing 0° to C/A Weak								
300.50	- 302.50	Shearing 0° to C/A								
301.00	- 302.00	Fracture 0° to C/A Weak								
326.00	- 346.00	2c, 8, sch <u>Quartz-chlorite schist, quartz-amphibole schist. Metamorphosed Gabbroic Rocks. Sch</u> Shear Zone. Strong calcite, chloritic slips, plagioclase grains are practically non-existent, which is gradational over 2 m from the above unit. Localized brecciation, minor quartz/ankerite/calcite veining. Traces of pyrite.								
336.55	- 340.00	8a <u>Gabbro, melagabbro</u> Unit becomes more massive grading into coarse grained anorthositic gabbro					22343	326.20	327.20	0.00
						22344	326.21	327.20	0.00	
						22345	330.55	331.55	0.00	
						22346	331.55	332.55	0.00	
						22347	332.55	333.55	0.00	
						22348	333.55	334.55	0.00	
						22349	334.55	335.55	0.00	
						22350	335.55	336.55	0.00	
<i>Mineralization:</i>										
326.20	- 346.00	Pyrite Trace 2.00% Plagioclase grains non-existent, localized brecciation								
326.21	- 327.20	Pyrite Disseminated 1.00% Brecciated								
330.55	- 331.55	Pyrite 4.00%, Marcasite 5.00%								
331.55	- 334.55	Pyrite Trace								
334.55	- 335.55	Pyrite Trace								
335.55	- 336.55	Pyrite Trace Massive								
<i>Alteration:</i>										
326.20	- 346.00	Chloritization Stringers , Calcareous Strong, Ankerite In Veins								

<i>Lithology</i>		<i>Assays</i>			<i>Au</i>	<i>Ag</i>	<i>Cu</i>
<i>FROM</i>	<i>TO</i>	<i>SAMPLE #</i>	<i>FROM</i>	<i>TO</i>	<i>ppm</i>	<i>ppm</i>	<i>ppm</i>
	Quartz/ankerite/calcite veining						
326.21	- 327.20 Chloritization , Calcareous Strong						
330.55	- 331.55 Calcareous In Veins Strong						
331.55	- 334.55 Chloritization Cemented , Calcareous Strong						
334.55	- 335.55 Silicification Patchy Moderate						
<i>Structure:</i>							
331.55	- 334.55 Fracture 0° to C/A						
332.55	- 334.55 Foliation 47° to C/A						
346.00	- 355.00 8a <u>Gabbro, melagabbro</u>						
	Dark green, fine grained, weakly sheared, strongly calcareous throughout as interstitial grains, weakly chloritic.						
		22351	346.00	347.00	0.00		
		22352	347.00	348.00	0.00		
		22353	348.00	349.00	0.00		
		22354	349.00	350.00	0.00		
		22355	350.00	351.00	0.00		
		22356	351.00	352.00	0.00		
		22357	352.00	353.00	0.00		
		22358	353.00	354.00	0.00		
		22359	354.00	355.00	0.00		
<i>Mineralization:</i>							
346.00	- 355.00 Pyrite Disseminated 0.50% Finely disseminated pyrite						
<i>Alteration:</i>							
346.00	- 355.00 Chloritization Weak, Calcareous Interstitial Strong						
346.01	- 355.00 Calcareous Massive						
353.00	- 354.00 Chloritization						
<i>Structure:</i>							
346.00	- 355.00 Shearing 0° to C/A Weak						
346.01	- 355.00 Foliation 44° to C/A						
346.02	- 355.00 Shearing 0° to C/A Weak						
353.00	- 354.00 Shearing 0° to C/A Increasing						

<i>Lithology</i>		<i>Assays</i>			<i>Au</i>	<i>Ag</i>	<i>Cu</i>
<i>FROM</i>	<i>TO</i>	<i>SAMPLE #</i>	<i>FROM</i>	<i>TO</i>	<i>ppm</i>	<i>ppm</i>	<i>ppm</i>
355.00	- 391.00	8b	<u>Leucogabbro, anorthosite</u>				
			Anorthositic Gabbro. Variable grain size, coarse to fine. 60% plagioclase, 40% clinopyroxene. Locally weakly sheared resulting in a finer grain size and weakly chloritic. Weakly calcareous throughout.				
385.55	- 386.45	QTV	<u>Quartz Tourmaline Vein</u>				
			5% veinlets in sheared and brecciated chloritic gabbro				
386.45	- 387.00	QTV	<u>Quartz Tourmaline Vein</u>				
			5% veinlets				
389.00	- 391.00	CV	<u>Calcite Vein</u>				
			Calcite veinlets cementing				
<i>Mineralization:</i>							
372.00	- 372.01		Chalcopyrite Blebs 1 cm bleb of chalcopyrite				
385.55	- 386.45		Pyrite Trace , Tourmaline Vein				
<i>Alteration:</i>							
355.00	- 391.00		Chloritization Weak, Calcareous Throughout Weak				
389.00	- 391.00		Calcareous Cemented Weak				
<i>Structure:</i>							
355.00	- 391.00		Shearing 0° to C/A Weak, local, resulting in a finer grain size				
385.55	- 386.45		Shearing 0° to C/A				
386.45	- 387.00		Fracture 0° to C/A Decreasing				
389.00	- 391.00		Brecciated 0° to C/A Weak				
		22360	385.55	386.45	0.00		
		22361	386.45	387.00	0.00		

02/05/2002 02:37 9287790107

G-GOLD RESOURCES

PAGE 04



Final : RL28306

Page 2 of 3

Element Method Det.Lim. Units	Au	Au (AR)	Au	Au (R)
	FAA303 0.01 g/t	FAA303 0.01 g/t	FAA303 0.001 g/t	FAA303 0.001 g/t
22309	<0.01	<0.01	<0.001	<0.001
22310	0.02	-	<0.001	-
22311	<0.01	-	<0.001	-
22312	<0.01	-	<0.001	-
22313	<0.01	-	<0.001	-
22314	<0.01	-	<0.001	-
22315	0.01	-	<0.001	-
22316	0.02	-	<0.001	-
22317	0.02	-	<0.001	-
22318	0.02	-	<0.001	-
22319	<0.01	-	<0.001	-
22320	<0.01	-	<0.001	-
22321	<0.01	-	<0.001	-
22322	<0.01	-	<0.001	-
22323	<0.01	-	<0.001	-
22324	0.01	-	<0.001	-
22325	<0.01	-	<0.001	-
22326	<0.01	-	<0.001	-
22327	<0.01	-	<0.001	-
22328	<0.01	-	<0.001	-
22329	<0.01	-	<0.001	-
22330	0.02	-	<0.001	-
22331	<0.01	-	<0.001	-
22332	<0.01	-	<0.001	-
22333	<0.01	-	<0.001	-
22334	<0.01	<0.01	<0.001	<0.001
22335	<0.01	-	<0.001	-
22336	<0.01	-	<0.001	-
22337	<0.01	-	<0.001	-
22338	<0.01	-	<0.001	-
22339	<0.01	-	<0.001	-
22340	<0.01	-	<0.001	-
22341	<0.01	-	<0.001	-
22342	<0.01	-	<0.001	-
22343	<0.01	-	<0.001	-
22344	<0.01	-	<0.001	-
22345	<0.01	-	<0.001	-
22346	<0.01	-	<0.001	-
22347	<0.01	-	<0.001	-
22348	<0.01	-	<0.001	-
22349	<0.01	-	<0.001	-
22350	<0.01	-	<0.001	-
22351	<0.01	-	<0.001	-
22352	<0.01	-	<0.001	-
22353	<0.01	-	<0.001	-
22354	<0.01	-	<0.001	-
22355	<0.01	-	<0.001	-
22356	<0.01	-	<0.001	-
22357	<0.01	-	<0.001	-

Q-07-03

The data reported on this certificate of analysis represents the sample submitted to SGS Minerals Services. Reproduction of this analytical report, in full or in part, is prohibited without prior written approval.

SGS Canada Inc. Minerals Services 18A Young St. PO Box 1249 Red Lake ON P0V 2M0 (807) 727-2938 (807) 727-5188 www.sgs.ca

Member of the SGS Group (Société Générale de Surveillance)

02/05/2002 02:37 9287790187

Q-GOLD RESOURCES

PAGE 03



Q-07-02

Final: RL28182

Page 3 of 3

Element Method Det.Lim. Units	Au	Au (AR)	Au	Au (R)
	FAA303 0.01 G/T	FAA303 0.01 G/T	FAA303 0.001 OZ/T	FAA303 0.001 OZ/T
22264	0.01	-	<0.001	-
22265	<0.01	-	<0.001	-
22270	<0.01	<0.01	<0.001	<0.001
22271	<0.01	-	<0.001	-
22272	<0.01	-	<0.001	-
22273	<0.01	-	<0.001	-
22274	<0.01	-	<0.001	-
22275	<0.01	-	<0.001	-
22276	<0.01	-	<0.001	-
22277	<0.01	-	<0.001	-
22278	<0.01	-	<0.001	-
22279	<0.01	-	<0.001	-
22280	<0.01	<0.01	<0.001	<0.001
22281	<0.01	-	<0.001	-
22282	<0.01	-	<0.001	-
22283	<0.01	-	<0.001	-
22284	<0.01	-	<0.001	-
22285	<0.01	-	<0.001	-
22286	<0.01	-	<0.001	-
22287	<0.01	-	<0.001	-
22288	<0.01	-	<0.001	-
22289	<0.01	-	<0.001	-
22290	<0.01	<0.01	<0.001	<0.001
22291	<0.01	-	<0.001	-
22292	<0.01	-	<0.001	-
22293	<0.01	-	<0.001	-
22294	<0.01	-	<0.001	-
22295	<0.01	-	<0.001	-
22296	<0.01	-	<0.001	-
22297	<0.01	-	<0.001	-
22298	<0.01	-	<0.001	-
22299	<0.01	-	<0.001	-
22300	<0.01	<0.01	<0.001	<0.001
22301	<0.01	-	<0.001	-
22302	<0.01	-	<0.001	-
22303	<0.01	-	<0.001	-
22304	<0.01	-	<0.001	-
22305	<0.01	-	<0.001	-
22306	<0.01	-	<0.001	-
22307	<0.01	-	<0.001	-
22308	<0.01	-	<0.001	-

Q-07-03

The data reported on this certificate of analysis represents the sample submitted to SGS Minerals Services. Reproduction of this analytical report, in full or in part, is prohibited without prior written approval.

SGS Canada Inc. Minerals Services 18A Young St. PO Box 1349 Red Lake ON P0V 2M0 t(807) 727-2938 f(807) 727-3183 www.sgs.ca

Member of the SGS Group (Société Générale de Métallurgie)

02/05/2002 02:37 9287790107

Q-GOLD RESOURCES

PAGE 05



Page 3 of 3

Final : RL28306

Element Method Det.Lim. Units	Au FAA303 0.01 G/T	Au (AR) FAA303 0.01 G/T	Au FAA303 0.001 OZ/T	Au (R) FAA303 0.001 OZ/T
	22368	<0.01	-	<0.001
22369	<0.01	-	<0.001	-
22380	<0.01	<0.01	<0.001	<0.001
22381	<0.01	-	<0.001	-

The data reported on this certificate of analysis represents the sample submitted to SGS Minerals Services. Reproduction of this analytical report, in full or in part, is prohibited without prior written approval.

SGS Canada Inc. Minerals Services 18A Young St. PO Box 1348 Red Lake ON P0V 2M0 t(807) 727-2835 f(807) 727-3183 www.sgs.ca

Member of the SGS Group (Société Canadienne de Services)