



Drillhole Log

Q-Gold (Ontario) Ltd

| | | | | | | | | | |
|---|----|----------------------------|---------------------------|--|---------------------|-----------------------------|---|---------------------------------|--|
| Province/State | | Co-ordinate System | | Grid/Property | | Hole Type | Length | Date Started | |
| Ontario | | UTM NAD83 Canada Zone 15 | | MG Grid | | Exploration hole | 177.00 | 11/13/2010 | |
| District | | UTM North | UTM East | Local Grid E | Local Grid N | Collar Survey Method | | Date Completed | |
| Kenora | | 5392388 | 523543 | -60.00 | 120.00 | MNR DEM | | | |
| Project | | UTM Elevation | Azimuth Astro. (°) | Azimuth Grid (°) | Dip (°) | Drill Contractor | | Date Logged | |
| McKenzie-Gray Project | | 349.00 | 43.80 | | -62.00 | C3 Drilling Company | | 11/26/2010 | |
| Area | | Claim No. | NTS Sheet | Supervised By | | Logged By | Verified | | |
| Mine Center | | K-475273 | 052C10 | | | Vincent Scime | <input type="checkbox"/> | | |
| Zone/Prospect | | Assessment Rpt. No. | Core Storage | | Plug Depth | Makes Water | Capped | Environmental Inspection | |
| | | | Fort Frances Office | | | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| Core Size (1) | NQ | 174 | Casing Pulled | Casing (1) | 3.00 | Steel | Plugged | Pulsed | |
| (2) | | | <input type="checkbox"/> | (2) | | | <input type="checkbox"/> | <input type="checkbox"/> | |
| Purpose | | | | Results | | | Comments | | |
| Intersect deeper section of MG and East Veins | | | | Intersected quartz rich zones and mineralized quartz veins | | | Drill Log Updated by D. Tortosa Dec 2010. NOTE: For samples with no assay data see ICP-MS multi-element results. | | |

Survey Tests

| Distance | Grid Azimuth (°) | | Astro. Azimuth (°) | | Dip (°) | | Use Test | Survey Method | Mag. Field (nT) | Comments |
|----------|------------------|-------|--------------------|-------|----------|-------|-------------------------------------|---------------|-----------------|------------------------------------|
| | Original | Final | Original | Final | Original | Final | | | | |
| 177.00 | | | 44.7 | | -60.4 | | <input checked="" type="checkbox"/> | Reflex EZ | | Dip -60.2 Azimuth 43.8 at 9 metres |

| <i>Lithology</i> | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> |
|------------------|-------------|-----------|--|------------|------------|-----------|-----------|-----------|
| <i>From</i> | <i>To</i> | | | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| <i>Sample #</i> | <i>From</i> | <i>To</i> | <i>Len.</i> | | | | | |
| 0.00 | - | 3.00 | OVB | | | | | |
| | | | <u>Overburden</u> | | | | | |
| 3.00 | - | 68.10 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> | | | | | |
| 3.00 | - | 5.70 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> massive, med grained weakly alt, plag xls have slight pink tint, minor sulphide | | | | | |
| 5.70 | - | 9.20 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> fine grained dark grey phase, rapid transition from previous becoming vfg, pink, siliceous below about 6.4m to 8.4; fine to med dark grey to end of section; 1-2% diss py | | | | | |
| 9.20 | - | 40.00 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> med grained, weak to mod alt, plag xls have pink/red tint, 1-2% diss py; distinct contact with previous @ 55-60deg to CA | | | | | |
| 40.00 | - | 58.70 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> similar to above but without pink tint in plag xls, med garined, grey, mod alt, locally strong; gradual transition from previous, contact is somewhat arbitrary; a few qtz-carb-tour stringers <1cm and fracture fillings @ 20deg to CA; 1-2% diss py | | | | | |
| 58.70 | - | 68.10 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> as 9.2-40m, med grained with pink tint in plag xls, gradual transition from previous across approx 1m; weak to mod alt becoming strong in bottom 2m | | | | | |
| 68.10 | - | 68.50 | FZ | | | | | |
| | | | <u>Fault Zone</u> strong pink/red tint; two 1-3cm sections of muddy breccia @ 20 and 40 deg to CA | | | | | |
| 68.50 | - | 75.50 | 9i | | | | | |
| | | | <u>Trondhjemite, altered</u> | | | | | |
| 586 | 68.50 | 69.00 | 0.50 | | | | | |
| 587 | 69.00 | 69.70 | 0.70 | | | | | |
| 588 | 69.70 | 70.45 | 0.75 | | | | | |
| 589 | 70.45 | 71.30 | 0.85 | | | | | |
| 591 | 71.30 | 72.20 | 0.90 | | | | | |
| 592 | 72.20 | 72.60 | 0.40 | | | | | |
| 593 | 72.60 | 72.85 | 0.25 | 0.015 | 1.5 | 0.005 | 0.14 | |
| 594 | 72.85 | 73.35 | 0.50 | | | | | |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> | | | |
|------------------|-----------|-----------|---|-----------------|-------------|-----------|-------------|------------|------------|----------|----------|----------|
| <i>From</i> | <i>To</i> | | | <i>Sample #</i> | <i>From</i> | <i>To</i> | <i>Len.</i> | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| 68.50 | - 72.85 | 9i | Trondhjemite, altered mod to strong alt; several narrow 1cm qtz and qtz-carb stringers @15-20deg to CA with coarse py xls; 72-6-72.85m carb fracture fillings with 2% coarse blebs cpy | | | | | | | | | |
| 75.50 | - 106.20 | 9c | Trondhjemite (quartz porphyritic) | 595 | 86.60 | 86.90 | 0.30 | | | | | |
| | | | | 596 | 92.80 | 93.80 | 1.00 | | | | | |
| | | | | 597 | 93.80 | 94.50 | 0.70 | | | | | |
| | | | | 598 | 94.50 | 95.00 | 0.50 | | | | | |
| | | | | 599 | 95.00 | 96.00 | 1.00 | | | | | |
| | | | | 601 | 96.00 | 97.00 | 1.00 | | | | | |
| | | | | 602 | 97.00 | 98.10 | 1.10 | | | | | |
| | | | | 603 | 98.10 | 99.00 | 0.90 | | | | | |
| | | | | 604 | 99.00 | 100.25 | 1.25 | | | | | |
| | | | | 605 | 100.25 | 100.75 | 0.50 | | | | | |
| | | | | 606 | 100.75 | 101.40 | 0.65 | | | | | |
| | | | | 607 | 101.40 | 102.00 | 0.60 | | | | | |
| | | | | 608 | 102.00 | 106.60 | 4.60 | | | | | |
| 75.50 | - 86.60 | 9c | Trondhjemite (quartz porphyritic) massive, med to coarse grained, leucocratic, local weak to mod alt, becoming mod altered in through bottom 2m; 1-2 % diss py with a few small patches <5cm containing up to 10% py blebs | | | | | | | | | |
| 86.60 | - 86.90 | 9i | Trondhjemite, altered strong alt, 12-15% py xls and blebs | | | | | | | | | |
| 86.90 | - 89.50 | 9c | Trondhjemite (quartz porphyritic) fin to vfg, grey, massive, 1% diss py; distinct upper contact @45-50deg to CA | | | | | | | | | |
| 89.50 | - 92.80 | 9c | Trondhjemite (quartz porphyritic) massive, coarse grained, leucocratic | | | | | | | | | |
| 92.80 | - 102.65 | 9c | Trondhjemite (quartz porphyritic) as above but contains several qtz and qtz-carb-tour stringers up to 3cmm @25deg to CA, offset by fractures @70-80deg to CA; 6-8% fine diss py in wallrock, coarse xls in qtz; local weak to mod alt | | | | | | | | | |
| 102.65 | - 103.80 | 9c | Trondhjemite (quartz porphyritic) fine to vfine grained, grey-green with a 10cm section intensely alt, aplitic; sharp upper contact @ 60deg to CA | | | | | | | | | |
| 103.80 | - 106.20 | 9c | Trondhjemite (quartz porphyritic) massive, med grained, leucocratic with pink and/or pale yellow-green, weakly sausseritized plag xls; minor py | | | | | | | | | |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> |
|------------------|-----------|-----------------|--|-------------|------------|------------|-----------|-----------|--------------------------------|
| <i>From</i> | <i>To</i> | | | <i>Len.</i> | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| | | <i>Sample #</i> | <i>From</i> | <i>To</i> | | | | | |
| 106.20 | - 106.60 | 9i | <u>Trondhiemite, altered</u> mod alt, 1-2% diss py | | | | | | |
| 106.60 | - 106.75 | MQV | <u>Mineralized Quartz Vein</u> 15cm qtz-tour-carb-chl vein @65-70deg to CA; 4-6% coarse xls py, 1-2% small blebs cpy, few small specks gal? | | 609 | 106.60 | 106.75 | 0.15 | 82.82 13.95 0.0075 0.31 0.0175 |
| 106.75 | - 131.20 | 9i | <u>Trondhiemite, altered</u> variably altered, mostly weak-mod with narrow sections of mod-strong alt and a few sections up to 2.5m unaltered; scattered qtz-tour-carb stringers @ 60-70deg to CA | | 611 | 106.75 | 107.25 | 0.50 | |
| | | | | | 612 | 107.25 | 107.75 | 0.50 | |
| | | | | | 613 | 107.75 | 108.60 | 0.85 | |
| | | | | | 614 | 108.60 | 108.90 | 0.30 | |
| | | | | | 615 | 108.90 | 110.25 | 1.35 | |
| | | | | | 616 | 110.25 | 110.70 | 0.45 | |
| | | | | | 617 | 110.70 | 111.65 | 0.95 | |
| | | | | | 618 | 111.65 | 112.30 | 0.65 | |
| | | | | | 619 | 112.30 | 113.60 | 1.30 | |
| | | | | | 621 | 113.60 | 114.80 | 1.20 | |
| | | | | | 622 | 114.80 | 115.40 | 0.60 | |
| | | | | | 623 | 115.40 | 116.70 | 1.30 | |
| | | | | | 624 | 116.70 | 117.70 | 1.00 | |
| | | | | | 625 | 117.70 | 118.65 | 0.95 | |
| | | | | | 626 | 118.65 | 119.80 | 1.15 | |
| | | | | | 627 | 119.80 | 120.80 | 1.00 | |
| | | | | | 628 | 120.80 | 121.80 | 1.00 | |
| | | | | | 629 | 121.80 | 122.75 | 0.95 | |
| | | | | | 631 | 122.75 | 123.30 | 0.55 | |
| | | | | | 632 | 123.30 | 124.20 | 0.90 | |
| | | | | | 633 | 124.20 | 125.15 | 0.95 | |
| | | | | | 634 | 125.15 | 126.60 | 1.45 | |
| | | | | | 635 | 126.60 | 127.60 | 1.00 | |
| | | | | | 636 | 127.60 | 128.60 | 1.00 | |
| | | | | | 637 | 128.60 | 129.90 | 1.30 | |
| | | | | | 638 | 129.90 | 131.20 | 1.30 | |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> |
|------------------|-----------|---|--|-------------|------------|------------|-----------|-----------|-----------|
| <i>From</i> | <i>To</i> | | | <i>Len.</i> | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| 106.75 | - 107.25 | 9i Trondhemite, altered first 20cm is fg, dark grey phase, remainder is weak alt 9i; 1-2 % diss py | | | | | | | |
| 107.25 | - 107.75 | 9i Trondhemite, altered mod alt, qtz-carb fractures @ 60deg to CA; 4-6% diss py | | | | | | | |
| 107.75 | - 108.60 | 9i Trondhemite, altered weak alt, 2cm qtz-carb-tour stringer @ 10deg to CA; 4-6% diss py | | | | | | | |
| 108.60 | - 108.90 | 9i Trondhemite, altered strong alt, qtz-carb fractures @ 65-70deg to CA; 6-8% diss py | | | | | | | |
| 108.90 | - 110.25 | 9c Trondhemite (quartz porphyritic) Trondhemite | | | | | | | |
| 110.25 | - 110.70 | 9i Trondhemite, altered mod to strong alt, two qtz-carb stringers, @ 70deg to CA with coarse py xls; 4-6% diss py | | | | | | | |
| 110.70 | - 111.65 | 9i Trondhemite, altered weak to mod alt, plag xls have slight pink tint, 1cm qtz-carb-tour stringer @ 10deg to CA, 4-6% diss py | | | | | | | |
| 111.65 | - 112.30 | 9i Trondhemite, altered strong alt, qtz-carb-tour stringers and fracture fillings @ 65deg to CA, 4-6% f diss py, a few small blebs | | | | | | | |
| 112.30 | - 114.80 | 9c Trondhemite (quartz porphyritic) 9c, variable to mod 9i near qtz stringers, plag xls have pink tint; 1-2% diss py; a few qtz-carb-tour stringers @ 15deg to CA | | | | | | | |
| 114.80 | - 115.40 | 9i Trondhemite, altered fg, dark grey phase, upper contact marked by 1cm qtz-carb-tour-chl vein with coarse py xls | | | | | | | |
| 115.40 | - 116.70 | 9i Trondhemite, altered weak-mod alt, plag xls have slight pink tint in less altered sections; 1-2% diss py | | | | | | | |
| 116.70 | - 118.65 | 9i Trondhemite, altered strong alt, qtz stringers @ 20-25deg to CA with coarse py xls crosscut by qtz-carb fractures and qtz-carb-tour stringers @ 55-60deg to CA | | | | | | | |
| 118.65 | - 119.80 | 9c Trondhemite (quartz porphyritic) 9c, weakly sausseritized pale green plag xls | | | | | | | |
| 119.80 | - 122.75 | 9i Trondhemite, altered mod alt, scattered qtz and qtz-carb-tour stringers @ 50-60deg to CA, 2-4% diss py | | | | | | | |
| 122.75 | - 123.30 | 9c Trondhemite (quartz porphyritic) Trondhemite | | | | | | | |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> |
|------------------|-----------|------------|---|-----|------------|------------|-----------|-----------|-----------|
| <i>From</i> | <i>To</i> | | | | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| 123.30 | - 125.15 | 9i | Trondhjemite, altered mod alt, several qtz and qtz-carb-tour stringers @ 60-65deg to CA; 2-4% diss py | | | | | | |
| 125.15 | - 126.60 | 9c | Trondhjemite (quartz porphyritic) 9c, leucocratic, 2-4 % diss py; good examples of the the two vein sets x-cutting each other | | | | | | |
| 126.60 | - 129.90 | 9i | Trondhjemite, altered mod to strong alt, scattered qtz-carb-tour stringers @ 60-65deg to CA; 4-6% diss py | | | | | | |
| 129.90 | - 131.20 | 9i | Trondhjemite, altered strong alt, silicified, sericitic, strong fol'n @30-35deg to CA; 6-8% v fine diss py | | | | | | |
| 131.20 | - 155.10 | QRZ | <u>Quartz Rich Zone</u> | | | | | | |
| | | | | 639 | 131.20 | 131.70 | 0.50 | | |
| | | | | 641 | 131.70 | 132.30 | 0.60 | | |
| | | | | 642 | 132.30 | 132.95 | 0.65 | | |
| | | | | 643 | 132.95 | 134.00 | 1.05 | 0.66 | 88.8 |
| | | | | 644 | 134.00 | 134.85 | 0.85 | 0.015 | 8.5 |
| | | | | 645 | 134.85 | 135.70 | 0.85 | 0.015 | 26.9 |
| | | | | 646 | 135.70 | 136.65 | 0.95 | 0.015 | 2.3 |
| | | | | 647 | 136.65 | 137.50 | 0.85 | 0.015 | 2 |
| | | | | 648 | 137.50 | 138.20 | 0.70 | 0.015 | 3.2 |
| | | | | 649 | 138.20 | 138.95 | 0.75 | 0.015 | 2.5 |
| | | | | 651 | 138.95 | 139.50 | 0.55 | 0.1 | 50.7 |
| | | | | 652 | 139.50 | 139.85 | 0.35 | 0.04 | 7.6 |
| | | | | 653 | 139.85 | 140.30 | 0.45 | 0.015 | 10.2 |
| | | | | 654 | 140.30 | 140.95 | 0.65 | 0.015 | 31.7 |
| | | | | 655 | 140.95 | 141.25 | 0.30 | 0.015 | 4.4 |
| | | | | 656 | 141.25 | 141.85 | 0.60 | 0.04 | 18.5 |
| | | | | 657 | 141.85 | 142.35 | 0.50 | 0.015 | 67.9 |
| | | | | 658 | 142.35 | 143.00 | 0.65 | 0.015 | 1.9 |
| | | | | 659 | 143.00 | 144.10 | 1.10 | 0.015 | 2.3 |
| | | | | 661 | 144.10 | 145.75 | 1.65 | 0.015 | 2.1 |
| | | | | 662 | 145.75 | 146.30 | 0.55 | 2.23 | 728.9 |
| | | | | 663 | 146.30 | 147.65 | 1.35 | 0.015 | 3.6 |
| | | | | 664 | 147.65 | 148.35 | 0.70 | 0.015 | 2.3 |
| | | | | 665 | 148.35 | 148.90 | 0.55 | 0.015 | 2 |
| | | | | 666 | 148.90 | 149.85 | 0.95 | 0.56 | 123.4 |
| | | | | 667 | 149.85 | 150.40 | 0.55 | 0.015 | 2.3 |
| | | | | 668 | 150.40 | 151.15 | 0.75 | 0.16 | 23.9 |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> | | | | |
|------------------|-----------|----------------------------------|--|-------------|------------|------------|-----------|-----------|-----------|------|------|--|-------|
| <i>From</i> | <i>To</i> | | | <i>Len.</i> | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> | | | | |
| | | | | | 669 | 151.15 | 151.50 | 0.35 | 0.015 | 2.1 | 0.01 | | |
| | | | | | 671 | 151.50 | 152.00 | 0.50 | 0.16 | 3.95 | 0.14 | | 0.005 |
| | | | | | 672 | 152.00 | 153.10 | 1.10 | 0.015 | 2.2 | 0.01 | | |
| | | | | | 673 | 153.10 | 153.60 | 0.50 | 0.015 | 2.2 | 0.03 | | |
| | | | | | 674 | 153.60 | 154.45 | 0.85 | | | | | |
| | | | | | 675 | 154.45 | 155.10 | 0.65 | | | | | |
| 131.20 | - 131.70 | QCV Quartz Carbonate Vein | white qtz-carb vein with coarse clots tour and chl; minor sulphides; sharp contact with previous @ 30deg to CA | | | | | | | | | | |
| 131.70 | - 132.95 | QV Quartz Vein | white qtz with fine tour-chl fractures @ 30-35deg to CA; minor sulphides | | | | | | | | | | |
| 132.95 | - 134.00 | QRZ Quartz Rich Zone | 80% white and grey qtz, 20% int alt 9i as diffuse inclusions; sharp contact with previous; 2-4% py mostly in 9i, minor py, a few vf specks cpy,gal in qtz | | | | | | | | | | |
| 134.00 | - 134.85 | QRZ Quartz Rich Zone | 80% int alt 9i, 20% white qtz as stringers and angular clasts; 10-12% py, a few small blebs gal | | | | | | | | | | |
| 134.85 | - 135.70 | QRZ Quartz Rich Zone | 80% white qtz, 20% int alt 9i; 1% py, a few small blebs gal in qtz | | | | | | | | | | |
| 135.70 | - 136.65 | QRZ Quartz Rich Zone | 30% qtz stringers and clasts with <1% diss py, a few diffuse blebs gal; 70% 9i with 10-12% vf py | | | | | | | | | | |
| 136.65 | - 137.50 | 9i Trondhemite, altered | 9i with a few qtz stringers, 6-8% vf diss py | | | | | | | | | | |
| 137.50 | - 138.95 | 9i Trondhemite, altered | 10% qtz as stringers and angular clasts, minor sulphides; 90% 9i with 4-6% py | | | | | | | | | | |
| 138.95 | - 139.50 | QV Quartz Vein | white qtz, 1-2% diss py, 1% gal, several specks acicular arg?, a few specks cpy | | | | | | | | | | |
| 139.50 | - 139.85 | QRZ Quartz Rich Zone | 50% qtz with 1% py, several vf specks gal?, 50% 9i with 6-8% py | | | | | | | | | | |
| 139.85 | - 140.95 | QRZ Quartz Rich Zone | 80% qtz with 1% diss py, 1% vf gal? in diffuse blebs and seams, numerous xls acicular arg?; section appears to be one or more narrow qtz veins @ 20deg to CA | | | | | | | | | | |
| 140.95 | - 141.25 | 9i Trondhemite, altered | 10% qtz as angular clasts and 1cm stringer @20deg to CA with vf gal? seam along stringer wall; | | | | | | | | | | |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> |
|------------------|-----------|---|--|-------------|------------|------------|-----------|-----------|-----------|
| <i>From</i> | <i>To</i> | | | <i>Len.</i> | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| 141.25 | - 141.85 | QRZ Quartz Rich Zone 50% qtz with <1% combined py,gal as few coarse blebs, 6-8% diss py in 9i | | | | | | | |
| 141.85 | - 142.35 | QRMZ Quartz Rich Mineralized Zone 70% qtz with 8-10% py as coarse xl aggregates, 1-2% vf gal/sph? as diffuse haloes to py aggregates, a few small blebs gal | | | | | | | |
| 142.35 | - 144.10 | 9i Trondhjemite, altered int alt 9i, silicified, vfg with a few remnant qtz phenos, 4-6% diss py | | | | | | | |
| 144.10 | - 145.75 | 9i Trondhjemite, altered 9i, int alt, med grained, minor qtz, 2-4% diss py | | | | | | | |
| 145.75 | - 146.30 | MQV Mineralized Quartz Vein white, massive qv with sharp contacts @35-40deg to CA; 1-2cm band of semi massive gal and sph at lower contact and a few small blebs scattered in rest of section; 1-2% overall | | | | | | | |
| 146.30 | - 147.65 | 9i Trondhjemite, altered int alt 9i, 2-4% diss py, minor qtz | | | | | | | |
| 147.65 | - 148.35 | QV Quartz Vein white qtz, sharp contacts @ 35-40deg to CA, a few small blebs py, gal | | | | | | | |
| 148.35 | - 148.90 | 9i Trondhjemite, altered int alt 9i, 6-8% diss py | | | | | | | |
| 148.90 | - 149.85 | QRZ Quartz Rich Zone 70% qtz, similar to 141.85-142.35; 6-8% py in xl aggregates with diffuse halos of vf gal? and a few small blebs gal; appears to be one or narrow stringers arcing near // to CA | | | | | | | |
| 149.85 | - 150.40 | 9i Trondhjemite, altered int alt 9i | | | | | | | |
| 150.40 | - 151.15 | QRZ Quartz Rich Zone v similar to 148.9-149.8; qtz stringers near // to CA; may be the same veins | | | | | | | |
| 151.15 | - 151.50 | 9i Trondhjemite, altered altered trondhjemite | | | | | | | |
| 151.50 | - 152.00 | QRZ Quartz Rich Zone 30% qtz as 1-2cm stringers @ 50-60deg to CA with a few blebs sph? (could be an fe-carb), gal, a few fine needles arg? | | | | | | | |
| 152.00 | - 153.10 | 9i Trondhjemite, altered altered trondhjemite | | | | | | | |
| 153.10 | - 153.60 | QCT Carbonate Tourmaline Quartz Vein 153.1-153.6m 15cm white qtz-carb-tour and a few stringers @ 50deg to CA, minor sulphides | | | | | | | |

| <i>Lithology</i> | | | | | <i>Au</i> | <i>Ag</i> | <i>Zn</i> | <i>Cu</i> | <i>Pb</i> | | | |
|------------------|-----------|------------|---|-----------------|-------------|-----------|-------------|------------|------------|----------|----------|----------|
| <i>From</i> | <i>To</i> | | | <i>Sample #</i> | <i>From</i> | <i>To</i> | <i>Len.</i> | <i>ppm</i> | <i>ppm</i> | <i>%</i> | <i>%</i> | <i>%</i> |
| 153.60 | - 154.50 | 9i | Trondhjemite, altered altered trondhjemite | | | | | | | | | |
| 154.50 | - 155.10 | QCT | Carbonate Tourmaline Quartz Vein qtz-car-tour vein @ 25deg to CA, minor sulphide | | | | | | | | | |
| 155.10 | - 156.20 | 9i | <u>Trondhjemite, altered</u> fine to med grained, int alt, slight pink tint, 4-6% diss py, | 676 | 155.10 | 156.00 | 0.90 | | | | | |
| 156.20 | - 156.50 | FZ | <u>Fault Zone</u> 5cm rusty gouge, slickensides, remainder has strong fol'n @ 35-40 deg to CA | | | | | | | | | |
| 156.50 | - 177.00 | 9i | <u>Trondhjemite, altered</u> | 677 | 161.30 | 162.80 | 1.50 | | | | | |
| 156.50 | - 161.30 | 9i | Trondhjemite, altered int alt, pale green-grey, fine to med grained, strong fol'n @ 35-40deg to about 159m; 2-4% vf diss py | | | | | | | | | |
| 161.30 | - 162.80 | 9i | Trondhjemite, altered as above with several qtz stringers <1cm with coarse py xls | | | | | | | | | |
| 162.80 | - 164.70 | 9i | Trondhjemite, altered med grained, massive, strong alt | | | | | | | | | |
| 164.70 | - 165.50 | 9i | Trondhjemite, altered blocky and broken with qtz-carb fracture fillings, weak fracture zone? | | | | | | | | | |
| 165.50 | - 170.90 | 9i | Trondhjemite, altered grey, med grained, massive, mod alt, minor sulphides | | | | | | | | | |
| 170.90 | - 174.00 | 9i | Trondhjemite, altered mod to strong alt, qtz stringers and int alt band <1cm near //to CA; 8-10% py xls and a few coarse blebs | | | | | | | | | |
| 174.00 | - 177.00 | 9i | Trondhjemite, altered mod alt 9i | | | | | | | | | |