

MECHANICAL TRENCHING AND PROSPECTING REPORT

ON THE


NARROWS GOLD PROPERTY
(WHITE LAKE NARROWS)

THUNDER BAY MINING DIVISION

DISTRICT OF THUNDER BAY

NTS 42C12

NTS 42C13



Thunder Bay Ontario.

March 28, 2010

Douglas. N .Kakeeway

Prospector

2 • 44760

TABLE OF CONTENTS

| | PAGE |
|---|------|
| 1. SUMMERY | 1 |
| 2. INTRODUCTION | 1 |
| 3. PROPERTY LOCATION AND ACCESS | 1 |
| 4. TOPOGRAPHY AND VEGETATION | 1 |
| 5. REGIONAL GEOLOGY | 2 |
| 6. LOCAL GEOLOGY ON CLAIM # 3005078 | 2 |
| 7. LOCAL GEOLOGY ON CLAIM # 3005075 AND # 3005077 | 3 |
| a) Lithologies | 3 |
| b) Mafic Metavolcanic | 4 |
| c) Course Grained Amphibolite | 4 |
| d) Migmatite | 4 |
| e) Metasediment | 5 |
| f) Metagabbro | 5 |
| g) Granodiorite | 5 |
| h) Granite | 5 |
| i) Felsic Intrusive | 6 |
| j) Lamprophyre | 6 |
| k) Diabase | 6 |
| 8. CONCLUSIONS | 6 |
| 9. RECOMMEDATIONS | 6 |

APPENDICES

APPENDIX "A" ASSAY CERTIFICATES

LIST OF FIGURES

FIGURE 1 KEY LOCATION MAP

FIGURE 2 LOCAL GEOLOGY MAP

FIGURE 3 REGIONAL GEOLOGY MAP

FIGURE 4 CLAIM MAP

LIST OF TABLES

TABLE 1 ROCK SAMPLE LOCATIONS

TABLE 2 ROCK SAMPLE DESCRIPTIONS

TABLE 3 DAILY PROSPECTING LOG

TABLE 4 BACKHOE DAILY LOG

LIST OF MAPS

MAP 1 NORTH EAST TRENCHES AND AREA GEOLOGY

MAP 2 NORTH STRIP ZONE AND AREA GEOLOGY

MAP 3 STRIP ZONE "A" AND AREA GEOLOGY

MAP 4 STRIP ZONE "B" AND AREA GEOLOGY

MAP 5 STRIP ZONE "C" AND AREA GEOLOGY

MAP 6 TRENCH AREA "D" AND AREA GEOLOGY

MAP 7 OVERVIEW OF TRAVERSES, TRENCHES, AND GEOLOGY

MAP 8 NORTH STRIPPED ZONE AND TRENCHES

MAP 9 STRIP ZONE "A"

MAP 10 STRIP ZONE "B" AND TRENCHES

MAP 11 STRIP ZONE "C" AND TRENCHES

MAP 12 TRENCH AREA "D"

MAP 13 TRENCH AREA "D" AND TRAVERSES





Figure 1: Location Map



MAP
 OGS M2614
 LEGEND ATTACHED

UTM/NAD 27
 Figure 2

SCALE
 2.8cm = 1km

LEGEND

PRECAMBRIAN

NEOPROTEROZOIC

- 10 Port Colville-Alaska Complex^{ab}
 - 10a Gabbro
 - 10b Pyroxene syenite
 - 10c Amphibole syenite
 - 10d Quartz syenite
 - 10e Heterogeneous syenite
 - 10f Neoproterozoic (?) amygdaloidal mafic flows (pendents)

INTRUSIVE CONTACT

PALEOPROTEROZOIC TO MESOPROTEROZOIC

- Mafic Intrusive Rocks^d
- Diorite dikes ± plagioclase phenocrysts

INTRUSIVE CONTACT

NEOARCHAIC

- 5 Felsic to Intermediate Intrusive Rocks^{ab}

Capures based on known and inferred ages (see note d, below)

- Plutons 2673-2677 Ma
- Plutons and Stocks 2688-2694 Ma
- Pluton 2697 Ma
- Bethulie - Midway Terrane 2720-2686 Ma

- 8a Leucocratic biotite tonalite to biotite granodiorite^d
- 8b Biotite tonalite^d
- 8c Biotite-hornblende tonalite
- 8d Hornblende-biotite tonalite^d
- 8e Plagioclase-phryic biotite tonalite
- 8f Plagioclase-phryic biotite-hornblende to hornblende-biotite tonalite
- 8g Plagioclase-phryic hornblende tonalite
- 8h Plagioclase-phryic biotite-hornblende tonalite gneiss
- 8i Biotite granodiorite
- 8j Biotite-hornblende granodiorite^d
- 8k Hornblende-biotite granodiorite^d
- 8l Plagioclase-phryic to eubphyric biotite granodiorite^d
- 8m Plagioclase-phryic biotite-hornblende granodiorite gneiss^d
- 8n Plagioclase-eubphyric biotite-hornblende to hornblende-biotite granodiorite^d
- 8p Vesically microcline-megacrystic hornblende-biotite granodiorite^d
- 8q Biotite-hornblende quartz monzonite
- 8r Hornblende-biotite quartz monzonite^d
- 8s Hornblende monzonite to hornblende quartz monzonite
- 8t Equigranular to plagioclase-subphyric hornblende diorite to quartz monzonite to granodiorite
- 8u Microcline-megacrystic hornblende-biotite diorite to quartz monzonite to granodiorite^d
- 8v Mainly foliated to gneissic tonalite to granodiorite; local massive to foliated phases; diverse minor phases^d
- 8w Plagioclase ± quartz porphyry^d
- 8x Aplita, pegmatite
- 8y Unsubdivided massive to weakly foliated granitoid rocks

INTRUSIVE CONTACT

- Metamorphosed Ultramafic Intrusive Rocks^d
- 7a Peridotite
- 7b Pyroxenite
- 7c Saponarite
- 7d Hornblende

- 6 Metamorphosed Mafic Intrusive Rocks^d
- 6a Dabing
- 6b Diorite^d
- 6c Unsubdivided, massive to gneissic, mafic to intermediate, intrusive and/or volcanic rocks
- 6d Schistose to gneissic rocks

INTRUSIVE CONTACT

- 5 Metasedimentary Rocks^{ab}
- 5a Mudstone (siltstone, claystone), minor wacke
- 5b Wacke, thin wacke, local minor conglomerate^d
- 5c Arenite, thin arenite, local minor conglomerate
- 5d Conglomerate ± thin wacke ± thin arenite^d
- 5e Oolite (magnesian) iron formation
- 5f Schistose rock
- 5g Gneissic rock
- 5h Migmatitic rock

- 4 Felsic Metavolcanic Rocks^d
- 4a Massive flows (rare flow layering), related subvolcanic intrusions; commonly plagioclase-quartz-phryic^d
- 4b Plagioclase-quartz-phryic tuff, lapilli tuff and reworked deposits^d
- 4c Plagioclase ± quartz-phryic tuff breccia, pyroclastic breccia and reworked deposits
- 4d Schistose rock

- 3 Intermediate Metavolcanic Rocks^d
- 3a Massive and pillowed flows; commonly plagioclase-phryic; locally amygdaloidal
- 3b Plagioclase-quartz-phryic tuff, lapilli tuff and reworked deposits^d
- 3c Plagioclase-quartz-phryic tuff breccia, pyroclastic breccia and reworked deposits
- 3d Schistose rock
- 3e Migmatitic rock

- 2 Mafic Metavolcanic Rocks^d
- 2a Massive to pillowed flows
- 2b Massive to pillowed flows with amygdalite and/or vesicles
- 2c Massive to pillowed flows with plagioclase phenocrysts
- 2d Tuff, lapilli tuff
- 2e Amphibolite
- 2f Schistose rock
- 2g Gneissic rock
- 2h Migmatitic rock
- 2i Pyroxene-spinel-textured flows

- Ultramafic Metavolcanic Rocks^d
- 1a Massive to pillowed peridotite flow
- 1b Olivine-spinel-textured flows
- 1c Polystratified flows
- 1d Schistose rock

^a Rocks in these units are subdivided lithologically and the order does not imply age relationships within the units.

^b Internal "contacts" within these units do not necessarily represent clastic igneous phases (units 10, 6) or sedimentary packages (unit 5).

^c The letter "G" preceding lithologic code 6 indicates a dike is inferred from aeromagnetic data.

^d See table of "U-Pb Zircon Geochronologic Ages for the Heron Greenstone Belt" (in text).

^e Some conglomerate-predominant sedimentary deposits may reflect Tivishaming-type sedimentation/tectonism.




Figure 3: Regional Geology (Muir et al, 1995)



figure 4

Claim Map

3.5cm is 1.6 km

1) Summary

Prospecting, rock sampling and power stripping and trenching were conducted on the Narrows Gold Property on the east side of the White Lake narrows from May 2008 to November 2010. A modified Clark skidder 664C quick coupled to an older version international backhoe was used in the stripping and trenching. A former cut grid that was established in 2002 and it was used to guide some of the prospecting using a geotul hammers. A total of 80 grab samples where taken.

2) Introduction

The Narrows Gold Property consisting of 3 contiguous claim blocks comprising of 44 units and are located 31 km west of the town of White River Ontario and 56 km east of Marathon Ontario. The claims lie within and on the east Shore of White Lake. The property is covered by N.T.S. map sheet 42C/12 and 42C/13 and by claim maps G-0623, G-0622.

3) Property Location and Access

The property is located 31 km west of the town of White River and 22.5 km east of Hemlo gold deposit in the Thunder Bay Mining Division. It includes portions of White Lake and surrounding land just north of Highway 17. A Public boat launch on the west side of white lake is available close to the Marathon boat club to access the property.

4) Topography and Vegetation

The Property consists of rolling hills rising from White Lake and generally ending in cliffs of 2-30m. The maximum relief is approximately 50m. Swampy and flat terrain is also prevalent. A few smaller ponds are contained within the property.

The higher areas are covered by large popular, birch and spruce, and smaller fir and spruce trees with local undergrowth of shrub maple and tag alder. Low lying areas contain cedar, tag alders and black spruce. Locally spruce budworm has devastated the trees and areas of deadfall are common.

In approximate year 2000 a forest fire has burnt about 1/2 of claim #3005078 in the east side of the claim and about 1/3 of claim #3005077 in the north east was also burnt.

5) Regional Geology

The Narrows property lies within the Heron Bay-Hemlo portion of the Schreiber-Hemlo greenstone belt in the Wawa subprovince of the Superior Province (Muir, 1983). This greenstone belt is composed of Archean metavolcanic and metasedimentary rocks surrounded by the regional granitic rocks and runs approximately east-west (Figure 3). The property region is underlain by mafic metavolcanics and metasediments with mafic, intermediate and felsic intrusives. Late intrusives consist of lamprophyre and diabase dikes. The metamorphic grade ranges from greenschist to amphibolite facies. A dominant north-northwest schistosity is found in this area.

The bedrock in the area is generally covered by a thin layer of surficial deposits consisting of humus and soil with a thin layer of glacial drift (Geddes, R.S. and Kristjansson, F.J., 1986). Locally the tills may be thicker. In some areas thick sections of glaciolacustrine deposits occur.

6) Local Geology on Claim #3005078

(Descriptions used from Graphite group property report year 1996)

The geology of the Claim is described in detail by Gallo (1990a, 1990b, 1991, 1992a, 1992b, and 1992c) and by McKay (1994). In summary, the claim is underlain primarily by a northwesterly-trending sequence of intercalated mafic metavolcanic and clastic metasedimentary rocks that have been intruded locally by foliation-parallel and foliation-crosscutting sills and dikes of both mafic and felsic compositions, and by narrow veins of quartz

The mafic metavolcanic rocks are typically locally rusty-weathering, dark green to black, generally moderately foliated, medium-grained, non-magnetic, and non-calcareous amphibolitic schists. The foliation in these rocks varies in strike from 145 to 165 degrees, and in dip from 75 to 85 degrees west. Trace amounts of fine-grained disseminated pyrite were present in most of the rocks examined.

The clastic metasedimentary rocks occur as relatively thin beds within the amphibolitic mafic metavolcanic sequence. The metasedimentary rocks were observed in two locations on the property. They outcrop intermittently along the ridge that bisects the southern part of the property, and are exposed on a small point of land jutting out into White Lake near the western

edge of the property. The metasedimentary rocks are typically locally sheared, medium-grained, biotitic, locally graphitic, non-calcareous, quartzo-feldspathic schists. The foliation in these rocks varies in strike and dip within the ranges quoted above for the mafic metavolcanic rocks. The biotitic-rich metasedimentary schists may represent metamorphosed arkose.

The mafic intrusive rocks examined during the present survey comprise of 2 large, norwesterly- and northerly-trending diabase dikes. These rocks are typically brownish-weathering, and dark green, massive, medium- to coarse-grained, locally moderately magnetic and contain less than 1% pyrite as fine- to medium-grained anhedral grains.

The felsic intrusive rocks occur primarily along the western and eastern margins of the property and envelope and intrude locally the supracrustal rocks. These intrusive rocks are typically buff-weathering, pinkish, weakly foliated, coarse-grained rocks of granitic to granodioritic composition. They were observed to contain numerous rounded xenoliths of amphibolite in several locations proximal to the western contact with the mafic metavolcanic rocks.

Quartz-feldspar and feldspar porphyritic, foliation-parallel sills occur widely distributed within the metavolcanic and metasedimentary rocks. Narrow (less than 1 mm to 50 cm wide), foliation-parallel and foliation-crosscutting quartz veins also occur locally. These quartz veins are glassy-white and generally barren-looking.

7) Local Geology on Claim #3005075 and #3005077

(Descriptions used from M.Stalker White Lake project year 2000)

a) Lithologies

The White Lake property is underlain by a sequence of metavolcanic and metasedimentary rocks which have been inundated by mafic, intermediate, and felsic intrusive. The close proximity of large batholiths has led to metamorphic aureoles and magma mixing causing heterogeneous outcrops. Lithologies change or grade from outcrop to outcrop or within the same outcrop and contacts may be crosscutting or gradational making it hard to outline individual units. Rocks have been metamorphosed to the amphibolites facies.

b) Mafic Metavolcanic

Much of the property is underlain by mafic flows which grade into coarser grained amphibolites. The mafic volcanic are usually comprised of amphiboles and chlorite and commonly could be termed an amphibolites. They can be biotite rich especially where shearing occurs, rarely, they are muscovite rich. Poorly developed pillows were observed at only one location but outcrops that have an indication of pillows or ropy lava but no definite selvages are more common. Garnets and a beaded mineral, probably sillimanite, are common especially in those outcrops suggesting pillows. All of the mafics exhibit a foliation but it can be strong to sheared over small zones. Locally the mafic flows are altered to light green to tan in bands at an angle to foliation. Many of the mafic flows have traces of medium grained cubes of pyrite but rarely outcrops are rusty and may contain up to 5% pyrite locally.

c) Coarse Grained Amphibolites

This unit covers a wide variety of rocks on the property. It is made of coarser grained amphibolites with up to 30% pink or white felsic matrix. This unit is probably the metamorphic equivalent of the mafic volcanic flows which have been affected by the intrusion of the intermediate to felsic intrusive. Grain size can be from 1mm to 5cm. It grades between the mafic volcanic and the granodirite or may be crosscutting these units. This unit is commonly without foliation but may be foliated or gneissic locally. Trace amount of pyrite and rare molybdenum may be found in the amphibolites, especially in the felsic matrix.

d) Migmatite

This unit has a light grey intermediate to felsic matrix with pods of mafic material and pods of granitic material which resemble slightly stretched clasts. These pods grade in composition with a number of different varieties. These pods look like they are replaced primary clasts, possibly originally a pyroclastic rock, but the unit may also be a hybrid of two different magmas. This unit occurs in seven different locations comprising several outcrops in the central part of the property. The largest observed width of the unit was >10m but it also appears in bands <1m. This lithology is often biotite rich and scattered pyrite cubes are common.

e) Metasediments

There are two main types of Lithologies on the property which appear to be meta-sediments. Biotite rich schist to gneiss with quartz and feldspar grains which grades to a more arkosic rock. A few outcrops are biotite schist that looks like a lamprophyre dike and it is difficult to tell between the two. Possibly a granitized sediment which is fine grained, laminated, and may be strongly sheared. It is commonly sericitized and locally muscovite rich with rare green mica grains. The unit gives the appearance of a mylonite and folding is commonly evident. Quartz eyes are found in some of these outcrops and it is possible this unit is altered porphyry with laminations due to alteration. Minor pyrite is found locally in both of these units and trace amounts of molybdenite is found in the granitized sediments.

f) Metagabbro

The gabbro is very similar to and is possibly the same unit as amphibolite with <5% felsic matrix. It is very coarse grained with grains up to 10 cm. It appears to be a true gabbro and not just a very coarse grained equivalent of the amphibolite because of its stronger magnetic signature, higher Ni content, and rarely observed cross cutting contacts with the amphibolite. Locally the gabbros' magnetic field is strong enough to disturb a compass. At some locations the gabbro and amphibolite contact does appear gradational. However, the two units do overlap and may easily be mistaken for one another.

g) Granodiorite

Granodiorite is a very prominent rock type on the grid. It grades between granite and amphibolite. It is mainly medium to coarse grained but may be very coarse grained. It is composed of amphibole and chlorite and less commonly biotite with pink and white feldspar grains and white quartz grains. The Granodiorite often grades in composition but slightly different compositions may also have sharp dike like contacts indicating a number of different intrusions of the Granodiorite magma. It is common on the property to see a Granodiorite outcrop with pods of Granodiorite in a more mafic matrix. On the north shore of the cut grid area are good examples of this with rounded pods of Granodiorite in a Matrix formed of amphibole.

h) Granite

Granite is not as prevalent on the property and probably is the most felsic end member of the Granodiorite batholith. It is mostly composed of feldspar, quartz, amphibole and biotite. It is commonly gneissic.

i) Felsic Intrusive

Felsic intrusives are prevalent throughout the property. Pegmatite's are common and consist of coarse grains of quartz, feldspar, and biotite.

Granite intrusives are also commonly found. Less common are aplitic intrusives. All of these intrusives are generally less than 1m in width.

j) Lamprophyre

Two lamprophyre dikes were observed on the property. These are fine to medium grained and biotite rich and are similar to the biotite schist sediments. They may be metasediment but appear to have intrusive contacts although these contacts are parallel to foliation.

k) Diabase

Three outcrops of diabase were observed but common diabase rubble indicates that it is more prevalent. Both fine grained and coarse grained diabase occur. The coarser diabase is magnetic. The diabase contains trace amounts of pyrite.

8) Conclusions

On claim #3005077 The stripping was successful in extending the former Carroll gold showing 35 meters north and sampling of the vein has given gold values up 8.4 Gr/ton. The prospecting program located 2 floats on claim #3005075 that assayed 4482 ppb and 15045 ppb gold. On claim #3005075 anomalous bedrock samples up to 259 ppb in trench "C"

9) Recommendations

Recut and repicket old grid lines to coordinate project. Continue stripping and trenching the Carroll vein north and south along strike. Continue prospecting to find where high grade floats came from. Continue stripping and trenching "trench C" to get higher gold values.

TABLE 1

WHITE LAKE NARROWS ROCK SAMPLE LOCATIONS 2008 AND 2009

| TAG NUMBER | NAD 83 ZONE | EASTING | NORTHING |
|-----------------------|------------------------|----------------|-----------------|
| 390620 | 16 U | 602346 | 5400919 |
| 390624 | 16 U | 601165 | 5402082 |
| 390625 | 16 U | 601164 | 5402082 |
| 390626 | 16 U | 601162 | 5402081 |
| 390627 | 16 U | 601184 | 5402065 |
| 390628 | 16 U | 601185 | 5402060 |
| 390629 | 16 U | 601186 | 5402056 |
| 390630 | 16 U | 601187 | 5402052 |
| 390631 | 16 U | 601188 | 5402047 |
| 390632 | 16 U | 601655 | 5401324 |
| 390633 | 16 U | 601654 | 5401324 |
| 390634 | 16 U | 601650 | 5401324 |
| 390642 | 16 U | 601345 | 5399805 |
| 390643 | 16 U | 601347 | 5399784 |
| 390646 | 16 U | 601322 | 5399840 |
| 390647 | 16 U | 601323 | 5399840 |
| 390648 | 16 U | 601324 | 5399840 |
| 390649 | 16 U | 601325 | 5399840 |
| 390650 | 16 U | 601326 | 5399840 |
| 716001 | 16 U | 601330 | 5399838 |
| 716002 | 16 U | 601331 | 5399838 |
| 716003 | 16 U | 601332 | 5399837 |
| 716004 | 16 U | 601333 | 5399836 |
| 716005 | 16 U | 601334 | 5399836 |
| 716006 | 16 U | 601335 | 5399836 |
| 716007 | 16 U | 601336 | 5399835 |
| 716008 | 16 U | 601337 | 5399834 |
| 716009 | 16 U | 601339 | 5399832 |
| 716010 | 16 U | 601340 | 5399831 |
| 716011 | 16 U | 601301 | 5398686 |
| 716012 | 16 U | 601327 | 5399840 |
| 716013 | 16 U | 601320 | 5399846 |
| 716014 | 16 U | 601319 | 5399857 |
| 716015 | 16 U | 601318 | 5399861 |
| 716016 | 16 U | 601317 | 5399862 |
| 716017 | 16 U | 601316 | 5399863 |
| 716018 | 16 U | 601315 | 5399864 |
| 716019 | 16 U | 601314 | 5399865 |
| 716020 | 16 U | 601313 | 5399866 |
| 716021 | 16 U | 601332 | 5399859 |

| TAG NUMBER | NAD 83 ZONE | EASTING | NORTHING |
|-----------------------|------------------------|----------------|-----------------|
| 716022 | 16 U | 601325 | 5399864 |
| 716023 | 16 U | 601326 | 5399865 |
| 716024 | 16 U | 601313 | 5399797 |
| 716025 | 16 U | 601302 | 5398683 |
| 716026 | 16 U | 601321 | 5399832 |
| 716027 | 16 U | 601333 | 5399839 |
| 716028 | 16 U | 601311 | 5399869 |
| 716029 | 16 U | 601312 | 5399872 |
| 716030 | 16 U | 601105 | 5400249 |
| 716031 | 16 U | 601107 | 5400229 |
| 716032 | 16 U | 601276 | 5398547 |
| 716033 | 16 U | 601328 | 5399836 |
| 716038 | 16 U | 601707 | 5401683 |
| 716039 | 16 U | 601726 | 5401639 |
| 716040 | 16 U | 601347 | 5401772 |
| 716041 | 16 U | 601347 | 5401773 |
| 716042 | 16 U | 601367 | 5401753 |
| 716043 | 16 U | 601170 | 5402183 |
| 716044 | 16 U | 601086 | 5402183 |
| 716045 | 16 U | 600901 | 5402594 |
| 759601 | 16 U | 601332 | 5398558 |
| 759602 | 16 U | 601323 | 5398521 |
| 759606 | 16 U | 601651 | 5398583 |
| 759609 | 16 U | 601448 | 5398779 |
| 759610 | 16 U | 601461 | 5398802 |
| 759611 | 16 U | 601474 | 5398785 |
| 759612 | 16 U | 601440 | 5398810 |
| 759613 | 16 U | 601440 | 5398811 |
| 759614 | 16 U | 601440 | 5398812 |
| 759615 | 16 U | 601475 | 5398786 |
| 759616 | 16 U | 601518 | 5398749 |
| 759617 | 16 U | 601379 | 5399038 |
| 759618 | 16 U | 601379 | 5399039 |
| 759619 | 16 U | 601379 | 5399040 |
| 759620 | 16 U | 601379 | 5399042 |
| 759621 | 16 U | 601379 | 5399042 |
| 759622 | 16 U | 601331 | 5399054 |
| 759623 | 16 U | 601333 | 5399053 |
| 759624 | 16 U | 601334 | 5399053 |
| 759651 | 16 U | 601906 | 5399427 |

TABLE 2

| SAMPLE NUMBER | PARTS PER BILLION GOLD | ROCK DESCRIPTION |
|---------------|------------------------|--|
| 390620 | 10 | Black foliated, medium grained amphibolite, 2" transparent qtz veins |
| 390624 | 16 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390625 | 13 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390626 | 12 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390627 | 164 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite and in Contact with feldspar porphyry? |
| 390628 | 19 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390629 | 20 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390630 | 15 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390631 | 12 | Very fine grained, laminated on cm scale, medium gray color, sediment? <1% pyrite |
| 390632 | 5 | Basalt? ,>4% pyrite |
| 390633 | 11 | Basalt? ,>4% pyrite |
| 390634 | 127 | Basalt? ,>4% pyrite |
| 390642 | 7 | Sediment, rusty, garnets |
| 390643 | 5 | Sediment, rusty, garnets |
| 390646 | 5 | Sediment, fine grain, black, no pyrite |
| 390647 | 5 | Sediment, fine grain, black, no pyrite, garnets |
| 390648 | 5 | Sediment, fine grain, black, no pyrite, |
| 390649 | 111 | Sediment, fine grain, black, no pyrite |
| 390650 | 70 | Sediment, fine grain, black, no pyrite |
| 716001 | 11 | Sediment, fine grain, black, no pyrite |
| 716002 | 348 | Sediment, fine grain, black, 2 inch transparent quartz, no pyrite, contact with white coarse grained pegmatite. |
| 716003 | 104 | Sediment, fine grain, black, no pyrite |
| 716004 | 8 | Sediment, fine grain, black, no pyrite |
| 716005 | 5 | Sediment, fine grain, black, no pyrite |
| 716006 | 20 | Sediment, fine grain, black, no pyrite |
| 716007 | 5 | Sediment, fine grain, black, no pyrite |
| 716008 | 9 | Sediment, fine grain, black, no pyrite |
| 716009 | 15 | Sediment, fine grain, black, no pyrite |
| 716010 | 94 | Sediment, fine grain, black, no pyrite |
| 716011 | 10 | Rusty angular sericitized schist float >1% pyrite |
| 716012 | 33 | Sediment, fine grain, black, no pyrite |
| 716013 | 11 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716014 | 49 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716015 | 8420 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716016 | 200 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716017 | 249 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716018 | 312 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716019 | 4402 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716020 | 18 | Quartz in bedrock, 12" to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716021 | 26 | Sediment, fine grain, black, no pyrite |
| 716022 | 5 | Sediment, fine grain, black, no pyrite |
| 716023 | 5 | Sediment, fine grain, black, no pyrite |
| 716024 | 7 | Sediment, fine grain, black, no pyrite |


| SAMPLE NUMBER | PARTS PER BILLION GOLD | ROCK DESCRIPTION |
|---------------|------------------------|--|
| 716025 | 10 | Round rusty basalt float |
| 716026 | 98 | Quartz float close to quartz in bed rock , transparent, rusty,<1% sulfide |
| 716027 | 15293 | Quartz float close to quartz in bed rock , transparent, rusty,<1% sulfide |
| 716028 | 222 | Quartz in bedrock, 12"to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment with garnets and epidote |
| 716029 | 279 | Quartz in bedrock, 12"to 18" wide, rusty, <1% sulfides. Host rock is sheared Sediment? with garnets and epidote |
| 716030 | 5 | Gabbro, medium grained, rusty,<1% pyrite |
| 716031 | 5 | Gabbro, medium grained, rusty,<1% pyrite |
| 716032 | 4482 | Black rusty amphibolite host, fine grained, rusty quartz cross cutting with Calcite, no visible sulfides.(float) |
| 716033 | 47 | Rusty Basalt float, >3% sulfides |
| 716038 | 14 | Amphibolite,black,medium grain and foliated, Qtz |
| 716039 | 5 | Sediment, medium grain foliated <1% pyrite |
| 716040 | 56 | Rusty Amphibolite, medium grained, <1% pyrite, qtz. |
| 716041 | 5 | Rusty Amphibolite, medium grained, <1% pyrite, qtz. |
| 716042 | 9 | Rusty qtz |
| 716043 | 5 | Rusty Fine grain sediment, foliated<1% pyrite |
| 716044 | 4 | Rusty Fine grain sediment, foliated,<1% pyrite |
| 716045 | 5 | Rusty Fine grain sediment, foliated<1% pyrite |
| 759601 | 15045 | Fine grained foliated sediment float, 1% pyrite, qtz. |
| 759602 | 10 | Sediment medium grained, foliated with biotite |
| 759606 | 6 | Fine grained foliated sediment,qtz,calcite,<1% pyrite |
| 759609 | 24 | Rusty quartz and fizzes with acid (calcite?) |
| 759610 | 8 | Creamy color fine grained sediment and qtz, <1% pyrite |
| 759611 | 18 | Very fine grained foliated sediment, dark gray color, <1% pyrite, qtz. |
| 759612 | 10 | Very fine grained foliated sediment, dark gray color, 1% pyrite, qtz. |
| 759613 | 13 | Very fine grained foliated sediment, dark gray color, 1% pyrite, qtz. |
| 759614 | 13 | Very fine grained foliated sediment, dark gray color, <1% pyrite, qtz. |
| 759615 | 13 | Very fine grained foliated sediment, dark gray color, <1% pyrite, qtz. |
| 759616 | 30 | Very fine grained foliated sediment, dark gray color, <1% pyrite, qtz. |
| 759617 | 102 | 3 to 4 parallel rusty quartz veins, average width of the quartz veins are 8 inches and the host rock is sediment? |
| 759618 | 37 | 3 to 4 parallel rusty quartz veins, average width of the quartz veins are 8 inches and the host rock is dark sediment? |
| 759619 | 36 | 3 to 4 parallel rusty quartz veins, average width of the quartz veins are 8 inches and the host rock is dark sediment? |
| 759620 | 259 | 3 to 4 parallel rusty quartz veins, average width of the quartz veins are 8 inches and the host rock is dark sediment? |
| 759621 | 100 | 3 to 4 parallel rusty quartz veins, average width of the quartz veins are 8 inches and the host rock is dark sediment? |
| 759622 | 5 | 2" quartz vein in basalt? |
| 759623 | 5 | 2" quartz vein in basalt? |
| 759624 | 5 | 2" quartz vein in basalt? |
| 759621 | 5 | Sericitized schist |

TABLE 3

DAILY PROSPECTING LOG

Two men prospecting on each traverse or job

| Date | Claim No. | Description of Work Done | traverse area location UTM/NAD 83/ Zone 16U | Hours Worked per day |
|--------------------------------|-----------|--------------------------|--|-------------------------|
| MAY19/09 | 3005075 | prospecting/sampling | 601225/5398715 | 4 |
| JUNE11/09 | 3005075 | prospecting/sampling | 601547/5398336 | 8 |
| JULY12/09 | 3005075 | prospecting/sampling | 601619/5398263 | 8 |
| JULY16/09 | 3005075 | prospecting/sampling | 601629/5398619 | 8 |
| AUG09/09 | 3005075 | prospecting/sampling | 602150/5398282 | 8 |
| AUG13/08 | 3005077 | prospecting/sampling | 601349/5399558 | 8 |
| OCT15/08 | 3005077 | prospecting/sampling | 601324/5399840 | 8 |
| MAY07/09 | 3005077 | <i>pressure wash</i> | 601323/5399620 | 8 |
| MAY19/09 | 3005077 | prospecting/sampling | 601093/5399113 | 4 |
| MAY20/09 | 3005077 | prospecting/sampling | 600998/5399955 | 8 |
| MAY24/09 | 3005077 | prospecting/sampling | 601023/5399856 | 8 |
| JUNE13/09 | 3005077 | prospecting/sampling | 601541/5399323 | 8 |
| OCT15/09 | 3005077 | prospecting/sampling | 601079/5399479 | 8 |
| JUNE02/08 | 3005078 | prospecting/sampling | 602326/5401482 | 8 |
| JUNE03/08 | 3005078 | prospecting/sampling | 601975/5401502 | 8 |
| JUNE04/08 | 3005078 | <i>pressure wash</i> | 601186/5401839 | 8 |
| JUNE05/08 | 3005078 | prospecting/sampling | 601188/5401828 | 8 |
| JULY04/08 | 3005078 | prospecting/sampling | 601652/5401087 | 8 |
| JUNE14/09 | 3005078 | prospecting/sampling | 601709/5401458 | 8 |
| JUNE15/09 | 3005078 | prospecting/sampling | 601349/5401546 | 8 |
| Total hours prospecting | | | | 152 |
| Two men | | | | 304 |
| | | | | 19 FULL DAYS |

I Andrew Baxter of Thunder Bay Ontario has worked the above mentioned dates for Doug Kakeeway
 Dated April 9 2010, Thunder Bay Ont. Signed 


I Doug Kakeeway of Thunder Bay Ontario has worked the above mentioned dates.
 Dated Apr 19 2010, Thunder Bay Ont. Signed 


TABLE 4

BACKHOE DAILY LOG


Note: Operator and one helper is included in the hourly price of backhoe

| Date | Claim No. | Description of Work Done | trench/stripping length | trench/stripped location UTM/NAD 83/ Zone 16U | Hours Worked per day |
|----------------------------|-----------|-----------------------------|-------------------------|---|----------------------|
| JULY21/09 | 3005075 | mechanical trenching | 86 meters | 601435/5398523 | 8 |
| JULY23/09 | 3005075 | mechanical trenching | 16 meters | 601439/5398576 | 5 |
| JULY24/09 | 3005075 | mechanical trenching | 18 meters | 601458/5398598 | 5 |
| AUG13/09 | 3005075 | mechanical trenching | 11 meters | 601482/5398566 | 4 |
| AUG14/09 | 3005075 | mechanical trenching | 31 meters | 601454/5398565 | 4 |
| AUG15/09 | 3005075 | mechanical trenching | 21 meters | 601488/5398578 | 5 |
| AUG16/09 | 3005075 | mechanical trenching | 19 meters | 601526/5398534 | 5 |
| SEPT19/09 | 3005075 | mechanical <i>stripping</i> | 324 sq meters | 601328/5398819 | 6 |
| SEPT20/09 | 3005075 | mechanical trenching | 19 meters | 601366/5398814 | 4 |
| SEPT21/09 | 3005075 | mechanical trenching | 38 meters | 601392/5398827 | 4 |
| SEPT22/09 | 3005075 | mechanical trenching | 24 meters | 601425/5398799 | 8 |
| OCT14/08 | 3005077 | mechanical <i>stripping</i> | 52 sq meters | 601312/5399563 | 4 |
| OCT16/08 | 3005077 | mechanical <i>stripping</i> | 1600 sq meters | 601323/5399626 | 8 |
| OCT17/08 | 3005077 | mechanical <i>stripping</i> | | 601323/5399626 | 8 |
| MAY21/09 | 3005077 | mechanical <i>stripping</i> | 105 sq meters | 601131/5400007 | 4 |
| JUNE09/09 | 3005077 | mechanical trenching | 28 meters | 601299/5399601 | 5 |
| JUNE12/09 | 3005077 | mechanical trenching | 70 meters | 601275/5399679 | 6 |
| JUNE01/08 | 3005078 | mechanical trenching | 8 meters | 602432/5400613 | 2 |
| JUNE01/08 | 3005078 | mechanical trenching | 23 meters | 602513/5400547 | 4 |
| Total Backhoe Hours | | | | | 99 |

I Andrew Baxter of Thunder Bay Ontario has worked the above mentioned dates for Doug Kakeeway

Dated April 9 2010, Thunder Bay Ont. Signed 

I Doug Kakeeway of Thunder Bay Ontario has worked the above mentioned dates.


Dated April 9 2010, Thunder Bay Ont. Signed 

APPENDIX A


LEGEND



BASALT



OVERBURDON



CLAIM POST



SWAMP



DDH



TRAIL



OUTCROP

WHITE LAKE (NARROWS GOLD PROPERTY) GEOLOGY NORTH EAST TRENCHES

MAP 1

CLAIM 3005078

NTS 42C/13

NAD 83/16U

YEAR 2009

57 METERS



SCALE


602424

602538

LEGEND




BASALT



OVERBURDON



CLAIM POST



SWAMP




DDH



TRAIL



OUTCROP



**WHITE LAKE
(NARROWS GOLD PROPERTY)
GEOLOGY
NORTH EAST TRENCHES
MAP 1
CLAIM 3005078
NTS 42C/13
NAD 83/16U
YEAR 2009**

MAGNETIC NORTH



57 METERS



SCALE

5400744

5400640



601165

601180

5402080

390627

390628

390629

**WHITE LAKE
(Narrows Gold Property)
NORTH STRIPPED ZONE
GEOLOGY
MAP 2**

**CLAIM #3005078
NTS 42C/13
NAD 83/16U
YEAR 2009**

**MAGNETIC
NORTH**



390627

390628

390629

390630

390631

7.5 METERS




SCALE

LEGEND




OVERBURDON




FELSIC



BASALT



DIABASE DIKE



OUTCROP






390624 **SAMPLE**



WHITE LAKE

5402050



LEGEND

 STRIPPED (GABBRO)

 OUTCROP (GABBRO)

 OVERBURDON

 TRAIL

 OLD GRID LINE

 SAMPLES

 CONTOUR

MAGNETIC
NORTH



716030
(FLOAT) ??

716031

WHITE LAKE
(NARROWS GOLD PROPERTY)
STRIPPED ZONE "A"
GEOLOGY
MAP 3

CLAIM #3005077
NTS 42C/12
NAD 83/16U
YEAR 2009

22 METERS



SCALE

5400250









5400228

601100

601150



LEGEND

-  SEDIMENT
-  FELSIC
-  GRANITE
-  SWAMP
-  OVERBURDON
-  OUTCROP
-  QUARTZ VEIN
-  TRAIL


5399850

5399800

601250

601300

601350




- 716028 to 716029
- 716026 to 716027 (FLOAT)
- 716013 to 716021
- 716022 to 716023
- 390646 to 390650
- 716009 to 716012
- 716001 to 716003
- 716004 to 716008

716033 (FLOAT)

716024

WHITE LAKE
(Narrows Gold Property)
GEOLOGY
ZONE "B"
MAP 4
NTS 42C12
NAD 83/16U
Claim #3005077
Year 2009



601400

601500

MAGNETIC NORTH



759612
759613
759614

759610

759611
759616

759616


759609

5398800

5398700

**WHITE LAKE
(NARROWS GOLD PROPERTY)
TRENCH ZONE D
GEOLOGY
MAP 6
CLAIM #3005075
NTS 42C/12
NAD 83/16U
YEAR 2009**

100 METERS



SCALE

LEGEND



BASALT



MINERALIZATION
(pyrite)



GRANITE



759111

SAMPLE




OUTCROP




HYDRO LINE





TRAIL





OVERBURDON











601000

601100

601200

601300

601400

5400400

5400400

5400300

5400300

5400200

5400200

5400100

5400100

601000

601100

601200

601300

601400

MAP

Legend

- StrippedAreas
- Sample#
- - - Traverse
- Trail
- Marsh
- Contours
- CLAIMPLY

Stripped Zone 'A'

3005077


716030

716031

MAY 20/09

MAY 24/09

0 10 20 40 60 80 100
Meters



601300

601400

5399100

5399100

5399000

5399000

601300

601400

Legend

- Sample#
- Trench
- StrippedAreas
- TBSutilline
- + Traverse
- Trail
- ▨ Marsh
- Contours



Stripped Zone 'C'
&
Trenches

3005075

759622 759623 759624

759617 to 21

MAP 5



601400

601500

MAY 19/09



Trench Area 'D'

3005075

759612'13'14

759610

42C/12

759611

759615

759609

759616

Legend

- Sample#
- Trench
- - - Traverse
- · - · - Hydro Line
- - - Trail
- ▨ Marsh
- · - · - Contours

5398800

5398800

5398700


5398700

601400

601500

MAP 6





MAP 7

0 15 30 60 90 120 150 Meters