

Brunswick Resources Inc.

DDH: MU-11-01RL Claims title: ABITIBI GOLD PROJECT Section:
 Township: Munro Level:
 Range: Work place: Gov. Road Coreshack
 Lot:
 Drilled by: NPLH
 Described by: W. MacRae From: 2011-05-12 Description date:
 To: 2011-05-15

Collar

Azimuth: 210.00°
 Dip: -50.00°
 Length: 347.50 m

UTM

East	554,361.00
North	5,381,554.00
Elevation	315.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	30.00	210.50°	-51.10°	No	
Reflex-Easy Shot	81.00	211.50°	-51.30°	No	
Reflex-Easy Shot	132.00	211.90°	-51.30°	No	
Reflex-Easy Shot	183.00	214.20°	-52.20°	No	
Reflex-Easy Shot	234.00	215.70°	-52.40°	No	
Reflex-Easy Shot	285.00	216.00°	-52.40°	No	
Reflex-Easy Shot	347.00	216.00°	-53.30°	No	

Description

UTM, NAD 83, Zone 17. MU-11-01RL is a re-log of MU-11-01. Re-logged by Les Kovacs Nov. 2012.

Core size: NQ Cemented: No Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	13.00	MO Overburden Casing							
13.00	35.10	VM; AK; SER Mafic Volcanic; Ankeritization; Sericite Light to medium pale olive green grey, aphanitic to fine grained, locally brecciated with scattered quartz/carb veins/patches. Local strong ankerite alteration due to proximity to faulting. Abundant micro-fracturing with quartz/carb infill predominantly at 85-90 degrees to CA but also at 35-50 degrees to CA.							
13.00	35.10	Ser Sericitisation Pale olive green to mustard yellow sericite alteration proximal to fault thereafter decreasing downhole. 13-21m intense ankerite alteration in hanging/footwall of fault.							
13.00	35.10	BRE; FRC; FAI Breccia; Fractured; Fault Fractures at 85-90 degrees to CA and 35-50 degrees with quartz/carb infill. 18.5 to 19.6 Fault with rubble and intense ankerite alteration. approx. 50cm of lost core.							
13.00	35.10	Cp; Py; As Chalcopyrite; Pyrite; Arsenopyrite Trace scattered clumpy CPY with lesser PY and ASP as very fine specks.	17.00	18.00	24451	1.00	10	11	
			18.00	20.20	24452	2.20	46		
18.90	22.20	FL;15%;;;55°;; Flooding 15% 55° 15% scattered quartz/carb veining/patches and fracture fill due to faulting. Tr. clumpy CPY.	20.20	20.50	24453	0.30	24		
			20.50	21.30	24454	0.80	26		
			21.30	21.70	24455	0.40	786		
			21.70	22.65	24456	0.95	204		
			22.65	24.00	24457	1.35	20		
			24.00	25.00	24458	1.00	18		
			25.00	26.00	24459	1.00	21		
			26.00	27.00	24460	1.00	11		
			27.00	28.00	24461	1.00	16		
			28.00	29.00	24462	1.00	8		
			29.00	30.00	24464	1.00	8		
			29.00	30.00	24463 (Std)	1.00	2,522		2.61
			30.00	31.00	24465	1.00	16		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
35.10	138.90	VMP Mafic Volcanic Pillowed Medium to dark green grey, aphanitic to very fine grained, hard, massive to very weakly foliated at 60 degrees to CA. Local and scattered chloritic and epidote altered selvages throughout. Local and scattered flow brecciation with minor carb-ankerite infill. 10% fracture controlled quartz/carb stringers.	31.00	32.25	24466	1.25	8		
			32.25	33.00	24468	0.75	8		
			32.25	33.00	24467 (Bln)	0.75	7		
			33.00	34.00	24469	1.00	5		
35.10	138.90	FL Foliation Poorly developed F1 from 60-70 degrees to CA.	68.60	69.60	010301	1.00	<5	<5	
			69.60	70.00	010302	0.40	<5		
			70.00	71.00	010303	1.00	<5		
			79.00	80.00	24470	1.00	7		
			80.00	80.75	24471	0.75	8		
			80.75	81.90	24473	1.15	5		
			80.75	81.90	24472 (Dbl)	1.15	5		
			81.90	83.00	24474	1.10	8		
			86.50	87.50	010304	1.00	<5		
			87.50	87.90	010305	0.40	<5		
87.90	88.90	010306	1.00	<5					
35.10	116.40	Py; Po Pyrite; Pyrrhotite Trace to <1% anhedral Py and lesser fine PO commonly rimming quartz/carb stringers 69.8-2cm quartz/carb altered selvage with 2-3% fine PY, Tr. PO. 87.65 to 87.75 10cm fractured pillow selvage with quartz/carb stringers and 3-5% fine anhedral PY and 1-2% PO.							
116.40	138.50	Py; Po Pyrite; Pyrrhotite <1% scattered fine specks and stringers anhedral PY, Tr. PO.							
138.70	153.00	Ser Sericitisation Light pale olive green sericite bleaching.							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
138.70	153.00	Py Pyrite <.5% fine PY.							
138.90	153.00	VM; SER Mafic Volcanic; Sericite Generally as from 13 to 35.1. Light to medium pale olive green grey, aphanitic to fine grained, somewhat hard, moderately sericite bleached-altered mafic volcanic. Barely visibly relict pillow selvages throughout with increased sericite and locally pale green chlorite alteration. Poorly developed F1 at 65-70 degrees to CA. 5% scattered quartz/carb stringers. Local faint flow brecciation.							
138.90	153.00	FL Foliation F1 from 65-70 degrees to CA.	138.95	139.75	24475	0.80	10	10	
			139.75	141.00	24476	1.25	12		
			141.00	142.00	24477	1.00	15		
			142.00	143.00	24478	1.00	17		
			143.00	144.00	24479	1.00	12		
			144.00	145.00	24480	1.00	14		
			144.85	151.00	24488 (Bln)	6.15	22		
			145.00	146.00	24481	1.00	12		
			146.00	147.00	24482	1.00	14		
			147.00	148.00	24483	1.00	11		
			148.00	148.80	24485	0.80	17		
			148.00	148.80	24484 (Std)	0.80	5,822		5.76
			148.80	149.40	24486	0.60	16		
			149.40	149.85	24487	0.45	21	25	
			149.85	151.00	24489	1.15	12		
			151.00	152.00	24490	1.00	15		
			152.00	153.00	24491	1.00	11		
153.00	202.30	VMP Mafic Volcanic Pillowed As from 35.1 to 138.9. Medium to dark green grey pillowed mafic volcanic. Abundant relict selvages. Local flow brecciation.							
153.00	202.30	FL Foliation Poorly developed F1 at 65-70 degrees to CA.	163.60	165.00	010307	1.40	<5		
			180.00	181.50	010308	1.50	<5		
			198.00	199.50	010309	1.50	<5		
201.00	202.30	Ser							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
201.00	202.30	Sericitisation Increasing sericite bleaching to 202.3m. Py Pyrite <.5% fine PY with local Tr. PO. commonly rimming selvages.							
202.30	213.00	VMP; SER Mafic Volcanic Pillowed; Sericite As from 138.9 to 153m. Section with light pale olive green sericite bleaching throughout. Local (10-15%) brecciated quartz/carb stringers and patched suggesting sutured shear/faulting. Minor fracturing with quartz/carb infill parallel F1 at 60-70 degrees to CA.							
202.30	213.00	Ser Sericitisation Moderate pale olive green sericite alteration.							
202.30	213.00	FL Foliation F1 at 60-70 degrees to CA. Possible sutured fault from 205.4 to 206.1parallel F1.							
202.30	213.00	Py; As Pyrite; Arsenopyrite Overall <.5% fine PY but 2-3% PY with 2-3% fine specks ASP rimming quartz/carb stringers within suggested sutured fault at 205.6m.	203.00	204.00	24493	1.00	11		
			203.00	204.00	24492 (Dbl)	1.00	11		
			204.00	205.35	24494	1.35	32		
			205.35	206.10	24495	0.75	1,201		1.20
			206.10	207.00	24496	0.90	23		
			207.00	208.00	24497	1.00	21		
213.00	339.00	VMP Mafic Volcanic Pillowed As from 153 to 202.3m. Pillowed mafic volcanic-abundant selvages and local flow breccia. Unit is somewhat homogeneous appearing. Local light sericite alteration. 5% scattered quartz/carb veins and stringers. 235.3 to 236.2 5% scattered irregular quartz/carb stringers commonly parallel F1 with .5 to 1% fine specks PY. 236.2 to 239.4 Increased flow brecciation with chlorite infill and .5% fine specks PY.							
213.00	216.00	Ser Sericitisation Light pale olive green sericite alteration from hangingwall unit decreasing abruptly after 216m.							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
213.00	347.50	FL Foliation Poorly developed F1 at 60-70 degrees to CA.	224.10	225.30	24498	1.20	11		
			225.30	225.65	24499	0.35	11	11	
			225.65	226.45	24500	0.80	18		
			234.00	235.00	010310	1.00	<5		
			235.00	236.20	010311	1.20	<5		
			236.20	237.20	010312	1.00	<5		
			237.20	238.50	010313	1.30	<5	<5	
			238.50	240.00	010314	1.50	<5		
213.00	339.00	Py Pyrite <.5% fine PY commonly rimming selvages or as fine fracture fill.							
239.40	315.40	VMP Mafic Volcanic Pillowed Homogeneous appearing flow with rare selvages.	264.00	265.50	010315	1.50	<5		
			279.00	280.50	010316	1.50	<5		
			294.00	295.50	010317	1.50	<5		
			314.40	315.40	24501	1.00	12		
315.20	318.00	Ser Sericitisation Disruptive appearance due to increased selvages and flow breccia with sericite and epidote alteration. (Sampled).	315.40	316.30	24502	0.90	10		
			316.30	317.50	24503	1.20	10		
			317.50	318.50	24504	1.00	17		
339.00	347.50	VM; SER Mafic Volcanic; Sericite As from 202.3 to 213m. Light to medium olive green to mustard yellow sericite overprinted pillowed mafic volcanic. (all sampled).							
339.00	347.50	Ser Sericitisation Light pale olive green to locally mustard yellow sericite alteration.							
339.00	347.50	Py Pyrite <.5% fine PY.	339.00	340.00	24505	1.00	13		
			340.00	341.00	24506	1.00	10		
			341.00	341.70	24507	0.70	11		
			341.70	342.70	24508	1.00	12		
			342.70	343.70	24509	1.00	16		
			343.70	345.00	24510	1.30	31		
			345.00	346.00	24511	1.00	29	34	
			346.00	347.00	24512	1.00	21		
			347.00	347.50	24513	0.50	18		
			347.00	347.50	24514 (Std)	0.50	5,518		5.69

347.50

End of DDH

Number of samples: 74

Number of QAQC samples: 7

Total sampled length: 74.70

Brunswick Resources Inc.

DDH: MU-11-02RL

Claims title: ABITIBI GOLD PROJECT

Section:

Township: Munro

Level:

Range:

Work place: Gov. Rd. Coreshack

Drilled by: NPLH

Lot:

Described by: W. MacRae

From: 2011-05-15

Description date:

To: 2011-05-17

Collar

UTM

Azimuth: 220.00°

East 554,976.00

Dip: -50.00°

North 5,381,590.00

Length: 386.00 m

Elevation 315.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	15.00	219.40°	-51.80°	No	
Reflex-Easy Shot	75.00	208.40°	-51.40°	No	
Reflex-Easy Shot	126.00	207.10°	-50.90°	No	
Reflex-Easy Shot	177.00	207.10°	-50.50°	No	
Reflex-Easy Shot	229.00	225.60°	-49.70°	No	
Reflex-Easy Shot	282.00	225.10°	-49.40°	No	
Reflex-Easy Shot	330.00	227.40°	-48.80°	No	
Reflex-Easy Shot	386.00	227.70°	-48.50°	No	

Description

UTM, Nad 83, Zone 17 MU-11-02RL is a re-log of MU-11-02. Re-logged by Les Kovacs Nov. 2012.

Core size: NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	1.30	MO Overburden Casing.							
1.30	55.70	VMP Mafic Volcanic Pillowed Medium to dark green grey, aphanitic to fine grained, hard, massive to weakly foliated at 60-70 degrees to CA. Abundant local pillow selvages with chlorite and locally epidote alteration. Lesser flow brecciation with chlorite, epidote and lesser sericite fracture fill. Local hematite on rare slips and fractures at 0 to 25 degrees to CA. <5% quartz/carb veins/stringers/patches.							
1.30	55.70	Ser Sericitisation Very local weak sericite bleaching from 37 to 42.3m due to weak slips.							
1.30	55.70	FL Foliation Poorly developed F1 at 60-70 degrees to CA. Weak slips with minor brecciation and quartz/carb injection at 37-37.2m and 41 to 41.2m.							
1.30	55.70	Py Pyrite Overall <5% fine PY commonly rimming selvages.	36.00	37.50	010318	1.50	<5		
			37.50	38.50	010319	1.00	<5		
			38.50	39.50	010320	1.00	<5		
			39.50	40.50	010321	1.00	<5		
			40.50	41.50	010322	1.00	<5		
			41.50	42.50	010323	1.00	<5		
55.70	95.40	I2 Intermediate intrusive Medium pale buff to buff grey, fine grained, very hard, massive intermediate intrusive. 55.7 to 60.2-assimilated mafic volcanics fragments. Other local sections of assimilated mafic volcanics common. Lower contact at 15 degrees to CA.							
55.70	158.50	FL Foliation Very poorly developed F1 from 60-70 degrees to CA. Minor brecciation with local brecciated quartz/carb stringers at 126.5 to 129.6.	56.00	57.00	24518	1.00	9		
			57.00	58.10	24515	1.10	11		
			58.10	59.00	24516	0.90	13		
			59.00	60.20	24517	1.20	10		
			60.20	61.20	24519	1.00	11		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
55.70	67.70	Py Pyrite Trace very fine PY.							
67.70	95.40	Py Pyrite Trace to <.5% very fine PY.	78.70	79.70	24520	1.00	10		
79.60	82.80	FBX Flow Breccia Interbedded flow breccia at what appears to be at 0-25 degrees to CA with .5 to 1% fine anhedral specks fracture/breccia controlled PY. (sampled).	79.70	80.70	24521	1.00	11		
			80.70	81.70	24523	1.00	18	18	
			80.70	81.70	24522 (Dbl)	1.00	10		
			81.70	82.70	24524	1.00	17		
			94.40	95.30	24525	0.90	15		
			95.30	96.00	24526	0.70	17		
95.40	158.50	VM Mafic Volcanic Generally as from 1.3 to 55.7 but with much decreased pillow selvages and flow breccia. Medium to dark green grey, hard, massive, aphanitic to fine grained, locally amygdaloidal mafic flow. Initially locally flow brecciated from 95.4 to 102.8m with minor sericite alteration (sampled). Somewhat homogeneous appearing from 102 to 158.5.	96.00	97.10	24528	1.10	16		
			96.00	97.10	24527 (Std)	1.10	2,280		2.26
			97.10	98.20	24529	1.10	14		
			98.20	99.00	24530	0.80	17		
			99.00	100.00	24531	1.00	15		
			100.00	101.00	24533	1.00	15		
			100.00	101.00	24532 (Dbl)	1.00	16		
			101.00	101.80	24535	0.80	14	13	
			101.00	101.80	24534 (Bln)	0.80	17		
			101.80	102.80	24536	1.00	12		
			126.00	127.00	010324	1.00	<5		
95.40	102.80	Ser Sericitisation Local and minor sericite alteration of flow breccia.							
95.40	126.90	Py Pyrite <.5% fine PY.							
126.90	129.60	Py Pyrite 1-2% PY as fine stringers, bands locally and as fine specks rimmins breccia fragments.	127.00	128.00	010325	1.00	<5	<5	
			128.00	129.00	010326	1.00	<5		
			129.00	130.00	010327	1.00	<5		
129.60	158.50	Py Pyrite <.5% fine PY.							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
158.50	179.50	I2 Intermediate intrusive Generally as from 55.7 to 95.4m. Medium grey to pale buff grey, fine grained, weakly, finely and locally hematitic, very hard and massive.							
158.50	169.90	Hem Hematisation Very weak, fine and spotty hematite altered.							
158.50	169.90	FL Foliation Very weakly developed F1 at 65-70 degrees to CA.							
158.50	179.50	Py Pyrite Trace PY.							
179.30	179.50	FAI Fault Fault at 50 degrees to CA with fissile core and mylonite.							
179.50	204.00	VM Mafic Volcanic As from 95.4 to 158.5m. Dark grey green, massive flow. 5% scattered locally brecciated quartz/carb veins and stringers. Local sericite alteration to end of hole as alteration proximal to footwall fault. Local minor hematitic slips.							
179.50	204.00	FL Foliation Massive to weakly foliated (F1) at 55-60 degrees to CA.							
179.50	204.00	Py Pyrite <.5% odd specks PY.	200.00	201.00	010328	1.00	<5		
200.30	200.40	VEI;0;Cl;40°;Py<.5; Vein 0 Chlorite 40° Pyrite <.5 10cm quartz/carb vein with Tr. PY at margins.							
201.00	204.00	Ser Sericitisation Weak increasing to moderate sericite bleaching due to footwall fault.							
204.00	273.60	VM; BX; SER Mafic Volcanic; Breccia; Sericite Overall light pale buff to olive green to mustard yellow, aphanitic to very fine grained, locally brecciated strongly to pervasively sericite altered mafic volcanic due to intense two faults. 5% scattered irregular and discontinuous quartz/carb veins and stringers. (Already sampled from 204							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
204.00	273.60	to 270m). Ser Sericitisation Light pale buff to mustard colored strongly to locally pervasively sericite alteration due to faulting.							
204.00	273.60	FAI Fault Poorly developed F1 at 55-60 degrees to CA. Fractures at 70-80 degrees to CA and less commonly at 25-30 degrees to CA with carb/ankerite infill. Fault at 210.4 to 211.2 with 100% rubble and fissile core. Fault at 222.5 to 223.7 with 100% dry, angular rubble-brittle fault.							
204.00	273.60	Py Pyrite Trace very fine Py commonly rimming brecciated and discontinuous quartz/carb stringers.	204.00	205.00	24537	1.00	16		
			205.00	206.00	24538	1.00	16		
			206.00	207.00	24539	1.00	15		
			207.00	208.00	24540	1.00	16		
			208.00	209.00	24541	1.00	31		
			209.00	210.00	24542	1.00	24		
			210.00	211.00	24543	1.00	19		
			211.00	212.00	24544	1.00	49		
			212.00	213.00	24545	1.00	45		
			213.00	214.00	24546	1.00	32		
			214.00	215.00	24547	1.00	26	22	
			215.00	216.00	24548	1.00	33		
			216.00	217.00	24549	1.00	27		
			217.00	218.00	24550	1.00	21		
			218.00	219.00	24551	1.00	22		
			219.00	220.00	24552	1.00	16		
			220.00	221.00	24553	1.00	16		
			221.00	222.00	24555	1.00	14		
			221.00	222.00	24554 (Dbl)	1.00	17		
			222.00	223.00	24556	1.00	11		
			223.00	224.00	24557	1.00	12		
			224.00	225.00	24558	1.00	11		

Brunswick Resources Inc.

Description	Assay						
	From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
	225.00	226.00	24559	1.00	13	12	
	226.00	227.00	24561	1.00	14		
	226.00	227.00	24560 (Std)	1.00	2,494		2.57
	227.00	228.00	24562	1.00	13		
	228.00	229.00	24563	1.00	13		
	229.00	230.00	24564	1.00	13		
	230.00	231.00	24565	1.00	14		
	231.00	232.00	24566	1.00	12		
	232.00	233.00	24567	1.00	11		
	233.00	234.00	24568	1.00	13		
	234.00	235.00	24570	1.00	18		
	234.00	235.00	24569 (Bln)	1.00	13		
	235.00	236.00	24571	1.00	30	28	
	236.00	237.00	24572	1.00	22		
	237.00	238.00	24573	1.00	24		
	238.00	239.00	24574	1.00	13		
	239.00	240.00	24575	1.00	17		
	240.00	241.00	24576	1.00	8		
	241.00	242.00	24578	1.00	18		
	241.00	242.00	24577 (Dbl)	1.00	8		
	242.00	243.00	24579	1.00	15		
	243.00	244.00	24580	1.00	36		
	244.00	245.00	24581	1.00	54		
	245.00	246.00	24583	1.00	12	14	
	245.00	246.00	24582 (Bln)	1.00	11		
	246.00	247.00	24584	1.00	8		
	247.00	248.00	24585	1.00	13		
	248.00	249.00	24586	1.00	13		
	249.00	250.00	24587	1.00	48		
	250.00	251.00	24588	1.00	15		
	251.00	252.00	24589	1.00	13		
	252.00	253.00	24590	1.00	13		
	253.00	254.00	24591	1.00	8		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
273.60	386.00	<p>VMP Mafic Volcanic Pillowed</p> <p>Medium to dark green grey, aphanitic to fine grained, massive to poorly developed F1, locally abundant pillow selvages and somewhat lesser flow breccia both with variable chlorite and lesser epidote alteration. Overall 5% quartz/carb veining and stringers with approx 50% being from .5 to 2cm and at 25-50 degrees to CA with the balance of 50% being an earlier generation and discontinuous at various angles to CA.</p> <p>280.3 to 287.2 10-15% quartz/carb stringers/veins from .5 to 2cm as well as 40% discontinuous and locally brecciated quartz/carb veins/patches and stringers with .5% fine PY.</p>	254.00	255.00	24593	1.00	21	8	5.86
			254.00	255.00	24592 (Std)	1.00	5,740		
			255.00	256.00	24594	1.00	9		
			256.00	257.00	24595	1.00	10		
			257.00	258.00	24596	1.00	10		
			258.00	259.00	24597	1.00	13		
			259.00	259.80	24598	0.80	7		
			259.80	261.00	24599	1.20	7		
			261.00	262.00	24600	1.00	9		
			262.00	263.00	24601	1.00	8		
			263.00	264.00	24602	1.00	13		
			264.00	265.00	24603	1.00	8		
			265.00	266.00	24604	1.00	11		
			266.00	267.00	24606	1.00	43		
			266.00	267.00	24605 (Bln)	1.00	9		
			267.00	268.00	24607	1.00	10		
			268.00	269.00	24608	1.00	8		
			269.00	270.00	24610	1.00	7		
			269.00	270.00	24609 (Dbl)	1.00	8		
			270.00	271.00	24611	1.00	8		
			271.00	272.00	24612	1.00	7		
			272.00	272.65	24613	0.65	6		
			272.65	273.60	24614	0.95	10		

Brunswick Resources Inc.

Description			Assay					
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)
273.60	386.00	290.2 to 290.7 50cm chlorite altered selvage at 0 degrees to CA with 10-15% coarse euhedral and fracture fill PY. Ser Sericitisation Weak and locally scattered sericite bleaching.						
273.60	313.20	FL Foliation Poorly developed F1 at 60-70 degrees to CA.						
273.60	386.00	Py Pyrite Overall <.5% fine specks PY with local coarser clumps euhedral PY to 10-15%.	279.00	280.50	010329	1.50	<5	
			280.50	282.00	010330	1.50	<5	
			282.00	283.50	010331	1.50	<5	
			283.50	285.00	010332	1.50	<5	
			285.00	286.50	010333	1.50	<5	
			286.50	288.00	010334	1.50	<5	
			288.00	289.50	010335	1.50	<5	
			289.50	291.00	010336	1.50	<5	
			291.00	292.50	010337	1.50	<5	<5
			313.00	314.00	010338	1.00	<5	
313.20	313.60	FAI Fault 40cm brittle fault with angular rubble.						
313.60	344.30	FL Foliation Poorly developed F1 from 60-70 degrees to CA.	344.00	345.00	010339	1.00	<5	
344.30	344.70	FAI Fault 40cm brittle fault with 90% angular rubble.						
344.70	386.00	FL Foliation Weak F1 at 60-70 degrees to CA.	385.00	386.00	24615 (Std)	1.00	3,782	3.90
386.00	End of DDH Number of samples: 110 Number of QAQC samples: 13 Total sampled length: 114.20							

Brunswick Resources Inc.

DDH: MU-11-03RL

Claims title: ABITIBI GOLD PROJECT

Section:

Township: Munro

Level:

Range:

Work place: Gov. Rd. Coreshack

Drilled by: NPLH

Lot:

Described by: W. MacRae

From: 2011-05-18

Description date:

To: 2011-05-19

Collar

UTM

Azimuth: 196.00°

East 554,530.00

Dip: -50.00°

North 5,381,725.00

Length: 405.00 m

Elevation 313.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	30.00	196.50°	-50.40°	No	
Reflex-Easy Shot	81.00	200.20°	-50.10°	No	
Reflex-Easy Shot	132.00	201.00°	-49.80°	No	
Reflex-Easy Shot	183.00	201.40°	-49.70°	No	
Reflex-Easy Shot	234.00	202.30°	-48.50°	No	
Reflex-Easy Shot	285.00	202.90°	-48.10°	No	
Reflex-Easy Shot	350.00	202.90°	-48.10°	No	
Reflex-Easy Shot	405.00	203.30°	-47.80°	No	

Description

UTM, Nad 83, Zone 17 MU-11-03RL is a re-logged of MU-11-03. Re-logged by Les Kovacs Nov. 2012.

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	14.25	MO Overburden Casing.							
14.25	34.50	MFP Mafic Pyroclastic Unit is medium to dark grey green, aphanitic to fine grained, hard, massive, locally siliceous and composed of variably altered mafic fragments/tuff with a siliceous/chlorite matrix. <5% mostly discontinuous and fracture fill carb/calcite. Rare slips with hematite staining.							
14.25	34.50	Ser Sericitisation Very weak and patchy sericite bleaching increasing after 32m due to footwall alteration.							
14.25	34.50	FL Foliation Very poorly developed F1 at 60-70 degrees to CA.							
14.25	34.50	Py Pyrite Trace fine PY.							
34.50	44.50	VM; SER Mafic Volcanic; Sericite Light medium pale olive green to pale mustard yellow, fine grained, massive, variably sericite bleached mafic volcanic. Locally texture is obliterated by intensity of alteration. (already sampled from 34.5 to 44.5m).							
34.50	44.50	Ser; Ank Sericitisation; Ankeritisation Predominantly variably moderate to strong sericite bleached. Local and patchy ankerite.							
34.50	44.50	FL Foliation Very poorly developed F1 at 60-70 degrees to CA.							
34.50	44.50	Py Pyrite Trace very fine PY.	34.50	35.65	24616	1.15	6		
			35.65	36.60	24617	0.95	7		
			36.60	37.60	24618	1.00	7		
			37.60	38.60	24619	1.00	10	9	
			38.60	39.60	24620	1.00	11		
			39.60	40.60	24621	1.00	19		
			40.60	41.60	24622	1.00	6		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
44.50	165.00	VMP Mafic Volcanic Pillowed Medium to dark green grey, aphanitic to fine grained, massive, hard, pillow mafic volcanic with scattered pillow selvages and lesser flow breccia with chlorite and lesser epidote and sericite alteration. Local hematitic slips and fractures. <5% scattered, discontinuous and locally brecciated quartz/carb stringers and within selvages.	41.60	42.60	24623	1.00	9		
			42.60	43.60	24624	1.00	8		
			43.60	44.60	24625	1.00	20		
44.50	165.00	Epi; Ser Epidotisation; Sericitisation Local, scattered and variably weak sericite and epidote bleaching most commonly within selvages and flow breccia.							
44.50	165.00	FL Foliation Very poorly developed F1 at 55-70 degrees to CA. Brittle fault with angular rubble and 2-3% PY at 111 to 111.3m.	58.50	60.00	010340	1.50	<5		
			91.50	93.00	010341	1.50	<5		
			109.70	110.70	010342	1.00	<5		
			110.70	111.70	010343	1.00	<5		
			111.70	112.70	010344	1.00	<5		
			122.50	123.50	010345	1.00	<5		
			123.50	124.50	010346	1.00	<5		
44.50	123.90	Py Pyrite <.5% PY as fine grains rimming local selvages and within flow breccia.							
123.90	124.20	Py Pyrite 30cm flow breccia section with grey siliceous material and stringers with 3-5% fine euhedral PY.							
124.20	165.00	Py Pyrite .5% fine PY most commonly within selvages and flow breccia.	124.50	125.50	010347	1.00	<5		
165.00	181.00	VM; SER Mafic Volcanic; Sericite Light to medium pale olive green to pale mustard yellow, aphanitic to very fine grained, massive to weakly sheared and schistose, sericite overprinted mafic volcanic with original texture almost obliterated by alteration. Locally up to 10-20% veins and stringers with fine fracturing as possible sutured fault. (already sampled from 165 to 181m).							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
165.00	181.00	Ser Sericitisation Moderate sericite alteration from 165 to 171m increasing to strong and pervasive from 171 to 176 then decreasing to moderate to low from 176 to 181m. (already sampled from 165 to 181m).							
165.00	173.50	FRC Fractured Finely fractured with main fracture pattern at 55 degrees to CA parallel F1 with a secondary and weaker fabric at 55-60 degrees perpendicular F1.							
165.00	181.00	Py Pyrite Overall <.5% fine accumulations and specks PY irregardless of alteration of quartz/carb injection.	165.00	165.90	24626	0.90	7		
			165.90	167.00	24627	1.10	5		
			167.00	168.00	24628	1.00	11		
			168.00	169.00	24630	1.00	15		
			168.00	169.00	24629 (Std)	1.00	3,844		3.84
			169.00	170.00	24631	1.00	8	8	
			170.00	171.00	24632	1.00	6		
			171.00	172.00	24633	1.00	9		
			172.00	173.00	24634	1.00	9		
			173.00	174.00	24635	1.00	7		
173.50	175.50	BRE; FRC; FAI Breccia; Fractured; Fault Increasingly brecciated and fractured with locally 10-20% quartz/carb veins & stringers parallel F1 as possible sutured fault.	174.00	175.00	24636	1.00	12		
			175.00	175.70	24638	0.70	7		
			175.00	175.70	24637 (Bln)	0.70	10		
175.50	181.00	FRC Fractured Much decreased fracturing within footwall of sutured fault with F1 and s1 at 55-60 degrees to CA.	175.70	176.70	24640	1.00	6		
			175.70	176.70	24639 (Dbl)	1.00	10		
			176.70	177.70	24641	1.00	6		
			177.70	178.70	24642	1.00	22		
			178.70	179.70	24643	1.00	14	12	
			179.70	180.00	24644	0.30	11		
			180.00	181.00	24645	1.00	9		
181.00	216.00	VMP Mafic Volcanic Pillowed As from 44.5 to 165. Pillow mafic volcanic with textbook selvages and local flow breccia. Local minor slips & fractures with hematite.							
181.00	216.00	Ser; Chl Sericitisation; Chloritisation							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
181.00	216.00	Weakly and locally chlorite and sericite altered selvages and flow breccia infill. FL Foliation Very poorly developed F1 at 55-60 degrees to CA.							
181.00	216.00	Py Pyrite <.5% fine specks PY commonly rimming selvages and within matrix of flow breccia.							
216.00	221.00	VM; SER Mafic Volcanic; Sericite Generally as from 165 to 181 although not as intensely sericite altered due to weaker sutured fault. Mafic volcanic unit weakly sericite overprinted. (already sampled from 216 to 221m).							
216.00	221.00	Ser Sericitisation Weak to locally moderate sericite bleaching.							
216.00	221.00	FL; FRC Foliation; Fractured Weak F1 but stronger S1 from 80-85 degrees to CA as sutured fault.							
216.00	221.00	30cm Sutured fault at 217.7 to 218 with siliceous granitic material injection and minor fuchsite stringers at lower contact. 2-3% fine fracture controlled PY. Py Pyrite Overall .5% fine PY. 2-3% fine fracture controlled PY within sutured fault.	216.00	216.90	24646	0.90	9		
			216.90	217.60	24647	0.70	13		
			217.60	218.10	24648	0.50	62		
			218.10	219.10	24650	1.00	11		
			218.10	219.10	24649 (Bln)	1.00	14		
			219.10	220.10	24651	1.00	11		
			220.10	221.00	24652	0.90	10		
221.00	362.80	VMP Mafic Volcanic Pillowed As from 44.5 to 165. Pillow and lesser flow brecciated mafic volcanic. Local 2-3mm amygdules. (already sampled from 281 to 284.8).							
221.00	251.00	Ser; Chl Sericitisation; Chloritisation Weak sericite, epidote and chlorite altered selvages and flow breccia matrix.							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
221.00	362.80	FL Foliation Massive to poorly developed F1 at 55-60 degrees to CA.	234.60	235.60	010348	1.00	<5		
			235.60	236.60	010349	1.00	<5	<5	
			236.60	237.60	010350	1.00	<5		
		Minor brittle fault with angular rubble at 236.1 to 236.3 with 2-3% euhedral PY.							
221.00	251.00	Py Pyrite .5 to <1% PY as local accumulations euhedral PY.							
251.00	362.80	Py Pyrite Overall .5% fine PY most commonly at rims of selvages or within matrix of flow breccia.	261.50	262.50	010351	1.00	<5		
			262.50	263.50	010352	1.00	<5		
			263.50	264.50	010353	1.00	<5		
		262.2m-2cm selvage with siliceous material and 5-8% very fine PY.							
		314.7m-3cm selvage with grey siliceous material and 5-8% fine PY.							
		317.6m-4cm selvage with chlorite and grey siliceous material and 4-6% clumpy PY.							
281.00	284.80	Ser Sericitisation Minor weak sericite bleaching with local silicification due to 10-15% quartz/carb stringers. (already sampled).	281.00	281.90	24653	0.90	12		
			281.90	283.00	24654	1.10	10		
			283.00	284.00	24656	1.00	10		
			283.00	284.00	24655 (Dbl)	1.00	10	9	
			284.00	284.80	24657	0.80	10		
284.80	362.80	Ser; Chl Sericitisation; Chloritisation Weak and local sericite and chlorite alteration of selvages and matrix of flow breccia.	313.50	314.50	010354	1.00	<5		
			314.50	315.50	010355	1.00	<5		
			315.50	316.50	010356	1.00	<5		
			316.50	318.00	010357	1.50	<5		
			318.00	319.00	010358	1.00	<5		
362.80	376.50	VM; SER Mafic Volcanic; Sericite As from 216 to 221m. Light pale olive green to mustard yellow, fractured, sericite overprinted mafic volcanic where original texture is almost obliterated by intensity of alteration possibly by sutured fault/slip from 369.2 to 369.8m. (already sample from 362.8 to 376.5m).							
362.80	376.50	Ser							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
362.80	376.50	<p>Sericitisation Variably sericite altered from weak to strong (369 to 370m) to weak from 370 to 376.5 (footwall of fault/slip).</p> <p>FL; FRC; FAI Foliation; Fractured; Fault Poorly developed F1 at 55-70 degrees to CA parallel to most fractures with a secondary (earlier) fabric at 60-70 degrees to CA perpendicular F1.</p> <p>Possible sutured fault/slip from 369.2 to 369.8m with stringers and patches brecciated quartz/carb at 30 degrees to CA perpendicular F1.</p>							
362.80	376.50	<p>Py Pyrite Overall <.5% fine diss PY with 2-3% very fine PY rimming quartz/carb injected stringers within slip fault at 369.2 to 369.8m.</p>	362.85	364.00	24659	1.15	17		
			362.85	364.00	24658 (Std)	1.15	2,484		2.54
			364.00	365.00	24660	1.00	12		
			365.00	366.00	24661	1.00	10		
			366.00	367.00	24662	1.00	11		
			367.00	368.00	24663	1.00	13		
			368.00	368.90	24664	0.90	26		
			368.90	369.80	24665	0.90	51		
			369.80	371.00	24666	1.20	16		
			371.00	372.00	24667	1.00	9	10	
			372.00	373.00	24668	1.00	9		
			373.00	374.00	24669	1.00	8		
			374.00	375.00	24670	1.00	8		
			375.00	375.50	24671	0.50	5		
			375.50	376.50	24672	1.00	6		
376.50	405.00	<p>VMP Mafic Volcanic Pillowed Generally as from 221 to 376.5m. Almost homogeneous appearing pillow mafic volcanic-medium to dark green grey, aphanitic to fine grained, massive, hard, with local selvages and lesser flow breccia.</p>							
376.50	405.00	<p>Ser Sericitisation Weak to moderate sericite bleaching from 376.5 to 380 due to proximity to hangingwall slip/fault-thereafter very weak and patchy sericite alteration.</p>							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
376.50	405.00	FL Foliation Poorly developed F1 at 55 to 70 degrees to CA.							
376.50	405.00	Py Pyrite <.5% fine PY commonly rimming local selvages.							
405.00	End of DDH Number of samples: 70 Number of QAQC samples: 6 Total sampled length: 69.05								

Brunswick Resources Inc.

DDH: MU-11-04RL

Claims title: BRUNSWICK RESOURCES INC.

Section:

Township: Munro

Level:

Range:

Work place: Gov. Rd. Coreshack

Drilled by: NPLH

Lot:

Described by: W.MacRae

From: 2011-05-24

Description date:

To: 2011-05-26

Collar

Azimuth: 191.00°
Dip: -50.00°
Length: 225.00 m

UTM

East	554,077.00
North	5,381,842.00
Elevation	307.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	51.00	191.90°	-48.50°	No	
Reflex-Easy Shot	102.00	192.60°	-48.80°	No	
Reflex-Easy Shot	153.00	193.50°	-49.00°	No	
Reflex-Easy Shot	204.00	194.40°	-49.70°	No	

Description

UTM, Nad 83, Zone 17 MU-11-04RL is a re-log of MU-11-04. Re-logged by Les Kovacs Nov. 2012.

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	6.00	MO Overburden Casing.							
6.00	19.00	VM; SER Mafic Volcanic; Sericite Light to medium green grey to locally pale yellow, aphanitic to fine grained, locally blocky, weathered and sericite altered mafic volcanic. (already sampled from 6 to 19m).							
6.00	9.80	AK; BKY Ankeritization; Blocky Blocky and ankeritic possibly due to near surface weathering.							
6.00	19.00	Ser Sericitisation Mostly sericite alteration from weak to moderate most likely due to injection from faulting.							
6.00	19.00	FL Foliation Massive to very poorly developed F1 at 50-60 degrees to CA. Fault with 100% angular rubble and fissile material from 13.4 to 14.1m.							
6.00	19.00	Py Pyrite Tr. to <.5% fine specks PY.	9.00	10.00	24673	1.00	7		
9.80	58.00	BKY Blocky Massive, decreased selvages and flow breccia and somewhat blocky with abundant hematitic slips.	10.00	11.00	24674	1.00	7		
			11.00	12.00	24675	1.00	7		
			12.00	13.00	24676	1.00	6		
			13.00	14.00	24677	1.00	18		
			14.00	15.00	24678	1.00	15		
			15.00	15.90	24679	0.90	6	7	
			15.90	17.00	24680	1.10	36		
			17.00	18.00	24682	1.00	7		
			17.00	18.00	24681 (Bln)	1.00	10		
			18.00	19.00	24683	1.00	10		
19.00	165.50	VMP Mafic Volcanic Pillowed Medium to dark green grey, aphanitic to fine grained, massive to poorly foliated, pillowed mafic volcanic. Scattered selvages and lesser flow breccia. Abundant slips and fractures with hematitic surface. 5% highly							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
		scattered quartz/carb veins & stringers from .1 to 5cm. Sharp lower contact at 35 degrees to CA.							
19.00	165.50	Hem Hematisation Abundant hematitic slips and fractures.							
19.00	165.50	FL Foliation Massive to poorly developed F1 from 55-70 degrees to CA.							
19.00	165.50	Py Pyrite Overall <.5% fine specks PY.	35.60	36.60	010359	1.00	<5		
35.70	36.50	VE;40%;Cl;;PyTr.; Veining 40% Chlorite Pyrite Tr. 40% irregular, discontinuous, barren, quartz/carb veining/patches up to 30cm with chlorite fracture fill and wall rock material. Tr. PY.							
58.00	104.60	VMP Mafic Volcanic Pillowed Increasingly foliated and pillowed with selvages and local flow breccia. 61.5 to 67 Medium grained, hard, intermediate mafic intrusive. Sharp lower contact at 45 degrees to CA.							
165.50	175.30	I2 Intermediate intrusive Medium grey, fine grained, massive, very hard, somewhat siliceous, locally blocky intermediate intrusive with 10-15% assimilated pillowed mafic volcanics. 5% local quartz/carb patches and fracture fill. Sharp but irregular lower contact.							
165.50	175.30	Sil Silicification Weak and fine pale grey silicification throughout.							
165.50	175.30	FL Foliation Massive to very poorly foliated (F1) at 55-60 degrees to CA.							
165.50	175.30	Py Pyrite <.5% fine diss PY.							
175.30	199.50	VMP Mafic Volcanic Pillowed As from 19 to 165.5 before interruption by intermediate intrusive. Pillowed and locally flow brecciated mafic volcanic. Local amygdules to 3-5mm.							

Brunswick Resources Inc.

Description			Assay					
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)
175.30	199.50	FL Foliation Poorly developed F1 at 60-65 degrees to CA.						
175.30	199.50	Py Pyrite .5% fine specks PY.						
199.50	202.70	I2 Intermediate intrusive As from 165.5 to 175.3. Internmediate intrusive. Sharp lower contact at 50 degrees to CA. 10-15% assimilated mafic volcanics. Generally finer grained that unit above.						
199.50	202.70	FL Foliation Massive to very poorly developed F1 at 60-65 degrees to CA.						
199.50	202.70	Py Pyrite Trace to <.5% fine diss PY.						
202.70	225.00	VMP Mafic Volcanic Pillowed As from 175.3 to 199.5m. Almost text book pillow mafic volcanic. Minor local flow breccia.						
202.70	225.00	FL Foliation Massive to weakly foliated (F1) at 55-60 degrees to CA.						
202.70	225.00	Py Pyrite <.5% odd specks euhedral PY.						
225.00	End of DDH Number of samples: 11 Number of QAQC samples: 1 Total sampled length: 11.00							

Brunswick Resources Inc.

DDH: MU-11-05RL

Claims title: ABITIBI GOLD PROJECT.

Section:

Township: Munro

Level:

Range:

Work place: Gov. Rd. Coreshack

Drilled by: NPLH

Lot:

Described by: W. MacRae

From: 2011-05-26

Description date:

To: 2011-05-27

Collar

Azimuth: 198.00°
Dip: -50.00°
Length: 201.00 m

UTM

East	554,043.00
North	5,381,854.00
Elevation	303.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	30.00	198.80°	-50.10°	No	
Reflex-Easy Shot	81.00	200.30°	-50.30°	No	
Reflex-Easy Shot	135.00	201.10°	-51.10°	No	
Reflex-Easy Shot	200.00	201.10°	-51.80°	No	

Description

UTM, Nad 83, Zone17 MU-11-05RL is a re-log of MU-11-05. Re-logged by Les Kovacs Nov. 2011.

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	3.00	MO Overburden Casing.							
3.00	6.60	VM Mafic Volcanic Medium to dark green grey, fine grained, massive , hard and blocky due to footwall faulting. Minor hematite on slips and fractures.							
3.00	6.60	FL Foliation Very poorly developed F1 at 55-60 degrees to CA.							
3.00	6.60	Py Pyrite Trace to <.5% fine specks PY.	5.60	6.60	010360	1.00	<5		
6.60	9.00	FZ; BKY Fault Zone; Blocky 100% angular rubble due to brittle faulting. Minor hematite staining.							
6.60	9.00	FAI Fault Brittle fault.							
6.60	9.00	Py Pyrite Trace to <.5% very fine PY.	6.60	7.60	010361	1.00	<5	<5	
			7.60	9.00	010362	1.40	<5		
9.00	99.80	VM; BKY Mafic Volcanic; Blocky As from 3 to 6.6. Massive mafic volcanic. Blocky from 9.0 to 24m due to proximity to hangingwall fault. Abundant hematitic slips and fracture fill. 24 to 99.9m-unit is increasingly massive and somewhat homogeneous appearing. 5% scattered quartz/carb veins/stringers.							
9.00	99.80	Epi; Hem Epidotisation; Hematisation Minor hematite on slips and local and minor epidote as relic selvage alteration.							
9.00	24.00	BL Blocky Blocky but decreasing to 24m due to footwall faulting.							
9.00	99.80	Py Pyrite Overall <.5% odd specks fine euhedral PY.	9.00	10.50	010363	1.50	<5		
24.00	99.80	FL Foliation Massive to poorly developed F1 from 55-70 degrees to CA.	31.30	32.30	010364	1.00	<5		
			32.30	33.30	010365	1.00	<5		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
32.60	33.00	Brittle fault from 98.1 to 98.6m with angular rubble. Str:80%;Cl;Py:5%; Stingers 80% Chlorite Pyrite .5%	33.30	34.50	010366	1.20	<5		
			98.00	99.00	010367	1.00	<5		
99.80	104.50	FZ; BKY Fault Zone; Blocky 50% angular rubble sections and 50% somewhat competent brecciated mafic volcanics. Lack of gouge/mud suggests a brittle fault. (already sampled from 102 to 104.5m).							
99.80	104.50	Ser; Epi Sericitisation; Epidotisation Minor fracture controlled sericite and epidote.							
99.80	104.50	FAI; BRE Fault; Breccia							
99.80	104.50	Py Pyrite Tr. to .5% odd fine specks fracture fill PY.	102.00	103.00	24684	1.00	7		
			103.00	104.00	24685	1.00	8		
			104.00	105.25	24686	1.25	6		
104.50	201.00	VM; TU3; VMP Mafic Volcanic; Mafic Tuff; Mafic Volcanic Pillowed Predominantly medium to dark green grey, aphanitic to fine grained, massive, hard flow brecciated mafic volcanic with sections of possible mafic tuff where composition appears to be predominantly of mafic minerals. Local relic selvages suggest some pillowed flows. <5% highly scattered mostly discontinuous quartz/carb stringers/patches at various angles to CA. Somewhat blocky throughout. (already sampled from 104.5 to 106m, 148 to 151m, 156 to 178m.)							
104.50	201.00	Epi; Ser; FU Epidotisation; Sericitisation; Fuchsite Minor and local epidote and sericite as breccia matrix and as selvage alteration and very minor fuchsite wisps and clots where sericite alteration slightly elevated.							
104.50	201.00	FL Foliation Massive to poorly developed F1 from 40 to 55 degrees to CA with fine tension fractures locally seen at 35 to 45 degrees perpendicular F1.							
104.50	201.00	Py Pyrite	105.25	106.00	24688	0.75	6		

Brunswick Resources Inc.

Description	Assay							
	From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)	
Overall <.5% fine specks PY commonly within flow breccia matrix or selvages.	105.25	106.50	24687 (Dbl)	1.25	<5			
	148.00	149.00	24690	1.00	16			
	148.00	149.00	24689 (Std)	1.00	2,020		2.30	
	149.00	150.00	24691	1.00	7	5		
	150.00	151.00	24692	1.00	5			
	156.00	157.00	24693	1.00	6			
	157.00	158.00	24694	1.00	5			
	158.00	159.00	24695	1.00	<5			
	159.00	160.00	24696	1.00	7			
	160.00	161.00	24697	1.00	5			
	161.00	162.00	24698	1.00	5			
	162.00	163.00	24699	1.00	5			
	163.00	164.00	24701	1.00	7			
	163.00	164.00	24700 (Bln)	1.00	7			
	164.00	165.00	24702	1.00	7			
	165.00	166.00	24703	1.00	6	5		
	166.00	167.00	24704	1.00	<5			
	167.00	168.00	24705	1.00	5			
	168.00	169.00	24707	1.00	13			
	168.00	169.00	24706 (Std)	1.00	3,778		3.84	
	169.00	170.00	24708	1.00	7			
	170.00	171.00	24709	1.00	<5			
	171.00	172.00	24711	1.00	<5			
	171.00	172.00	24710 (Dbl)	1.00	<5			
	172.00	173.00	24712	1.00	6			
	173.00	174.00	24713	1.00	5			
	174.00	175.00	24714	1.00	5			
	175.00	176.00	24715	1.00	10	9		
	176.00	177.00	24716	1.00	10			
	177.00	178.00	24717	1.00	10			
	201.00	End of DDH Number of samples: 37 Number of QAQC samples: 5 Total sampled length: 38.10						

Brunswick Resources Inc.

DDH: MU-11-06RL
 Claims title: BABITIBI GOLD PROJECT
 Section:
 Township: Munro
 Level:
 Range:
 Work place: Gov. Rd. Coreshack
 Lot:
 Drilled by: NPLH
 Described by: W.MacRae
 From: 2011-05-27
 Description date:
 To: 2011-06-28

Collar

Azimuth: 198.00°
 Dip: -50.00°
 Length: 216.00 m

UTM

East	553,980.00
North	5,381,866.00
Elevation	301.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	51.00	198.20°	-50.00°	No	
Reflex-Easy Shot	102.00	200.30°	-51.10°	No	
Reflex-Easy Shot	153.00	201.50°	-51.80°	No	
Reflex-Easy Shot	204.00	202.70°	-52.20°	No	

Description

UTM, Nad 83, Zone 17 MU-11-06RL is a re-log of MU-11-06. Re-logged by Les Kovacs Nov.-Dec., 2012.

Core size: NQ Cemented: No Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	6.00	MO Overburden Casing.							
6.00	13.60	VM Mafic Volcanic Medium to dark green grey, aphanitic to fine grained, hard massive mafic flow with massive to poorly developed foliation. Unit is strongly blocky (60-70%) due to proximity to footwall fault. Abundant hematitic slips and open fractures. 5% late, white band barren quartz/carb veining.							
6.00	13.60	Hem Hematisation Blocky with abundant hematitic slips and fractures due to proximity to faulting.							
6.00	13.60	FL Foliation Poorly developed F1 at 55-70 degrees to CA.							
6.00	13.60	Py Pyrite Trace to <.5% very fine PY.							
13.60	25.50	FZ; BKY; VM Fault Zone; Blocky; Mafic Volcanic 100% blocky, rubble core from 17 to 23.7m. Lack of gouge/mud suggests a brittle fault. The remainder of unit shows 50-70% blocky rubble core with drillers reporting 1.2m of lost core at 24m. Medium to dark green massive mafic flow where rock is somewhat competent and identifiable. Abundant hematitic slips and fracture fill. 5% irregular quartz/carb injection. (already sampled from 15.7 to 18.0m).							
13.60	25.50	Hem Hematisation Mostly hematite on slips and within fractures.							
13.60	25.50	FAI Fault Intense brittle faulting.							
13.60	25.50	Py Pyrite Overall <.5% odd specks fine PY.	15.70	16.60	24718	0.90	8		
			16.60	16.90	24719	0.30	11		
			16.90	18.00	24720	1.10	11		
25.50	64.10	VM Mafic Volcanic Medium green grey, aphanitic to fine grained, hard, massive mafic flow							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
		as above. 5% scattered quartz/carb veining from .5 to 2cm commonly from 35 to 60 degrees to CA. Blocky core with hematitic slips and fracture fill from 25.5 to 34m due to proximity to hangingwall fault-more massive and competent thereafter. (already sampled from 63 to 64.1m)							
25.50	37.40	Hem Hematization Hematization on slips/fractures due to faulting.							
25.50	64.10	FL Foliation Very poorly developed F1 at 60-70 degrees to CA.							
58.50	62.00	Car Carbonatisation 30% carb/calcite stringers/veining at 0 to 5 degrees to CA.							
58.50	64.10	Py Pyrite Overall <.5% fine PY.							
62.00	71.90	Car Carbonatisation Carb-ankerite alteration.	63.00	64.50	24721	1.50	11		
64.10	71.90	QZ/CA Quartz/Carb Veining Unit is a mafic volcanic flow with 40-50% quartz/carb/calcite stringers and veinlets at 0 to 5 degrees to CA with wisps and slivers of chlorite and chlorite altered wallrock material. (sampled from 64.1 to 71.9m).							
64.10	71.90	BRE Breccia Brecciated quartz/carb/calcite stringers at 0-5 degrees to CA.							
64.10	71.90	Py Pyrite .5% fine specks PY.	64.50	64.70	24722	0.20	6		
			64.70	65.80	24724	1.10	8		
			64.70	65.80	24723 (Dbl)	1.10	5		
			65.80	66.60	24725	0.80	9		
			66.60	68.20	24726	1.60	15		
			68.20	69.00	24728	0.80	10		
			68.20	69.00	24727 (Std)	0.80	2,113		2.13
			69.00	70.00	24729	1.00	7		
			70.00	71.00	24730	1.00	7		
			71.00	71.90	24731	0.90	<5		
71.90	171.00	VM Mafic Volcanic							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
		Massive, somewhat homogeneous appearing fine grained mafic flow as above at 25.5 to 64.1m. Local and scattered flow top breccia-increasing after 160m. Minor residual quartz/carb/calcite stringers from 71.9 to 73m. <5 scattered quartz/carb stringers/veins. (Already sampled from 71.9 to 73m and 170 to 171.0m)							
71.90	171.00	FL Foliation Massive to poorly developed F1 from 55-70 degrees to CA. Flow top breccia infill with quartz/carb at 0 degrees to CA.							
71.90	171.00	Py Pyrite <.5% odd fine specks PY commonly within flow top breccia.	71.90	73.00	24733	1.10	7		
			71.90	73.00	24732 (Bln)	1.10	9		
			109.50	111.00	010368	1.50	<5		
110.60	112.70	Car Carbonatisation Predominantly quartz but 30% quart/carb stringers/patches as flow breccia infill at 0 degrees to CA.	111.00	112.50	010369	1.50	<5		
			112.50	114.00	010370	1.50	<5		
			170.00	171.00	24734	1.00	7		
171.00	204.00	VM; FBX Mafic Volcanic; Flow Breccia General description as above at 71.9 to 171m but with much increased flow top breccia to 90% of unit. Rare pillow selvages. Breccia matrix composed of darker green to black wisps, clots chlorite with lesser lighter green sericite and epidote. (Already sampled from 171 to 198.0m).							
171.00	204.00	Chl; Ser Chloritisation; Sericitisation Increased chlorite as flow top breccia matrix with minor sericite and lesser epidote.							
171.00	204.00	FL; FRC Foliation; Fractured Massive to very poorly developed F1 from 60-70 degrees to CA with a local fine fracture/fabric pattern commonly seen at 80-90 degrees							
171.00	204.00	Py Pyrite <.5% very fine PY confined to flow top breccia matrix and fine fracture fill.	171.00	172.00	24735	1.00	9		
			172.00	173.00	24736	1.00	5		
			173.00	174.00	24737	1.00	6		
			174.00	175.00	24738	1.00	6		
			175.00	176.00	24739	1.00	<5	<5	
			176.00	177.00	24740	1.00	<5		
			177.00	178.00	24741	1.00	6		
			178.00	179.00	24742	1.00	9		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
204.00	216.00	VM Mafic Volcanic Generally as from 71.9 to 171m-more massive appearing mafic volcanic flow with local flow top breccia and lesser pillow selvages.	179.00	180.00	24744	1.00	13	5	2.26
			179.00	180.00	24743 (Std)	1.00	2,141		
			180.00	181.00	24745	1.00	6		
			181.00	182.00	24746	1.00	6		
			182.00	183.00	24747	1.00	9		
			183.00	184.00	24748	1.00	6		
			184.00	185.00	24749	1.00	5		
			185.00	186.00	24750	1.00	13		
			186.00	187.00	24751	1.00	<5		
			187.00	188.00	24752	1.00	<5		
			188.00	189.00	24753	1.00	<5		
			189.00	190.00	24754	1.00	7		
			190.00	191.00	24755	1.00	<5		
			191.00	192.00	24757	1.00	<5		
			191.00	192.00	24756 (Bin)	1.00	7		
			192.00	193.00	24759	1.00	26		
			192.00	193.00	24758 (Dbl)	1.00	5		
			193.00	194.00	24760	1.00	13		
			194.00	195.00	24761	1.00	8		
			195.00	196.00	24762	1.00	<5		
196.00	197.00	24763	1.00	<5	<5				
197.00	198.00	24764	1.00	5					
204.00	216.00	FL Foliation Massive to very poorly developed F1 at 60-70 degrees to CA.							
204.00	216.00	Py Pyrite <5% odd fine specks PY commonly within rare flow top breccia or selvages.							

216.00 End of DDH
Number of samples: 44
Number of QAQC samples: 6
Total sampled length: 44.80

Brunswick Resources Inc.

DDH: MU-11-07RL

Claims title: ABITIBI GOLD PROJECT.

Section:

Township: Munro

Level:

Range:

Work place: Gov. Rd. Coreshack

Drilled by: NPLH

Lot:

Described by: W.MacRae

From: 2011-05-29

Description date:

To: 2011-05-31

Collar

Azimuth: 200.00°
Dip: -50.00°
Length: 300.00 m

UTM

East	554,476.00
North	5,380,919.00
Elevation	314.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	51.00	201.70°	-50.70°	No	
Reflex-Easy Shot	102.00	201.70°	-51.00°	No	
Reflex-Easy Shot	153.00	202.10°	-51.70°	No	
Reflex-Easy Shot	198.00	206.30°	-52.30°	No	
Reflex-Easy Shot	252.00	208.40°	-52.70°	No	
Reflex-Easy Shot	297.00	209.60°	-52.90°	No	

Description

UTM, Nad 83, Zone 17 MU-11-07RL is a re-log of MU11-07. Re-logged by Les Kovacs Dec. 2012. No additional samples added.

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	6.00	MO Overburden Casing.							
6.00	166.50	VMP; FBX Mafic Volcanic Pillowed; Flow Breccia Medium to dark green, aphanitic to fine grained, massive to poorly foliated mafic pillowed and flow top breccia volcanic. Unit is mottled in appearance due to 5-10% torn-up, discontinuous quartz/carb/calcite stringers as fracture fill, selvage replacement and flow top breccia matrix. (already sampled from 90 to 93m, 165 to 166.5m).							
6.00	91.90	Ser; Sil Sericitisation; Silicification Sericite mostly as selvage replacement and local olive green to buff harder siliceous material.							
6.00	132.00	FL Foliation Very poorly developed F1 at 60-70 degrees to CA.							
6.00	166.50	Py Pyrite Overall <.5% fine specks PY but up to 2-3% euhedral PY over 20cm from 36.9 to 37.1m. And 2-3% fine euhedral specks PY from 156.1 to 156.15m.	35.50	36.50	10371	1.00	9	11	
			36.50	37.50	10372	1.00	8		
			37.50	38.50	10373	1.00	6		
			90.00	91.00	24765	1.00	<5		
			91.00	92.00	24766	1.00	7		
			92.00	93.00	24767	1.00	5		
132.00	138.00	FL; BRE Foliation; Breccia Poorly developed F1 from 60-70 degrees to CA with a secondary fracture pattern from 50-90 degrees to CA cross cutting F1 with grey siliceous material infill. (Already sampled from 132 to 138m).	132.00	133.00	24768	1.00	5		
			133.00	134.00	24769	1.00	7		
			134.00	135.00	24770	1.00	5		
			135.00	136.00	24772	1.00	<5		
			135.00	136.00	24771 (Bln)	1.00	7		
			136.00	137.00	24773	1.00	5		
			137.00	138.00	24774	1.00	<5		
138.00	166.50	FL Foliation Massive to very poorly developed F1 from 60-70 degrees to CA. Somewhat more blocky from 144 to 151m with little evidence of faulting. Minor quartz/carb stringers/veins at various angles to CA. (already sampled from 148.7 to 158.4m)	148.70	149.70	24775	1.00	7	5	
			149.70	150.40	24776	0.70	8		
			150.40	151.40	24777	1.00	<5		
			151.40	152.40	24779	1.00	8		
			151.40	152.40	24778 (Std)	1.00	2,095		2.09
			152.40	153.40	24780	1.00	7		

Brunswick Resources Inc.

Description			Assay								
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)		
166.50	169.30	FZ; GR Fault Zone; Graphitic Fault zone with rubble, minor gouge and 70% highly graphitic material-most likely altered mafic volcanic. (already sampled from 166.5 to 169.3m).	153.40	154.40	24782	1.00	<5	8			
			153.40	154.40	24781 (Dbl)	1.00	7				
			154.40	155.40	24783	1.00	6				
			155.40	156.40	24784	1.00	5				
			156.40	157.40	24785	1.00	<5				
			157.40	158.40	24786	1.00	<5				
			165.00	166.00	24787	1.00	9				
			166.00	167.00	24788	1.00	7				
166.50	169.30	FL Foliation Fault with rubble and graphitic material. F1 from competent core is 60-65 degrees to CA.	167.00	167.90	24789	0.90	11				
167.40	169.30	GR Graphite Muddy, slippery graphite alteration.									
167.40	169.30	Py Pyrite Trace to <.5% very fine PY.	167.90	168.70	24790	0.80	9				
			168.70	170.00	24792	1.30	6				
			168.70	170.00	24791 (Bln)	1.30	<5				
169.30	221.00	VM Mafic Volcanic Medium to dark green grey to weakly olive green, aphanitic to very fine grained, massive to poorly developed F1. Mafic volcanic flow with local weak flow breccia and relic selvages although not abundant. <5% fine quartz/carb stringers, veins and fracture fill at various low angles to CA. Somewhat softer and locally serpentinized starting from 175m to 221. (already sampled from 169.3 to 174m and 178 to 190m).									
			169.30	221.00	FL Foliation Poorly developed F1 from 65-70 degrees to CA.						
			169.30	221.00	Py Pyrite Overall <.5% fine diss PY.	170.00	171.00	24793	1.00	<5	
						171.00	172.00	24794	1.00	6	
			172.00	173.00	24796	1.00	<5				
			173.00	174.00	24797	1.00	6				

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
175.00	221.00	Spt Serpentinisation Locally, unit is somewhat softer and turning somewhat olive green grey which may suggest early alteration to komatiite-ultramafic.	178.00	179.00	24798	1.00	<5		
			179.00	180.00	24799	1.00	<5	<5	
			180.00	181.00	24801	1.00	<5		
			180.00	181.00	24800 (Std)	1.00	5,632		5.83
			181.00	182.00	24802	1.00	6		
			182.00	183.00	24803	1.00	5		
			183.00	184.00	24804	1.00	<5		
			184.00	185.00	24805	1.00	8		
			185.00	186.00	24806	1.00	7		
			186.00	187.00	24807	1.00	<5		
			187.00	188.00	24808	1.00	<5		
188.00	188.70	VEI;;Sr Qz;;40°;; Vein Sericite Quartz 40° 70cm hard, buff to pale pink, hard, siliceous material with two generations of torn-up white and opaque quartz. Minor sericite altered wallrock material. (already sampled).	188.00	188.70	24809	0.70	<5		
			188.70	190.00	24810	1.30	<5		
			214.00	215.20	24811	1.20	10	9	
215.00	221.00	FBX Flow Breccia Lighter green flow breccia interbedded with possible interflow sediments with darker green to black, soft serpentinization.	215.20	216.00	24812	0.80	13		
			216.00	217.00	24814	1.00	10		
			216.00	217.00	24813 (Bln)	1.00	14		
			217.00	218.00	24815	1.00	5		
			218.00	219.00	24816	1.00	9		
			219.00	220.00	24818	1.00	9		
			219.00	220.00	24817 (Dbl)	1.00	11		
			220.00	221.30	24819	1.30	6		
221.00	300.00	V4A; UM Komatiite; Ultramafic Medium to dark green grey, aphanitic to very fine grained, soft, massive to poorly foliated with local with flow brecciation. Local and scattered well developed spinifex texture suggests a komatiite grading to an ultramafic locally where unit is darker green to black, serpentinized and softer. (already sampled from 214 to 223.3m).	221.30	222.30	24820	1.00	<5		
			222.30	223.30	24821	1.00	7		
			299.00	300.00	24822 (Std)	1.00	3,670		3.70
221.00	297.00	Spt Serpentinisation Soft, locally serpentinized due to ultramafic alteration from komatiite.							
221.00	297.00	FL Foliation							

Brunswick Resources Inc.

Description		Assay						
		From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
221.00	297.00							
Very poorly developed F1 from 50-70 degrees to CA. Somewhat blocky throughout but increasingly from 257.1 to 257.6 with little evidence of faulting. Py Pyrite <.5% fine diss PY.								
300.00	End of DDH Number of samples: 52 Number of QAQC samples: 8 Total sampled length: 52.00							

Brunswick Resources Inc.

DDH: MU-11-08RL	Claims title: ABITIBI GOLD PROJECT.	Section:
	Township: Munro	Level:
	Range:	Work place: Gov. Rd. Coreshack
Drilled by: NPLH	Lot:	
Described by: W.MacRae	From: 2011-06-01	Description date:
	To: 2011-06-03	

Collar

Azimuth: 200.00°	UTM
Dip: -50.00°	East 554,686.00
Length: 327.40 m	North 5,381,355.00
	Elevation 318.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	15.00	201.50°	-48.90°	No	
Reflex-Easy Shot	66.00	201.50°	-49.40°	No	
Reflex-Easy Shot	117.00	201.60°	-50.00°	No	
Reflex-Easy Shot	168.00	201.90°	-50.10°	No	
Reflex-Easy Shot	219.00	201.70°	-49.90°	No	
Reflex-Easy Shot	270.00	201.60°	-49.70°	No	

Description

UTM, Nad 83, Zone 17 MU-11-08RL is a re-log of MU-11-08. Re-logged by Les Kovacs Dec. 2012.

Core size: NQ	Cemented: No	Stored: Yes
---------------	--------------	-------------

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	6.00	MO Overburden Casing.							
6.00	146.80	VM Mafic Volcanic Dark green, aphanitic to very fine grained, massive, somewhat soft with 5-10% wisps, clots and fracture fill dark green to black chlorite throughout. <5% highly scattered relic pillow selvages with sericite & epidote alteration. <5% scattered discontinuous/torn-up and fracture fill quartz/carb at various low angles to CA (0 to 35 degrees). Rare hematitic slips/fracture fill. (already sampled from 10.5 to 15m and 115.9 to 120.6).							
6.00	146.80	FL Foliation Very poorly developed F1 from 60-70 degrees to CA. 11.6 to 13.8m Possible sutured fault due to ragged/brecciated quartz/carb vein with accicular tourmaline at 12.5 to 12.9m. Rusty ankeritic halo with 5% scattered quartz/carb stringers from 11.6 to 13.8m. <.5% very fine PY.							
6.00	146.80	Py Pyrite <.5% odd fine specks PY with local accumulations to 2%.	10.50	11.70	24823	1.20	24	20	
11.60	13.70	Ank Ankeritisation Moderate to strong ankerite alteration due to possible sutured fault.	11.70	12.50	24824	0.80	71		
			12.50	12.90	24825	0.40	21		
			12.90	13.70	24826	0.80	120		
			13.70	15.00	24827	1.30	14		
15.00	106.80	VM Mafic Volcanic Somewhat homogeneous appearing throughout broken up by odd quartz/carb veins/stringers and rare selvages.	116.00	117.20	24828	1.20	8		
117.20	118.50	VE;60%;Cl;;Cp.5% Cv; Veining 60% Chlorite Chalcopyrite .5% Covellite 50-60% irregular, brecciated, quartz/carb veining with chlorite altered wallrock material. .5 to 1% CPY and covellite at irregular lower contact.	117.20	117.80	24829	0.60	8		
			117.80	118.55	24830	0.75	6		
			118.55	119.60	24831	1.05	5		
			119.60	120.60	24832	1.00	6		
146.80	191.00	FBX; VM Flow Breccia; Mafic Volcanic Predominantly as above at 6 to 146.8 but with increased flow brecciation and somewhat lighter color from 147 to 178.9 due to very weak and local sericite and lesser epidote alteration and breccia matrix infill. <5% fine,							

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
146.80	191.00	brecciated, fracture fill and fine irregular quartz/carb patched and stringers. BRE; FL Breccia; Foliation Mostly flow brecciated. Massive to very poorly foliated at 60-70 degrees to CA.							
147.00	179.00	Ser; Epi Sericitisation; Epidotisation Very weak, light and highly scattered sericite and lesser epidote flow breccia matrix alteration.							
147.00	191.00	Py; Cp Pyrite; Chalcopyrite Highly scattered specks and small mm scale clumps of PY and CPY from .5 to 1%.	147.00	147.80	24833	0.80	5		
			147.80	148.50	24834	0.70	7		
			148.50	149.50	24835	1.00	9	10	
			149.50	150.55	24836	1.05	11		
			150.55	152.00	24837	1.45	8		
			152.00	153.00	24838	1.00	7		
			153.00	154.00	24839	1.00	9		
			154.00	155.00	24840	1.00	7		
			155.00	156.00	24841	1.00	8		
			156.00	157.00	24843	1.00	8		
			156.00	157.00	24842 (Bln)	1.00	13		
			157.00	158.00	24844	1.00	8		
			158.00	159.00	24845	1.00	11		
			159.00	160.00	24846	1.00	9		
			160.00	161.00	24848	1.00	9		
			160.00	161.00	24847 (Std)	1.00	3,693		3.81
			161.00	162.00	24849	1.00	11		
			162.00	163.00	24850	1.00	8		
			163.00	164.00	24851	1.00	6		
			164.00	165.00	24852	1.00	<5		
			165.00	166.00	24853	1.00	8		
			166.00	167.00	24854	1.00	9		
			167.00	168.00	24856	1.00	13		
			167.00	168.00	24855 (Dbl)	1.00	8		
			168.00	169.00	24857	1.00	<5		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
191.00	327.40	VM Mafic Volcanic Generally as from 6 to 146.8m. Dark green to grey green mafic flow with local and scattered pillow selvages and flow breccia. Fine tension fractures with chlorite and possibly serpentinite suggest early alteration to komatiite. Increasingly flow brecciated from 253 to 270m but returns to massive flow after 270m. (already sampled from 252.5 to 257.2m, 287 to 294m and 325.5 to 327.5). <5% scattered quartz/carb stringers and fracture fill.	169.00	170.00	24858	1.00	8	6	
			170.00	171.00	24859	1.00	5		
			171.00	172.00	24860	1.00	<5		
			172.00	173.00	24861	1.00	7		
			173.00	174.00	24862	1.00	5		
			174.00	175.00	24864	1.00	6		
			174.00	175.00	24863 (Dbl)	1.00	<5		
			175.00	175.80	24865	0.80	<5		
			175.80	176.45	24866	0.65	<5		
			176.45	177.50	24867	1.05	8		
			177.50	178.70	24868	1.20	<5		
191.00	208.80	FL Foliation Massive to very poorly developed F1 from 60-70 degrees to CA.							
191.00	327.50	Py Pyrite Overall trace to .5% very fine PY, trace CPY commonly within selvages and flow breccia.							
208.80	327.50	FL Foliation Massive to very poorly developed F1 from 60-70 degrees to CA. Minor brittle faulting with angular weakly hematite stained rubble core but no gouge at 277.6 to 279m.							
252.50	257.20	Epi; Ser Epidotisation; Sericitisation 30% mixed epidote and sericite alteration of flow breccia.	252.60	253.60	24869	1.00	6	7	
			253.60	254.80	24870	1.20	6		
			254.80	255.40	24871	0.60	8		
			255.40	256.50	24872	1.10	6		
			256.50	257.50	24873	1.00	6		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
288.00	294.00	Ank Ankeritisation Section with 5-8% carb(ankerite)/quartz/calcite fracture fill, stringers and brecciated veins with main brecciated veining from 292.6 to 292.65m as a possible weak sutured fault.	287.00	288.20	24874	1.20	6		2.16
			288.20	289.00	24875	0.80	6		
			289.00	290.00	24876	1.00	12		
			290.00	291.00	24878	1.00	11		
			291.00	292.00	24880	1.00	6		
			291.00	292.00	24879 (Bln)	1.00	8		
			292.00	293.00	24882	1.00	15		
			292.00	293.00	24881 (Std)	1.00	2,071		
			293.00	294.00	24883	1.00	<5	<5	
			305.00	327.50	Spt Serpentinisation Local, faint and primitive spinifex texture suggests this section is grading to a komatiite.	325.60	326.60	24884	
326.60	326.90	24885				0.30	<5		
326.90	327.40	24886				0.50	<5		
326.90	327.40	24877 (Bln)				0.50	12		
327.40	End of DDH Number of samples: 57 Number of QAQC samples: 7 Total sampled length: 54.50								

Brunswick Resources Inc.

DDH: MU-11-09RL

Claims title: ABITIBI GOLD PROJECT.

Section:

Township: Munro

Level:

Range:

Work place: Gov. Rd. Coreshack

Drilled by: NPLH

Lot:

Described by: W.MacRae

From: 2011-05-29

Description date:

To: 2011-05-31

Collar

Azimuth: 200.00°
Dip: -50.00°
Length: 350.00 m

UTM

East	554,874.00
North	5,380,831.00
Elevation	314.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	18.00	202.50°	-48.80°	No	
Reflex-Easy Shot	69.00	205.90°	-48.50°	No	
Reflex-Easy Shot	120.00	206.60°	-49.00°	No	
Reflex-Easy Shot	171.00	207.40°	-48.70°	No	
Reflex-Easy Shot	222.00	212.40°	-48.40°	No	
Reflex-Easy Shot	273.00	211.80°	-48.00°	No	
Reflex-Easy Shot	324.00	210.70°	-47.80°	No	

Description

UTM, Nad 83, Zone 17, MU-11-09RL is a re-log of MU-11-09. Re-logged by Les Kovacs, Nov. 2014.

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	4.50	CA Casing Overburden							
4.50	31.10	MP Mafic Intrusive Dark green, very fine grained, massive, hard, magnetic mafic intrusive. Very weak foliation at 60-65 degrees to CA. Sharp but irregular lower contact. Overall Tr. PY.							
31.10	51.40	VM Mafic Volcanic Medium to dark green, aphanitic to very fine grained, predominantly homogeneous appearing mafic volcanic. Massive to very weak foliation at 60-65 degrees to CA. Local but rare relict pillow selvages. <5% odd quartz/carb stringers. Overall Tr. PY.							
51.40	89.00	VMP; FBX Mafic Volcanic Pillowed; Flow Breccia General description as above at 4.5 to 51.4 but with increasing relict pillow selvages and flow breccia zones. Local zones of tension fractures with dark green-black chlorite infill. 5-8% odd quartz/carb stringers, patches and selvage replacement. Local minor epidote fracture fill. Weakly developed F1 at 65-70 degrees to CA. Overall Tr. to <.5% fracture fill and specks PY.							
89.00	90.80	MP Mafic Intrusive Light to medium green grey, fine grained, massive, hard, carb/quartz/carb speckled non-magnetic mafic intrusive. Sharp lower contact at 25 degrees to CA. Overall Tr. PY.							
90.80	350.00	VMP; FBX Mafic Volcanic Pillowed; Flow Breccia Generally as from 51.4 to 89.0m but with less pillow selvages and flow breccia zones to 144.0m. Thereafter to 207.1m pillow selvages and flow brecciation increase with locally abundant dark green to black chlorite tension fracture, brecciation and selvage infill. <5% torn, crenulated & irregular quartz/carb stringers. Very weak F1 at 65-70 degrees to CA. Overall Tr to <.5% PY. 231.3 to 242.5m Decreased pillow selvages and flow breccia-section is a more homogeneous mafic flow. 242.5 to 280.0m Predominantly flow breccia.	143.00	144.10	24887	1.10	<5		
			144.10	145.00	24888	0.90	7		
			145.00	146.00	24889	1.00	7		
			146.00	147.00	24891	1.00	<5		
			146.00	147.00	24890 (Dbl)	1.00	<5		
			147.00	148.00	24892	1.00	<5		
			148.00	149.00	24893	1.00	<5		
			149.00	150.00	24894	1.00	9		
			150.00	151.00	24895	1.00	<5	<5	
			151.00	152.00	24896	1.00	<5		

Brunswick Resources Inc.

Description	Assay						
	From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
280.0-298.2m Lighter green grey mafic flow possibly due to minor sericite bleaching.	152.00	153.00	24897	1.00	<5		
	153.00	154.00	24898	1.00	<5		
298.2 to 308.5m Medium grey green, harder, fine grained and possibly weakly silicified mafic flow.	154.00	155.00	24899	1.00	6		
	155.00	156.00	24900	1.00	<5		
308.5 to 348.0m Flow brecciated with local variolithic sections. 10-15% irregular quartz/carb veins/stringers and fractute fill from .5 to 2cm commonly at 0 to 25 degrees to CA.	156.00	157.00	24902	1.00	10		
	156.00	157.00	24901 (Bln)	1.00	5		
	157.00	158.00	24903	1.00	<5		
	158.00	159.00	24905	1.00	<5		
	158.00	159.00	24904 (Std)	1.00	5,678		5.83
	159.00	160.00	24906	1.00	<5		
	160.00	160.70	24907	0.70	6	<5	
	160.70	162.00	24908	1.30	<5		
	238.00	239.40	24909	1.40	<5		
	239.40	240.60	24910	1.20	14		
	240.60	241.60	24912	1.00	<5		
	240.60	241.60	24911 (Bln)	1.00	<5		
	248.00	249.00	24913	1.00	<5		
	249.00	250.00	24914	1.00	<5		
250.00	251.00	24915	1.00	<5			
251.00	252.00	24916	1.00	<5			
252.00	253.00	24917	1.00	<5			
253.00	254.00	24918	1.00	7			
254.00	255.00	24920	1.00	<5			
254.00	255.00	24919 (Dbl)	1.00	6	6		
255.00	256.00	24921	1.00	<5			
256.00	257.00	24922	1.00	40			
257.00	258.00	24923	1.00	5			
327.00	328.20	24925	1.20	50			
327.00	328.20	24924 (Std)	1.20	2,192		2.13	
328.20	329.20	24926	1.00	13			
329.20	330.20	24927	1.00	<5			
330.20	331.20	24928	1.00	<5			
331.20	332.20	24929	1.00	<5			

Brunswick Resources Inc.

Description	Assay						
	From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
	332.20	333.00	24930	0.80	30		
	333.00	334.00	24931	1.00	<5	<5	
	334.00	335.00	24932	1.00	<5		
	335.00	336.00	24933	1.00	<5		
	336.00	337.00	24934	1.00	<5		
	337.00	338.00	24935	1.00	<5		
	338.00	339.00	24936	1.00	<5		
350.00 End of DDH Number of samples: 44 Number of QAQC samples: 6 Total sampled length: 44.60							

Brunswick Resources Inc.

DDH: MU-11-10RL

Claims title: ABITIBI GOLD PROECT.

Section:

Township: Munro

Level:

Range:

Work place:

Gov. Road Core Shack

Drilled by: NPLH

Lot:

Described by: W.MacRae

From: 2011-06-06

Description date:

To: 2011-06-08

Collar

UTM

Azimuth: 200.00°

East 555,108.00

Dip: -50.00°

North 538,089.00

Length: 384.00 m

Elevation 314.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	18.00	198.90°	-49.80°	No	
Reflex-Easy Shot	69.00	199.80°	-49.60°	No	
Reflex-Easy Shot	120.00	201.00°	-49.30°	No	
Reflex-Easy Shot	171.00	200.40°	-47.90°	No	
Reflex-Easy Shot	222.00	202.10°	-46.80°	No	
Reflex-Easy Shot	273.00	202.40°	-46.40°	No	
Reflex-Easy Shot	324.00	204.40°	-45.40°	No	
Reflex-Easy Shot	381.00	205.90°	-45.10°	No	

Description

UTM, Nad 83, Zone 17. MU-11-10RL is a re-log of MU-11-10. Re-logged by Les Kovacs, Nov. 2014. One additional samples added. (010374)

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	3.00	CA Casing Casing/Overburden.							
3.00	253.30	VMP; FBX Mafic Volcanic Pillowed; Flow Breccia Mostly medium to dark green, massive and aphanitic to fine grained, weakly foliated with F1 at 50-60 degrees to CA. Predominantly pillowed flows with local selvages and flow breccia. Local hematitic slips. <5% carbonate as selvage infill and fine fractures. Locally weakly to moderately magnetic. Overall <.5% odd specks PY 3.0 to 17.4m Blocky and locally vuggy with hematitic slips due to footwall faulting. 17.4 to 18.0m 60cm Fault. Blocky with rubble. 181.0 to 182.0 1.0 meter Fault Zone. Mostly angular rubble suggesting a brittle fault. 182.6 to 183.0 40cm Fault Zone. Mostly angular rubble suggesting a brittle fault.	18.00	18.30	24937	0.30	<5		
			18.30	19.30	24938	1.00	6		
			109.30	110.30	24939	1.00	7		
			110.30	111.35	24940	1.05	9		
			111.35	111.60	24941	0.25	<5		
			111.60	112.70	24942	1.10	17		
			112.70	113.50	24943 (Std)	0.80	2,262		2.23
			112.70	113.70	24944	1.00	12		
			113.70	114.70	24945	1.00	<5		
			114.70	115.70	24946	1.00	9		
			115.70	116.70	24947	1.00	<5		
			116.70	117.70	24949	1.00	<5		
			116.70	117.70	24948 (Dbl)	1.00	<5		
			117.70	118.70	24950	1.00	6		
			118.70	119.70	24951	1.00	7		
			119.70	120.70	24952	1.00	7		
			120.70	121.10	24953	0.40	9		
			121.10	122.10	24955	1.00	10	8	
			121.10	122.10	24954 (Bln)	1.00	<5		
			122.10	123.00	24956	0.90	<5		
			133.00	134.15	24957	1.15	<5		
			134.15	134.50	24958	0.35	<5		
			134.50	135.50	24959	1.00	<5		
			135.50	136.50	24960	1.00	<5		
			136.50	137.50	24961	1.00	7		
			137.50	138.50	24962	1.00	6		
			138.50	139.50	24963	1.00	<5		
			139.50	140.50	24964	1.00	<5		
			140.50	141.50	24965	1.00	<5		
			141.50	142.50	24967	1.00	10	12	
			141.50	142.50	24966 (Std)	1.00	5,630		5.83

Brunswick Resources Inc.

Description	Assay						
	From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
	142.50	143.50	24968	1.00	5		
	143.50	144.50	24970	1.00	8		
	143.50	144.50	24969 (Dbl)	1.00	<5		
	144.50	145.50	24971	1.00	45		
	145.50	146.50	24973	1.00	<5		
	145.50	146.50	24972 (Bln)	1.00	8		
	146.50	147.50	24974	1.00	<5		
	147.50	148.50	24975	1.00	<5		
	148.50	149.50	24976	1.00	<5		
	149.50	150.50	24977	1.00	22		
	150.50	151.40	24978	0.90	<5		
	151.40	152.20	24979	0.80	<5	<5	
	152.20	153.00	24980	0.80	<5		
	192.00	193.10	24981	1.10	<5		
	193.10	194.00	24982	0.90	<5		
	194.00	195.00	24983	1.00	<5		
	195.00	196.00	24984	1.00	8		
	196.00	197.00	24986	1.00	<5		
	196.00	197.00	24985 (Bln)	1.00	<5		
	197.00	198.00	24987	1.00	<5		
	198.00	199.00	24988	1.00	<5		
	199.00	200.00	24989	1.00	<5		
	200.00	201.00	24991	1.00	<5	<5	
	200.00	201.00	24990 (Std)	1.00	3,848		3.87
	201.00	202.00	24992	1.00	<5		
	202.00	203.00	24993	1.00	<5		
	203.00	204.00	24994	1.00	<5		
	204.00	205.00	24995	1.00	<5		
	205.00	206.00	24997	1.00	<5		
	205.00	206.00	24996 (Dbl)	1.00	<5		
	206.00	207.00	24998	1.00	5		
	207.00	208.00	24999	1.00	<5		
	208.00	209.00	25000	1.00	<5		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
			209.00	210.00	25001	1.00	<5		
			210.00	211.00	25002	1.00	7		
253.30	259.80	MP Mafic Intrusive Medium green, very fine grained, hard, massive, very weakly magnetic with minor quartz/carb speckling. <5% odd random calcite stringers of mm scale at 45 to 90 degrees to CA. Sharp but irregular upper/lower contacts. Trace PY.							
259.80	263.30	VMP Mafic Volcanic Pillowed Pillowed and locally flow brecciated mafic volcanics as from 3 to 253.3m. Tr. PY.							
263.30	275.80	MP Mafic Intrusive Description as from 253.3 to 259.8m. Mafic intrusive with sharp but irregular upper/lower contacts. Tr. PY.							
275.80	290.00	VMP; FBX Mafic Volcanic Pillowed; Flow Breccia Description as from 259.8 to 263.3m Hard, massive, mafic pillowed flow with local flow breccia. Magnetic throughout. Local mafic intrusive inclusions. Overall <.5% PY.							
		281.4 to 281.7m 30cm section with 5-8% fine tension fracture fill PY.							
290.00	299.00	MP Mafic Intrusive Hard, very fine grained mafic intrusive with description as from 263.3 to 275.8m. Sharp but irregular upper/lower contact. Tr. PY.							
299.00	341.60	VMP; FBX Mafic Volcanic Pillowed; Flow Breccia Hard, massive, magnetic pillowed and flow brecciated mafic volcanic with description as from 3 to 253.3m. Up to 10% assimilated mafic intrusive. Tr. PY.							
341.60	361.65	MP Mafic Intrusive Description as from 290 to 299.0m. Local hematitic/carb slips. Sharp but irregular upper/lower contact. 5% quartz/carb speckling from .5 to 3mm throughout. Tr. Py.							
361.65	384.00	VMP Mafic Volcanic Pillowed Description as from 299.0 to 341.6m. Pillowed and lesser flow brecciated mafic volcanic. Hard, massive and locally magnetic. Relict selvages with							

Brunswick Resources Inc.

DDH: MU-11-11RL Claims title: ABITIBI GOLD PROJECT. Section:
 Township: Munro Level:
 Range: Work place: Gov. Road Coreshack
 Lot:
 Drilled by: NPLH
 Described by: W.MacRae From: 2011-06-09 Description date:
 To: 2011-06-11

Collar

Azimuth: 200.00°
 Dip: -50.00°
 Length: 354.00 m

UTM
 East 555,067.00
 North 5,381,838.00
 Elevation 314.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	30.00	194.80°	-49.90°	No	
Reflex-Easy Shot	81.00	195.20°	-49.50°	No	
Reflex-Easy Shot	132.00	195.70°	-49.20°	No	
Reflex-Easy Shot	183.00	196.30°	-49.00°	No	
Reflex-Easy Shot	234.00	197.40°	-48.60°	No	
Reflex-Easy Shot	285.00	197.70°	-48.50°	No	
Reflex-Easy Shot	336.00	198.10°	-48.30°	No	

Description

UTM Nad 83, Zone 17 Hole MU-11-11RL is a re-log of MU-11-11. Re-logged by Les Kovacs Nov. 2014. No additional samples added.

Core size: NQ Cemented: No Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	10.60	CA Casing Overburden	2.05	21.00	25012 (Dbl)	18.95	<5		
10.60	99.30	VMP Mafic Volcanic Pillowed Mostly medium to dark green, aphanitic to fine grained, hard, massive mafic volcanic pillowed. Local well preserved pillow selvages replaced by chlorite, quartz/carb and lesser biotite with rare local flow breccia. Very weak F1 at 30-40 degrees to CA. Local <1.0m mafic intrusives. Overall <.5% odd specks, fracture fill PY. 10.6 to 41.0m Lighter pale green possibly due to sericite/epidote bleaching.	12.00	13.00	25003	1.00	8	10	
			13.00	14.00	25004	1.00	6		
			14.00	15.00	25005	1.00	<5		
			15.00	16.20	25006	1.20	9		
			16.20	16.80	25007	0.60	7		
			16.80	17.40	25008	0.60	57		
			17.40	18.30	25009	0.90	38		
			18.30	19.05	25010	0.75	8		
			19.05	20.05	25011	1.00	43		
			20.05	21.00	25013	0.95	<5		
			21.00	22.00	25014	1.00	<5		
			39.00	40.00	25015	1.00	<5	<5	
			40.00	41.20	25016	1.20	<5		
			41.20	42.00	25017	0.80	<5		
			42.00	42.70	25018	0.70	<5		
			42.70	43.70	25020	1.00	5		
99.30	112.20	MP Mafic Intrusive Medium to dark green, very fine to fine grained, hard, massive mafic intrusive. Somewhat homogeneous appearing throughout. Sharp but irregular upper/lower contacts. Appears to be a very weak F1 at 40-50 degrees to CA. Overall Tr. PY.	42.70	43.70	25019 (Bln)	1.00	<5		
112.20	151.50	VMP Mafic Volcanic Pillowed Major description as from 10.6 to 99.3m-mafic volcanic-pillowed with very local flow breccia. Very weak F1 at 40-50 degrees to CA. Overall Tr to <.5% PY. 133.0 to 145.0m Lighter green bleached (chlorite depleted) section due to FAULT at 140.25 to 140.65m. 140.25 to 140.65m Fault Zone. 40cm fault with rubble, fissile material and local gouge. S1 at 60 degrees to CA.	136.00	137.10	25022	1.10	10		
			136.00	137.10	25021 (Std)	1.10	3,526		3.77
			137.10	138.00	25023	0.90	10		
			138.00	139.00	25024	1.00	<5		
			139.00	140.00	25025	1.00	19		
			140.00	141.00	25026	1.00	41		
			141.00	142.00	25027	1.00	62	57	
			142.00	143.00	25028	1.00	7		
			143.00	144.00	25029	1.00	13		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
151.50	157.10	FZ Fault Zone Unit is composed of 20% bleached but competent mafic volcanic core, 50% blocky core and rubble, 20% breccia and mylonite and 10% local gouge. S1 appears to be at 60 degrees to CA. Drillers note 2.0 meters of lost core.	144.00	145.00	25031	1.00	<5		
			144.00	145.00	25030 (Bin)	1.00	6		
157.10	255.40	VMP Mafic Volcanic Pillowed Description as from 112.2 to 151.5m. Predominantly pillowed mafic volcanics with a somewhat lesser volume of pillow selvages and more massive mafic flow with very local flow breccia. Very weakly developed F1 at 55-60 degrees to CA. Somewhat blocky from 157.1 to 180m due to faulting. Overall <.5% odd specks and fracture fill PY.							
		179.0 to 181.1m Fault Zone. Mostly blocky & rubble core with gouge from 179.7 to 181.0m.							
255.40	265.60	VM Mafic Volcanic Medium green grey, very fine grained, massive mafic flow. Very weak F1 at 55-60 degrees to CA. Overall Tr. PY.							
265.60	273.60	MP Mafic Intrusive Dark green to green black, fine grained, massive, hard, locally magnetic with local abundant leucoxene? speckling. Sharp but irregular upper/lower contacts. Overall Tr. PY.							
273.60	285.00	VM Mafic Volcanic General description as from 255.4 to 265.6m. Green grey, aphanitic to fine grained, massive, hard, mafic volcanic flow. Very weak F1 at 50-55 degrees to CA. Overall Tr. PY.							
		274.0 to 274.4m Fault Zone-brittle fault with angular rubble.							
285.00	298.20	VMP Mafic Volcanic Pillowed Description as from 157.1 to 255.4. Pillowed volcanic with very good relict selvages and lesser flow breccia. Odd and local quartz amygdules to 5mm. Overall <5% quartz/carb stringers most commonly from 0 to about 15 degrees to CA. Tr. to <.5% PY.							
298.20	324.40	VM	298.60	299.60	25032	1.00	12		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
324.40	354.00	Mafic Volcanic	299.60	300.30	25033	0.70	<5	12	2.13
		Somewhat lighter green grey mafic flow as from 273.6 to 285.0m. Local sections show fine tuffaceous texture and stratigraphy (304 to 310m) .	300.30	301.30	25034	1.00	<5		
		Minor local epidote-sericite bleaching. Weakly developed F1 at 55-60 degrees to CA. Overall Tr. PY.	301.30	302.30	25035	1.00	10		
			302.30	303.30	25036	1.00	30		
			303.30	303.60	25037	0.30	33		
			303.60	304.60	25038	1.00	10		
			304.60	305.60	25040	1.00	<5		
			304.60	305.60	25039 (Db1)	1.00	13		
			305.60	306.60	25041	1.00	<5		
			306.60	307.60	25042	1.00	26		
			307.60	308.60	25043	1.00	<5		
			308.60	309.30	25044	0.70	<5		
			353.00	354.00	25045 (Std)	1.00	2,012		
354.00	End of DDH Number of samples: 37 Number of QAQC samples: 6 Total sampled length: 34.40								

Brunswick Resources Inc.

DDH: MU-11-12RL

Claims title: ABITIBI GOLD PROJECT.

Section:

Township: Munro

Level:

Range:

Work place:

Gov. Road Core Shack

Drilled by: NPLH

Lot:

Described by: W.MacRae

From: 2011-06-11

Description date:

To: 2011-06-13

Collar

Azimuth: 20.00°
Dip: -60.00°
Length: 252.00 m

UTM

East	554,048.00
North	5,381,760.00
Elevation	316.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	51.00	16.50°	-60.00°	No	
Reflex-Easy Shot	102.00	17.10°	-60.20°	No	
Reflex-Easy Shot	151.00	17.20°	-60.40°	No	
Reflex-Easy Shot	201.00	16.70°	-58.50°	No	
Reflex-Easy Shot	252.00	16.10°	-55.30°	No	

Description

UTM, Nad 83, Zone17 MU-11-12RL is a re-log of MU11-12. Re-logged by Les Kovacs November 2014

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	5.20	CA Casing Overburden							
5.20	103.30	VM; TU3 Mafic Volcanic; Mafic Tuff Mostly medium to dark green, aphanitic to fine grained, massive, locally tuffaceous mafic volcanic. Very weak F1 from 50-55 degrees to CA. Locally blocky due to several brittle & ductile faults. Sharp but irregular lower contact. Overall Tr. to <.5% PY. 18.0 to 22.3m Blocky due to open fracturing from 0 to 15 degrees to CA. with calcite/carb slickensides. 22.3 to 25.5m Lighter grey green, harder, siliceous and tuffaceous. 25.5 to 27.0m Blocky due to open fracturing from 0 to 15 degrees to CA. 29.4 to 29.7m Fault-blocky brittle faulting at 15-20 degrees to CA. 31.5 to 32.1m Faulting-angular blocky core with hematitic slip surface due to brittle faulting. 32.1 to 64.8m Mostly homogeneous mafic flow. 64.8 to 74.8m Lighter grey green, harder, silicified and tuffaceous section. 74.8 to 101.1m Mostly medium green grey, fine grained mafic flow-blocky due to local brittle faulting. 80.3 to 81.2m Fault Zone-mostly fine angular rubble suggesting a brittle fault. 81.7 to 82.0m Fault Zone-mostly fine angular rubble suggesting a brittle fault. 82.0 to 99.8m Mostly medium grey green, fine grained mafic flow with local hematitic slips. 99.8 to 100.6m Fault Zone-fine angular rubble suggesting a brittle fault.	101.60	102.40	25046	0.80	8		
			102.40	103.30	25047	0.90	<5		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
103.30	111.00	FZ; BX Fault Zone; Breccia Fault zone with mafic volcanics. Unit is composed predominantly of 75% brecciated rock with 10% rubble and minor gouge from 105.4 to 106.1m Also composed of approx. 10-15% injected quartz/carb. An open fracture can be seen from 104.7 to 105.6 at 0 degrees to CA with rubble and gouge. Overall <.5% fine specks PY.	103.30	104.20	25048	0.90	<5	<5	
			104.20	105.00	25049	0.80	<5		
			105.00	106.00	25050	1.00	<5		
			106.00	107.00	25051	1.00	<5		
			107.00	108.00	25052	1.00	<5		
			108.00	109.00	25053	1.00	<5		
			109.00	110.00	25054	1.00	<5		
111.00	123.50	MP Mafic Intrusive Medium to dark grey green, fine grained with coarser quartz/carb phenocrysts from 1-5mm, hard, massive and somewhat silicified. Locally dioritic in appearance. Very weakly foliated-F1 at 60 degrees to CA. Overall Tr. PY.	111.00	112.00	25056	1.00	<5		
			112.00	113.00	25057	1.00	<5		
			113.00	114.00	25058	1.00	<5		
123.50	138.60	FBX Flow Breccia Medium green, aphanitic to fine grained, medium hard, very weakly foliated with F1 at 55-60 degrees to CA. Composed predominantly of poorly defined flow breccia and tension fractures with carb/calcite and chlorite infill. Overall Tr. to <.5% PY.							
138.60	155.60	TU3 Mafic Tuff Light to medium green grey, overall fine grained matrix with angular to sub-rounded quartz shards from 1-4mm. (mostly clast supported). Hard, massive with local weak F1 at 60 degrees to CA. 153 to 155.6 approx. 30% assimilated flow breccia as from 123.5 to 138.6m. Overall <.5% PY.	141.00	142.25	25060	1.25	6	11	3.87
			141.00	142.25	25059 (Std)	1.25	3,522		
			142.25	143.00	25061	0.75	<5		
			143.00	144.00	25062	1.00	<5		
155.60	171.30	FBX; TU3 Flow Breccia; Mafic Tuff Medium green grey, fine grained, mottled and disruptive appearance due to the blending of flow breccia and mafic tuff sections and the brecciation and fracturing due to proximity to faulting. Overall Tr. Py.	162.00	163.00	25063	1.00	9	11	
			163.00	164.00	25064	1.00	6		
			164.00	165.00	25066	1.00	<5		
			164.00	165.00	25065 (Dbl)	1.00	<5		
			165.00	166.00	25067	1.00	<5		
			166.00	167.00	25069	1.00	<5		
			166.00	167.00	25068 (Bln)	1.00	<5		
			169.10	170.70	25070	1.60	<5		
170.70	171.20	25071	0.50	<5					

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
171.30	172.90	FZ Fault Zone 1.6m Fault zone composed predominantly of 70% rubble, 10% gouge, 15% somewhat competent mafic tuff and 5% quartz injection. Overall Tr. to <.5% odd specks PY.	171.20	172.50	25072	1.30	6		
			172.50	174.00	25073	1.50	5		
172.90	192.60	FBX; TU3 Flow Breccia; Mafic Tuff General description as from 155.6 to 171.3. Unit is composed of approx 60% lighter green-yellow mustard colored beds of mafic tuff and approx 40% flow breccia all with a disruptive appearance due to proximity to faulting. Tuffaceous sections diminish after approx 187.0m. Unit is hard to very hard suggesting moderate silicification. Overall Tr. to <.5% odd specks PY.	174.00	175.00	25074	1.00	6		
			175.00	176.00	25075	1.00	12	9	
			176.00	177.00	25076	1.00	6		
			177.00	178.00	25078	1.00	7		
			177.00	178.00	25077 (Dbl)	1.00	6		
			178.00	179.00	25079	1.00	7		
			179.00	180.00	25080	1.00	11		
			180.00	181.00	25082	1.00	14		
			180.00	181.00	25081 (Bln)	1.00	<5		
			181.00	182.00	25083	1.00	6		
			182.00	183.00	25084	1.00	<5		
			183.00	184.00	25085	1.00	<5		
			184.00	185.00	25086	1.00	18		
			185.00	186.00	25087	1.00	<5	<5	
			186.00	187.50	25088	1.50	<5		
			187.50	189.00	25089	1.50	<5		
192.60	212.80	FBX; VMP Flow Breccia; Mafic Volcanic Pillowed Light green buff to pale olive green, aphanitic to very fine grained, hard to very hard, sericite & silica bleached flow breccia with local altered pillow selvages with quartz/carb/chlorite infill. Weak F1 at 45-50 degrees to CA. Overall Tr to <.5% PY.							
212.80	240.40	VM Mafic Volcanic Medium to dark green grey, aphanitic to fine grained, hard, massive to weak F1 at 50 degrees to CA. Very local faint flow breccia and possible pillow selvages. Most areas show homogeneous mafic flow. Overall Tr. PY.							
240.40	252.00	VMP Mafic Volcanic Pillowed Medium green grey, aphanitic to fine grained, massive to weakly foliated	251.00	252.00	25090 (Std)	1.00	5,702		5.93

Brunswick Resources Inc.

Description	Assay						
	From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
<p>with F1 from 50-60 degrees to CA. Composed predominantly of chlorite, carb altered pillow selvages and local flow breccia. Flow breccia and tension fractures with chlorite infill. <5% quartz/carb stringers at various angles to CA. Overall Tr to <.5%PY.</p> <p>252.0 EOH</p>							
<p>252.00 End of DDH Number of samples: 39 Number of QAQC samples: 6 Total sampled length: 40.30</p>							

Brunswick Resources Inc.

DDH: MU-11-13RL
 Claims title: ABITIBI GOLD PROJECT.
 Township: Munro
 Range:
 Lot:
 Drilled by: NPLH
 Described by: W.MacRae
 From: 2011-06-13
 To: 2011-06-15
 Section:
 Level:
 Work place: Gov. Road Coreshack
 Description date:

Collar

Azimuth: 20.00°
 Dip: -60.00°
 Length: 255.00 m

UTM

East	553,946.00
North	5,381,746.00
Elevation	313.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	51.00	15.40°	-59.90°	No	
Reflex-Easy Shot	102.00	16.50°	-59.30°	No	
Reflex-Easy Shot	153.00	17.30°	-58.80°	No	
Reflex-Easy Shot	204.00	17.50°	-58.50°	No	

Description

UTM, Nad 83, Zone17 MU-11-13RL is a re-log of MU-11-13. Re-logged by Les Kovacs November 2014. No additional samples added.

Core size: NQ Cemented: No Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	4.20	CA Casing Overburden							
4.20	22.30	FBX Flow Breccia Medium green, aphanitic to fine grained, hard, massive to very poorly foliated at 60 degrees to CA. Composed predominantly of prolific flow breccia sections with rare pillow selvages with mostly chlorite and lesser epidote alteration. Overall Tr PY.	4.60	6.00	25091	1.40	10		
			6.00	7.00	25092	1.00	9		
			7.00	8.00	25093	1.00	<5		
			8.00	9.00	25094	1.00	5		
			9.00	10.00	25095	1.00	5		
			10.00	11.00	25096	1.00	17		
			11.00	12.00	25097	1.00	6		
			12.00	13.00	25098	1.00	12		
			13.00	14.00	25099	1.00	<5	<5	
			14.00	15.00	25100	1.00	<5		
			15.00	16.00	25101	1.00	13		
			16.00	17.00	25102	1.00	15		
			17.00	18.00	25103	1.00	<5		
			18.00	19.00	25104	1.00	<5		
			19.00	20.00	25105	1.00	5		
			20.00	21.00	25107	1.00	<5		
			20.00	21.00	25106 (Bln)	1.00	7		
			21.00	22.00	25108	1.00	6		
22.30	68.20	VM; FBX Mafic Volcanic; Flow Breccia General description as from 4.2 to 22.3m. Unit is composed mainly of aphanitic mafic flow with 15-20% local flow breccia of which has been infilled by chlorite and lesser quartz/cab. Very poorly foliated with F1 being 55-60 degrees to CA. Rare pillow selvages. Increasing mafic intrusive after 62.5m. Overall Tr to <.5% fine PY.	22.00	23.00	25109	1.00	<5		
			23.00	24.00	25110	1.00	<5		
			31.00	32.00	25111	1.00	10	12	
			32.00	32.50	25112	0.50	17		
			32.50	33.50	25113	1.00	7		
68.20	82.30	MP Mafic Intrusive Medium to dark green, fine to medium grained, harder, massive, with local gabbroic texture. Tension fractures with chlorite infill. Sharp but irregular upper/lower contacts. Overall Tr. PY.							
82.30	100.00	FBX Flow Breccia General description as from 4.2 to 22.3m. Flow breccia with dark green to	86.00	87.00	25115	1.00	9		
			86.00	87.00	25114 (Std)	1.00	2,128		2.16

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
100.00	211.80	<p>MP Mafic Intrusive General description as from 68.2 to 82.3m but with approx. 20% assimilated mafic flow and flow breccia from 100 to 130m. Unit shows increased tension fracturing over intrusive as above with dark green to black chlorite infill. 5% white local quartz/carb veins from 3 to 40cm. Local epidote & hematite slips. Locally coarser with near gabbroic texture. Overall Tr. PY.</p> <p>153 to 153.4m 40cm white, barren quartz/carb vein with 20% wallrock material and sharp but irregular upper/lower contacts.</p> <p>158.1 to 158.5m 40cm barren white quartz/carb vein as from 153 to 153.4m.</p>	87.00	88.00	25116	1.00	<5	<5	
			88.00	89.00	25118	1.00	<5		
			88.00	89.00	25117 (Dbl)	1.00	<5		
			89.00	90.00	25119	1.00	<5		
			90.00	91.00	25120	1.00	<5		
			91.00	92.00	25121	1.00	<5		
			92.00	93.00	25122	1.00	<5		
			93.00	94.00	25123	1.00	<5		
			94.00	95.00	25125	1.00	<5		
			94.00	95.00	25124 (Dbl)	1.00	<5		
			95.00	96.00	25126	1.00	<5		
			96.00	97.00	25127	1.00	<5		
			97.00	98.00	25128	1.00	<5		
			98.00	99.00	25130	1.00	<5		
			98.00	99.00	25129 (Bln)	1.00	<5		
99.00	100.00	25131	1.00	<5					
211.80	255.00	<p>VMP Mafic Volcanic Pillowed Medium to dark green, aphanitic to fine grained, hard, massive to weakly foliated at 60 degrees to CA. Mafic pillowed volcanic with abundant well preserved selvages as well as local but less common flow breccia. <.5% odd irregular quartz/carb stringers/veins. Overall Tr to <.5% fine PY.</p>							

255.00 End of DDH
Number of samples: 36
Number of QAQC samples: 5
Total sampled length: 35.90

Brunswick Resources Inc.

DDH: MU-11-14RL

Claims title: ABITIBI GOLD PROJECT.

Section:

Township: Munro

Level:

Range:

Work place:

Gov. Road Coreshack

Drilled by: NPLH

Lot:

Described by: W.MacRae

From: 2011-06-15

Description date:

To: 2011-06-17

Collar

UTM

Azimuth: 90.00°

East 553,910.00

Dip: -50.00°

North 5,381,797.00

Length: 357.00 m

Elevation 309.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	51.00	90.50°	-49.50°	No	
Reflex-Easy Shot	102.00	90.70°	-49.40°	No	
Reflex-Easy Shot	153.00	101.10°	-49.00°	No	
Reflex-Easy Shot	204.00	91.40°	-48.80°	No	
Reflex-Easy Shot	255.00	103.00°	-48.60°	No	
Reflex-Easy Shot	306.00	104.20°	-48.40°	No	
Reflex-Easy Shot	357.00	96.10°	-48.10°	No	

Description

UTM, Nad 83, Zone17 MU-11-14RL is a re-log of MU-11-14. Re-logged by Les Kovacs November 2014

Core size:

NQ

Cemented: No

Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	3.00	CA Casing Overburden							
3.00	21.30	FBX Flow Breccia Medium to dark green grey, aphanitic to fine grained, massive to very weakly foliated at 55-60 degrees to CA. Unit is composed of abundant section of flow breccia and tension fractures with chlorite infill. Overall Tr. PY. 3.6 to 4.0 40cm Fault Zone-brittle faulting with angular rubble. 12.0 to 12.4m 40cm Fault Zone-brittle faulting with angular rubble. 21.0 to 21.1m 1.0cm Fault Zone-brittle faulting with rubble and a ankeritic/carb halo in hanging/footwall of fault.	6.50 7.50 8.50 8.50 9.50 17.00 18.00 19.00 19.50 20.00	7.50 8.50 9.50 9.50 10.50 18.00 19.00 19.50 20.00 21.30	25132 25133 25135 25134 (Std) 25136 25137 25138 25139 25140 25141	1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.50 0.50 1.30	9 <5 9 3,564 <5 <5 <5 9 7 6	6	3.70
21.30	65.90	MP Mafic Intrusive Medium to dark green grey, fine grained with local coarser sections, massive to very weakly foliated at 50-60 degrees to CA. Very locally magnetic with locally abundant tension fractures with chlorite infill. <10% local quartz/carb stringers, veins and patches. Overall Tr. PY.							
65.90	90.40	VM Mafic Volcanic Medium green, aphanitic to fine grained, massive to weakly foliated at 60 degrees to CA. mafic flow. Local lighter green epidotitic fracture fill. Local hematitic slips. Rare pillow selvages. Overall Tr. PY.	90.00	91.00	25142	1.00	<5		
90.40	102.00	FBX Flow Breccia Mafic flow breccia with description as from 3 to 21.3m. 5% local irregular quartz/carb patches with 1-3% anhedral PY accumulations confined to flow brx. Overall .5% fine specks and Bx/fracture fill PY.	91.00 92.00 93.00 94.00 95.00 96.00 97.00 98.00 99.00 99.70 99.70 101.00	92.00 93.00 94.00 95.00 96.00 97.00 98.00 99.00 99.70 101.00 101.00 102.00	25143 25144 25145 25146 25147 25148 25149 25150 25151 25153 25152 (DbI) 25154	1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.70 1.30 1.30 1.00	8 <5 6 <5 <5 <5 <5 <5 <5 <5 <5 <5	<5	

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
102.00	116.40	VM Mafic Volcanic Mafic flow as described from 65.9 to 90.4m. Minor tension fracturing with chlorite infill. Very weak F1 at 50-55 degrees to CA. Overall <.5% PY.	115.00	116.00	25155	1.00	<5		
			116.00	117.00	25156	1.00	11		
116.40	127.10	FBX Flow Breccia Predominantly flow breccia as from 90.4 to 102.0m. Very weakly foliated at 50-60 degrees to CA. Overall <.5% fine PY.	117.00	118.00	25157	1.00	<5		
			118.00	119.00	25158	1.00	<5		
			119.00	120.00	25159	1.00	<5	<5	
			120.00	121.00	25161	1.00	<5		
			120.00	121.00	25160 (Bln)	1.00	<5		
			121.00	122.00	25162	1.00	<5		
			122.00	123.00	25163	1.00	<5		
			122.00	123.00	25164 (Std)	1.00	2,236		2.23
127.10	181.25	MP Mafic Intrusive Mafic Intrusive with general description as from 21.3 to 65.9m. Overall Medium to dark green, fine grained, massive, hard, locally coarser and appearing gabbroic. Locally magnetic and with minor local tension with chlorite infill. Local irregular quartz/carb to 40% mostly from 144.2 to 144.8m. Very weak F1 at 60 degrees to CA. Overall Tr. PY.							
181.25	330.30	VM; VMP Mafic Volcanic; Mafic Volcanic Pillowed Generally as from 102 to 116.4. Somewhat lighter pale green where weakly sericite bleached within local flow breccia. Rare pillow selvages to 204.0m then selvages become more numerous and better defined. Selvages composed of chlorite, epidote, sericite and lesser quartz/carb. Overall Tr to <.5% PY. 189.6 to 191.2m 1.6m section of mafic intrusive. Sharp but irregular upper/lower contacts. 244.0 to 244.65m 65cm dark grey, fine grained, massive, hard, diabase dike inclusion with sharp but irregular upper/lower contacts. 244.65 to 306.5m Mostly medium green grey, aphanitic to fine grained, weakly foliated with F1 at 55-60 degrees to CA. Predominantly mafic flow with poorly defined flow breccia and rare pillow selvages. 297.65 to 297.9-25cm diabase dike with sharp upper/lower contacts at 40 degrees to CA. Tension fractures with chlorite infill. Overall Tr to <.5% PY.	188.60	189.60	25165 (Std)	1.00	7		
			189.60	190.30	25166	0.70	<5		
			190.30	191.00	25167	0.70	<5		
			191.00	192.00	25168	1.00	<5		
			192.00	193.00	25169	1.00	<5		
			193.00	193.80	25170	0.80	<5		
			193.80	195.00	25171	1.20	<5	<5	
330.30	335.40	MP7 Diabase							

Brunswick Resources Inc.

Description		Assay					
		From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)
335.40	339.30	<p>Dark grey to black, aphanitic to fine grained, massive, magnetic diabase dike. Sharp upper/lower contacts at 40 degrees to CA. Locally blocky. Tr. PY at margins.</p> <p>VM; VMP Mafic Volcanic; Mafic Volcanic Pillowed Mafic volcanic with local pillow selvages and coarser flow as from 181.25 to 330.3 before interruption by diabase dike. Overall Tr. PY Minor local quartz/carb stringers/veins.</p> <p>335.6 to 335.7m 10cm fault zone with rubble.</p>					
339.30	340.20	<p>336.2 to 336.45m 25cm brittle fault zone with angular rubble.</p> <p>FZ Fault Zone 90cm brittle fault zone with 100% angular rubble. Tr. PY.</p>					
340.20	357.00	<p>VM Mafic Volcanic Medium to dark green, aphanitic to fine grained, massive to weakly foliated mafic flow. Very weak F1 at 45-50 degrees to CA. Tension fractures with dark green chlorite infill. Lighter green olive green section suggest sericite/epidote bleaching. Overall Tr to <.5% PY.</p> <p>346.5 to 346.8m 30cm brittle fault zone with angular rubble.</p>					
357.00	<p>End of DDH Number of samples: 35 Number of QAQC samples: 5 Total sampled length: 33.70</p>						

Brunswick Resources Inc.

DDH: MU-11-15RL
 Claims title: ABITIBI GOLD PROJECT.
 Township: Munro
 Range:
 Level:
 Work place: Gov. Road Coeshack
 Drilled by: NPLH
 Lot:
 Described by: W.MacRae
 From: 2011-06-18
 Description date:
 To: 2011-06-20

Collar

Azimuth: 255.00°		UTM	
Dip: -45.00°		East	554,127.00
Length: 351.00 m		North	5,381,842.00
		Elevation	312.00

Down hole survey

Type	Depth	Azimuth	Dip	Invalid	Description
Reflex-Easy Shot	18.00	247.10°	-44.20°	No	
Reflex-Easy Shot	69.00	247.00°	-43.90°	No	
Reflex-Easy Shot	120.00	246.90°	-43.70°	No	
Reflex-Easy Shot	171.00	246.10°	-43.40°	No	
Reflex-Easy Shot	222.00	248.10°	-43.30°	No	
Reflex-Easy Shot	273.00	249.60°	-43.20°	No	

Description

UTM, Nad 83, Zone17 MU-11-15RL is a re-log of MU-11-15. Re-logged by Les Kovacs November 2014

Core size: NQ Cemented: No Stored: Yes

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
0.00	4.10	CA Casing Overburden							
4.10	7.00	FZ Fault Zone 2.9m brittle fault zone with light green pale beige aphanitic to fine grained, sericite? bleached mafic volcanic. Rubble from 4.9 to 5.3m. rest of unit is bleached and blocky. Overall Tr. to <.5% PY.	6.00	7.10	25172	1.10	16		
7.00	11.50	QZ/CA; BX; FZ Quartz/Carb Veining; Breccia; Fault Zone 60% ragged quartz/carb stringers, veins, patches with altered wallrock material all becciated due to hangingwall fault at 4.1 to 7.0m. Competant core is pale beige to buff and is most likely a bleached mafic volcanic. 11.4 to 11.5-fine angular rubble-brittle fault zone. Overall Tr to <.5% PY.	7.10	7.60	25173	0.50	192		
			7.60	8.20	25175	0.60	108		
			7.60	8.20	25174 (Dbl)	0.60	198		
			8.20	9.00	25176	0.80	42		
			9.00	9.50	25177	0.50	73		
			9.50	10.00	25178	0.50	421		
			10.00	10.50	25179	0.50	185		
			10.50	11.00	25180	0.50	154		
			11.00	11.70	25182	0.70	774		
11.50	40.50	VM; SER Mafic Volcanic; Sericite Mostly light to medium pale olive green to buff green, aphanitic to very fine grained, massive to very weakly foliated at 50-55 degrees to CA. Unit is most likely a sericite bleached mafic volcanic due to proximity to hangingwall fault. Bleached diminishes downhole. Abundant micro and tension fractures with silica and chlorite infill at various angles to CA. Overall Tr. PY.	11.00	11.70	25181 (Bln)	0.70	5		
			11.70	12.80	25183	1.10	461	442	
			12.80	14.00	25184	1.20	25		
			14.00	15.00	25185	1.00	11		
			15.00	16.00	25187	1.00	10		
			15.00	16.00	25186 (Std)	1.00	5,668		5.83
			16.00	17.00	25188	1.00	7		
			17.00	18.00	25189	1.00	5		
40.50	74.05	VM; FBX Mafic Volcanic; Flow Breccia Medium to dark green, aphanitic to fine grained, weakly foliated at 55 degrees to CA. Predominantly mafic flow with local poorly preserved flow breccia. Locally blocky due to weak faulting. Overall Tr PY. 40.5 to 40.9m 40cm brittle faulting. 59.2 to 66.5m Somewhat blocky section with abundant slips at 0 to 10 degrees to CA locally with minor gouge. 69.1 to 69.5m 40cm medium grained QFP/syenite inclusion.	68.00	69.00	25190	1.00	19		
			69.00	69.70	25191	0.70	<5		
			69.70	70.70	25192	1.00	<5		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
74.05	181.80	MP Mafic Intrusive Dark green, fine to medium grained, massive, locally magnetic mafic dike with local gabbroic phases. Abundant slips with hematite slickensides. Very weak F1 at 55-60 degrees to CA. Up to 5% local quartz/carb stringers/veins with wallrock material. Locally blocky with no evidence of faulting. Overall Tr. PY. 129.0 to 181.8 Increased tension fracturing with dark green to black chlorite infill.							
181.80	209.90	VM Mafic Volcanic Medium green, aphanitic to fine grained, massive mafic volcanic flow. Locally blocky with no evidence of faulting. Tension fractures with chlorite infill and local but poorly defined flow breccia and pillow selvages not common. 5% quartz/carb and calcite stringers and veinlets most commonly at 50-60 degrees to CA parallel to sub-parallel very weak F1. Overall Tr. Py.							
209.90	245.50	FBX Flow Breccia Medium green, aphanitic to fine grained, massive and brecciated appearing flow breccia. Unit is composed of well preserved and defined flow breccia sections with dark green to black chlorite & quartz/carb breccia infill. Rare pillow selvages. Very weak F1 at 60 degrees to CA. hematitic slips are common. Tr to .5% anhedral PY and pyrrhotite mainly within flow breccia.	210.00	211.00	25193	1.00	<5		
			211.00	212.00	25194	1.00	10		
			212.00	213.00	25195	1.00	<5	<5	
			213.00	214.00	25196	1.00	9		
			214.00	215.00	25197	1.00	<5		
			215.00	216.00	25198	1.00	<5		
			216.00	217.20	25199	1.20	<5		
245.50	254.20	VM Mafic Volcanic Description very much like 181.8 to 209.9m. Medium green, aphanitic to fine grained, massive mafic homogeneous flow with <5% local odd flow BX and lesser rare pillow selvages. Very weakly foliated with F1 at 60 degrees to CA. Overall Tr. Py.							
254.20	288.40	FBX Flow Breccia Description as from 209.9 to 245.5. Predominantly flow breccia (80%) with chlorite and lesser quartz/carb infill. Tr to <.5% fine PY and pyrrhotite. Very weak F1 at 60 degrees to CA.	254.20	255.00	25200	0.80	<5		
			255.00	256.00	25201	1.00	<5		
			256.00	257.00	25202	1.00	<5		
			257.00	258.00	25203	1.00	<5		
			258.00	259.00	25204	1.00	<5		
			259.00	260.00	25205	1.00	<5		
			260.00	261.00	25206	1.00	<5		

Brunswick Resources Inc.

Description			Assay						
			From	To	Sample number	Length	Au (ppb)	Au-Dup (ppb)	Au (g/t)
288.40	298.50	VM Mafic Volcanic Description as from 245.5 to 254.2m. Homogeneous mafic interflow between flow breccia units. Medium green, aphanitic to fine grained with very weak F1 at 60 degrees to CA. Overall Tr. PY.	261.00	262.00	25208	1.00	<5	<5	2.23
			261.00	262.00	25207 (Dbl)	1.00	<5		
			262.00	263.00	25209	1.00	<5		
			263.00	264.00	25210	1.00	<5		
			298.00	299.00	25212	1.00	5		
			298.00	299.00	25211 (Std)	1.00	2,018		
298.50	331.10	FBX Flow Breccia Predominantly (70%) well preserved and defined flow breccia with mostly chlorite and lesser quartz/carb infill. Very weak F1 at 60 degrees to CA. Overall Tr to <.5% fine PY and Tr. pyrrhotite. 306.4 to 319.2m Somewhat weaker flow brecciation. 319.2 to 320.0m 80cm ductile fault zone with 50% blocky rubble core and 50% gouge and breccia. Possible S1 at 45 degrees to CA. Tr. PY. 320.0 to 331.1 Flow breccia with much decreased brecciation and increasing mafic intrusive inclusions to end of unit.	299.00	300.00	25213	1.00	<5		
			300.00	301.20	25214	1.20	<5		
			301.20	301.70	25215	0.50	<5		
			301.70	302.40	25217	0.70	<5		
			301.70	302.40	25216 (Bln)	0.70	<5		
			302.40	303.40	25218	1.00	<5		
331.10	351.00	MP Mafic Intrusive Medium to dark green, fine to locally medium grained, massive, hard, magnetic mafic intrusive with local gabbroic phases. 5% quartz/carb veining. Very weak F1 at 60 degrees to CA. Overall <.5% PY but up to 2-3% very locally as anhedral clumps to 1-3mm. Tr pyrrhotite. 339.35 to 339.9m 55cm quartz/carb vein with wallrock material with sharp upper/lower contacts at 50 degrees to CA. Tr. PY at margins. 346.8 to 347.1m 30cm quartz/carb vein with minor wallrock material with sharp upper/lower contacts at 50 degrees to CA with Tr PY at margins.							
351.00	End of DDH Number of samples: 41 Number of QAQC samples: 6 Total sampled length: 37.10								

