We are committed to providing <u>accessible customer service</u>. If you need accessible formats or communications supports, please <u>contact us</u>.

Nous tenons à améliorer <u>l'accessibilité des services à la clientèle</u>. Si vous avez besoin de formats accessibles ou d'aide à la communication, veuillez <u>nous contacter</u>.



DIAMOND DRILLING BEILBY, McBRIDE AND CLARKE on the SHINING TREE PROJECT for PLATINEX INC.

CHURCHILL AND ASQUITH TOWNSHIPS, LARDER LAKE MINING DIVISION ONTARIO, CANADA

Prepared By: Iain Trusler Under the advisement of James R. Trusler P.Eng, President PLATINEX INC. March 7th, 2017

INDEX

Section	No.	Title	Page No.
1.0		Introduction	1
2.0		Property Location and Access	1
3.0		Previous Work	2
4.0		Topography	2
5.0		Geology	3
	5.1	Quaternary Geology	3
	5.2	General Bedrock Geology	3
	5.3	Metamorphism and Structure	4
6.0		Drilling Program	4
	6.1	Logistics	4
	6.2	Results	5
7.0		Conclusions and Recommendations	6
8.0		References	7
	Quali	ifications	9

LIST OF FIGURES

Figure No.	Title	Page No.
1	Location and Contiguity Map	11
2-5	Drill Hole Plan Maps	12 - 15
6-10	Drill Hole Cross Sections	16 - 20

Appendix I	Drill Hole Logs	21
Appendix II	Certificates of Analysis	53

1.0 INTRODUCTION

This report presents the results of five reconnaissance drill holes on the Beilby, McBride, and Clarke areas of the Platinex Inc. Shining Tree Project. The project is located in Churchill, Asquith and Macmurchy townships, Larder Lake Mining Division, District of Sudbury, Ontario. The work was completed between March 16th 2012 and November 24th 2016 as part of ongoing exploration by Platinex Inc. Over the course of two years prior to drilling, several surveys were carried out to explore for gold using glacial till. Prospecting of gold in till anomalies, powerstripping and channel sampling during the summer of 2011 in the Beilby area located a new gold occurrence. An IP/Resistivity survey was completed by JVX Ltd. by March 11 of 2012 covering the new gold occurrence as well as several other areas with high gold in till results.

Targets were selected on four separate claims; three of the holes were centered on targets south of Beilby Lake. Two of them on claim L4207971, and one on claim L4245864 which at depth crosses onto claim L1217520. Another hole was situated about a kilometre east of the first three holes and about a half of a kilometre north of highway 560 on claim L1192177 dubbed the McBride area. The last hole was located around three kilometres south of the others and just north of Macdonald Lake on claim L4209215, on what is known as the Clarke showing.

Diamond drilling was performed by Laframboise Drilling, of Earlton, Ontario under supervision from Platinex field geologist Dean Cutting P.Geo. Drill Core from all holes (BP12-01 through BP12-03, MP12-01, and CKP12-01) was logged, sawn and stored on property rented by Platinex from Bruce Cupskey in Shining Tree, Ontario. Twenty three samples were taken to Accurassay Laboratories of Thunder Bay, Ontario and 197 samples were taken to SGS minerals of Toronto, with sample prep completed at SGS Sudbury, Ontario for gold analysis.

2.0 PROPERTY LOCATION AND ACCESS

At the time that this drill program was completed the Shining Tree property consisted of 148 contiguous claim units and one separate claim containing 3 claim units in Churchill, Asquith and MacMurchy townships, Larder Lake Mining Division, District of Sudbury, Ontario. Of the claim units, 139 were and are subject to an underlying agreement with Skead Holdings Ltd. The remaining 12 were but are not currently under option agreement with Canadian Prospecting Ventures Inc.

Figure 1 shows the location of the Shining Tree area in Ontario as well as the claim locations and numbers with respect to major topographic and cultural features of the area.

Primary access to the property is obtained using a paved secondary highway (560) which runs through the centre of the property. Highway 560 connects with Highway 144 to the west and with Highway 65 at Elk Lake to the east. The claims are about 3 kilometers northeast of the village of Shining Tree and about 50 kilometers west of Gowganda. A number of logging trails accessible by 4-wheel drive vehicle provide access to portions of the property, and boat access is possible using Michiwakenda Lake, Cryderman Lake and West Shining Tree Creek.

Access to the Beilby Lake area was obtained via an old logging road and by cutting drill trails to allow the drill rig to gain access to the collar sites. The entrance of the road is located north of the Spruce Shilling Camp on Highway 560, on the west side. Access to the McBride area was similarly obtained by another old logging road and by cutting a drill trail to allow the drill rig to gain access. The entrance of the road is located on the north side of Highway 560, on claim L1199655. Access to the Clarke area was obtained by an old logging road on the east side of Highway 560 just north of Moorecamp Lake.

3.0 PREVIOUS WORK

Several shafts with limited underground development are situated on the project claims, and existed within separate properties pre 1940's. These were best known as the Herrick, Churchill, and Caswell properties. Relatively little diamond drilling has been done on the Churchill, and only sporadic programs have been carried out on the Caswell, including seven holes drilled by Platinex in 2011. Within the last eight years 51 drill holes have targeted the Herrick deposit, bringing the total number of diamond drill holes to 66. Exploration on the remainder of the property has been limited to prospecting, hand dug trenches, mapping and local sporadic geophysical and diamond drilling programs. Several glacial till sampling programs have also been carried out.

4.0 TOPOGRAPHY

The area has relatively low relief between 350 and 420 metres above sea level. Terrain is hummocky and gently rolling, with the remnant bases of Nipissing diabase sills forming several of the higher ridge lines, along with positive relief Matachewan diabase dykes. The area is generally well drained with numerous lakes and rivers. Logging for pine, spruce and poplar has taken place in small areas of the property at various times in the past, and continues. Regrowth is generally jackpine and poplar. Cedar is common in poorly drained areas. Outcrop ranges from 5% to greater than 50% in some areas, with outwash sands obscuring the eastern-most part of the property.

5.0 GEOLOGY

5.1 QUATERNARY GEOLOGY

The glacial deposits preserved in the area are products of the latest continental ice sheet, the Laurentide of Wisconsinan age. The Keewatin lobe advanced from the northeast approximately 100,000 years ago, and extended south into the northern United States. By 11,000 years ago, the ice sheet had receded back to the Shining Tree area and deposited a variety of surficial material, dominated by thin sandy till ground moraine over bedrock knobs (Roed and Hallett, 1979). Sand and gravel outwash deposits begin to predominate on the eastern edge of the project area, and can often be found as a thin deposit overlying ground moraine tills.

5.2 GENERAL BEDROCK GEOLOGY

The Shining Tree greenstone belt is located approximately 100 km north of Sudbury, and is located within the southern portion of the Abitibi Sub province, Superior Province, northeast Ontario. The supracrustal rocks in the Shining Tree area have been divided into the Pacaud, Deloro, Kidd-Munro, Tisdale and Timiskaming assemblages in keeping with the rest of the Abitibi greenstone belt (Ayer 1999; Ayer et al. 1999; Johns 1999b; Oliver et al. 1999b). The ~2669-2678 Ma Timiskaming assemblage is separated from the older assemblages (>2.7 Ga) by an unconformity. The Timiskaming assemblage is also composed of a considerably different array of rocks than the older supracrustal rocks (Ayer 2000).

The Pacaud, Deloro, Kidd-Munro and Tisdale assemblages are dominated by volcanic supracrustal rocks, which were formed before the first phase of deformation. Felsic volcanic units close to the presumed tops of the assemblages in the Shining Tree area have been dated: The ages of the older three assemblages (Pacaud, Deloro and Kidd-Munro) indicate that the greenstone belt is younging to the northeast (Ayer 2000).

The Pacaud assemblage is mainly composed of massive and pillowed basalts and is associated with minor spinifex or cumulate textured komatiites.

The Deloro assemblage is dominated by felsic volcanic rocks and is capped in many places by chemical meta-sedimentary rocks, seen as banded chert and jasper.

The Kidd-Munro assemblage is a varied assemblage dominated by tholeiitic basalts and komatiites, with minor felsic volcanic rocks, and the Tisdale assemblage comprises mafic flows and intermediate to felsic pyroclastics and/or volcaniclastics (Johns 1999a).

The 2012 drilling program intersected rock units from both the Pacaud assemblage and the Deloro assemblage. It is suggested that drill holes BP12-01 and BP12-02, and BP12-03 have intersected volcanic rocks in close proximity to the Pacaud-Deloro boundary, possibly with mafic volcanic rocks belonging to the Pacaud assemblage and felsic volcanic rocks belonging to the Deloro assemblage. MP12-01 is entirely composed of a low-quartz intrusive (syentite?) of unknown age. CKP12-01 was drilled entirely with Pacaud assemblage mafic to ultramafic lithologies.

5.3 METAMORPHISM AND STRUCTURE

The metamorphic grade throughout most of the Shining Tree area is mid to low greenschist facies (Oliver et al. 1999a, 1999b). Amygdules are filled with chlorite, carbonate or quartz. There are two main phases of deformation and associated metamorphism in the Shining Tree area (Oliver et al. 1999a, 1999b) with rocks older than 2.7 Ga having undergone two periods of deformation. There are multiple deformation zones in the older volcanic rocks in which gold has been found, especially in MacMurchy and Tyrrell Townships (Johns 1996, 1997 and 1999a). The Timiskaming assemblage has undergone a single period of deformation and is metamorphosed to a lesser degree than the older volcanic rocks (Oliver et al. 1999a, 1999b). The Timiskaming assemblage was formed between the two deformation events and lies unconformably above the pre-deformational volcanics (Ayers 2000).

6.0 DRILLING PROGRAM

6.1 LOGISTICS

Diamond drilling was done under contract by Laframboise Drilling, of Earlton, Ontario from March 22th until March 29th 2012. The project was conducted under the supervision of Dean Cutting P.Geo., of Rouyn-Noranda, Quebec, from mid-March 2012 through mid-November 2016. Drill core was sawn in preparation for sampling, and stored on property rented by Platinex from Bruce Cupskey in Shining Tree, Ontario. Preliminary logging and sampling of the core was completed in the early spring of 2012 by Dean Cutting with the assistance of Bruce Cupskey. Relogging and additional sampling work was

completed in the fall of 2016 by Dean Cutting, with core being cut by Canadian Exploration Services in Larder Lake, Ontario. Preliminary drill logs from 2012 as well as final drill logs from 2016 have been included in this report.

The NQ drill core was sawn in half, with one half archived and the other sent for gold analysis at SGS Minerals of Toronto, Ontario for the 2012 sampling, and Accurassay Laboratories of Thunder Bay, Ontario for the 2016 sampling. Standards, and blank samples were inserted into the sample stream at the core facility roughly every 20 samples.

6.2 RESULTS

The Drilling program comprised of 5 holes on the properties totaling 869 metres. The holes were drilled to test various IP anomalies coincident to gold in till anomalies. Plan views of the drilling are illustrated in figures 2 to 5 (pages 12-15). Drill cross sections are presented in figures 6 to 10 (pages 16-20)

BP12-01 (See figure 6) This drill hole tested a strong surface chargeability anomaly. Rock types logged in the hole consist of mafic volcanics, either massive or pillowed. Chargeability appears to be caused by sporadic to locally strong pyrite mineralization controlled by quartz-carbonate veinlets. The highest gold value was 140 ppb associated with 15 to 20% disseminated pyrite.

BP12-02 (See figure 7) This hole tested a strong surface chargeability anomaly. Rock types logged in the hole range from felsic volcanics to mafic volcanics as well as gabbroic intrusive units. Several late diabase dykes were also intersected. The chargeability anomaly appears to be caused by wide zones of disseminated pyrite in quartz-carbonate-sericite alteration within felsic volcanic rocks. Most gold values however were below 5 ppb, with a high value of 65 ppb.

BP12-03 (See figure 8) This hole tested a weak surface chargeability anomaly. Rock types logged in the hole include a number of intrusive units described as gabbro, trachyte, feldspar porphyry, and fine grained felsic dykes. Mafic to felsic flow rocks are also noted. The chargeability anomaly again appears to be caused by zones of disseminated pyrite in quartz-carbonate-sericite alteration within felsic volcanic rocks, although narrow zones of 5% disseminated pyrite also occur within a feldspar porphyry dyke, as well as a gabbroic unit. The highest gold value of 179 ppb was returned from a narrow pyritic quartz carbonate healed fracture zone in gabbro. A gold value of 79 ppb was returned from a pyritic feldspar porphyry dyke with 5% disseminated pyrite.

MP12-01 (See figure 9) This hole tested a very weak chargeability anomaly in area with high gold in till counts. The entire hole was logged as a syenite to syenodiorite, generally fractured and altered with a carbonate-chlorite stockwork. Very little sulphide was noted. Gold values are locally anomalous up to 324 ppb.

CKP12-01 (See figure 10) This hole tested a strong surface chargeability anomaly. Rock types logged in the hole consist mainly of pillowed and massive porphyritic mafic volcanics, cut by several diabase dykes. The chargeability does not appear to be explained by the relatively minor amounts of disseminated pyrite in mafic flows and magnetite in the diabase dykes. Gold values are generally below 5 ppb with a high value of 31 ppb.

7.0 CONCLUSIONS AND RECOMMENDATIONS

The 2012 Beilby-Clarke drilling program generally focused on testing of strong chargeability anomalies outlined by a relatively small (10.9 km) Pole-Dipole IP/Resisitivity survey. These anomalies were explained by disseminated pyrite zones within volcanic as well as intrusive rocks. Gold values associated with chargeability anomalies were generally at or below detection limit, with locally anomalous values (20 to 140 ppb). It should be noted that much of the core remains to be sampled.

The most significant gold mineralization occurred in MP12-01, which targeted a very weak chargeability anomaly and intersected only trace to minor amounts of pyrite mineralization. Gold mineralization occurs within what appears to be a relatively large body of syenite or syenodiorite that has been chloritized and carbonatized along a stockwork style fracture system. Pyrite concentrations generally occur in trace amounts, but locally can approach concentrations of 1%.

It is recommended that the remainder of drill hole MP12-01 be split and sampled for gold. Selected, representative multi-element, whole rock analyses, and thin section description are suggested for each distinct lithology intersected in each drill hole of the 2012 drilling program. Additional gold analyses should be considered in locations where anomalous gold values (>20 ppb) occur.

The new gold occurrence south of Beilby Lake was not directly tested during the 2012 drilling program. Additional mapping in the Beilby area should be undertaken, integrating the results of the 2012 geophysics and drilling programs.

8.0 REFERENCES

Ayer, J.A. 2000. The Abitibi greenstone belt: a program overview; in Summary of Field Work and Other Activities 2000, Ontario Geological Survey, Open File Report 6032 p 6-1 to 6-14.

Ayer, J.A. 1999. Geological compilation of the Abitibi greenstone belt in Ontario; toward a revised stratigraphy based on compilation and new geochronology results; Summary of Field Work and Other Activities, Ontario Geological Survey, Miscellaneous Paper 169, p.14-24.

Carter, M.W. 1987. Geology of the Shining Tree Area, districts of Sudbury and Timiskaming; Ontario Geological Survey, Report 240, 48p.

Cluff, G.R. 1990. Exploration Activities – 1989 Herrick and Churchill Properties, Churchill Township, Ontario. Unocal Canada Limited.

Johns, G.W. 1996. Reappraisal of the Geology of the Shining Tree Area, districts of Sudbury and Timiskaming; Summary of Field Work and Other Activities 1997, Ontario Geological Survey, Miscellaneous Paper 166, p.13-15.

Johns, G.W. 1997. Reappraisal of the geology of the Shining Tree area, districts of Sudbury and Timiskaming. Summary of Field Work and Other Activities 1997, Ontario Geological Survey, Miscellaneous Paper 168, p.26-29.

Johns, G.W and Amelin, Y. 1999. Reappraisal of the geology of the Shining Tree area (East Part), districts of Sudbury and Timiskaming; Summary of Field Work and Other Activities 1999, Ontario Geological Survey, Miscellaneous Paper 169, p.43-50.

Johns, G.W. 1999a. Reappraisal of the geology of the Shining Tree area (West Part), District of Sudbury; Summary of Field Work and Other Activities 1999, Ontario Geological Survey, Open File Report 6000, p.6.1-6.7. Johns, G.W. 1999b. Precambrian geology, Shining Tree area (east half); Ontario Geological Survey, Preliminary Map P.3389.

Johns, G.W. 2000. Precambrian geology, Shining Tree area (west half); Ontario Geological Survey, Preliminary Map P.3420.

Oliver, H.S., Hughes, D.J., Hall, R.P. and Johns, G.W. 1999a. Preliminary Geochemistry of Metavolcanic rocks of the Shining Tree Area; Abitibi Subprovince, Ontario; Summary of Field Work and Other Activities 1998, Ontario Geological Survey, Miscellaneous Paper 169, p.51-58.

Oliver, H.S., Hughes, D.J., Hall, R.P. and Johns, G.W. 1999b. The Mafic And Ultramafic Volcanics Of The ShiningTree Greenstone Belt, Northeastern Ontario, Canada; Summary of Field Work and Other Activities 1999, Ontario Geological Survey, Open File Report 6000, p.11.1-11.12.

Oliver, H.S., Johns, G.W., Thurston, P.C., Hughes, D.J. and Hall R.P. 1998 Preliminary geochemistry of the Shining Tree area; Abitibi Subprovince, Ontario; in Summary of Field Work and Other Activities 1998, Ontario Geological Survey, Miscellaneous Paper 169, p.51-58.

Webster, B., 2013. Geophysical Report on Spectral IP/Resistivity Surveys on the Beilby Lake Project, Shining Tree Area, Northeastern Ontario. REF 12-009.

Certificate of Qualifications: James R. Trusler

I, James R. Trusler at 20 William Roe Blvd, Suite 807, Newmarket, Ontario do hereby certify that:

- 1) I am a Geological Engineer employed as President and CEO of Platinex Inc. and I am also the principal shareholder of Platinex Inc.;
- I graduated from the University of Toronto with BA.Sc. in Geological Engineering in 1967. I obtained a Master of Science (Geology) from Michigan Technological University in 1972. I have practiced my profession full-time from 1967-1969 and from 1970 to present;
- I am a Professional Engineer registered with the Professional Engineers Ontario (PEO #47064019);
- 4) I have not received, nor do I expect to receive, any interest, directly or indirectly, in the Shining Tree property;
- 5) As of the date of this certificate, to the best of my knowledge, information and belief, this report contains all scientific and technical information that is required to be disclosed to make the Diamond Drilling Beilby, McBride and Clarke on the Shining Tree Project for Platinex Inc. not misleading;
- 6) I have read National Instrument 43-101 and supervised the completion of the Diamond Drilling Beilby, McBride and Clarke on the Shining Tree Project for Platinex Inc. which has been prepared in compliance with the intent of National Instrument 43-101 and Form 43-101F1 but is not a Technical Report as defined by National Instrument 43-101;
- 7) I have collaborated with lain Trusler who prepared Diamond Drilling Beilby, McBride and Clarke on the Shining Tree Project for Platinex Inc. under my supervision;
- 8) I have visited the property on several occasions and during the program.

James R. Truster, BASC, MS, PEng

Dated at Newmarket, ON March 7th, 2017

Certificate of Qualifications: Iain S. Trusler

I, Iain S. Trusler at 32 Richmond St., Richmond Hill, Ontario do hereby certify that:

- 1) I am a GIS consultant employed as such by Platinex Inc.;
- 2) I have practiced my profession full-time from 2010 to present;
- 3) I have not received, nor do I expect to receive, any interest, directly or indirectly, in the Shining Tree property;
- 4) As of the date of this certificate, to the best of my knowledge, information and belief, this report contains all scientific and technical information that is required to be disclosed to make the Diamond Drilling Beilby, McBride and Clarke on the Shining Tree Project for Platinex Inc. not misleading;
- 5) I have collaborated with James R Trusler who supervised Diamond Drilling Beilby, McBride and Clarke on the Shining Tree Project for Platinex Inc.;
- 6) I have visited the property once in October of 2011, and twice in March and April of 2012

Dated at Richmond Hill, ON March 7th, 2012

Iain S. Trusler wind





















Appendix I

Drill Hole Logs

PLATINEX INC.

SHINING TREE PROJECT, ONTARIO -

Grid Coordinates:		UTM - NAD83 - Zone 17							
		N:	5271434	E:	428848				
Dip:	-45		Elevation:		410 m				
Azimuth:	270		Total Depth:		170 m				
Core Size:	NQ		Core Boxes:		40				

Туре	Depth	Dip	Azimuth
EZ-Shot	30	-45.9	269.70
	60	-46.3	271.90
	90	-46.9	275.70
	120	-46.9	273.60
	150	-46.8	274.20

subtracting 10.5 degrees from the EZ shot instrument reading.

HOLE#: BP12-01

Page	1	of	3
Date Started:	March 21	, 2012 (N)	
Date Completed:	March 22	, 2012 (N)	
Claim#:	4207971		
Contractor:	Laframbo	oise Drillin	g
			~

Target: IP Anomaly in the vicinity of the Beilby Lake Gold Occurrence.

Logged by: Dean R. Cutting, P.Geo. Sampled by: CXS

De	pth	Rock Type	Description	Struct. core angles Strai		Struct. core angles		Struct. core		Struct. core angles Strain Alter		Struct. core angles		t. core angles		Alteration Characteristics			Sample Assays						
From	То			S.		Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t								
0.00	1.50	Overburden	Casing left in the hole.																						
		OVB																							
1.50	100.00	Mafic Flow - Pillowed	Medium to dark green colour. Moderately hard. Pillow selveges are locally quite																						
		PMFLOW	evident. Medium to fine grained locally. Amygdules locally notable over short																						
			intervals, may be round or elongate, mostly 2-5 mm diameter but may be up to																						
			1 cm on the long axis. Orientations of the long axes do not seem preferential.																						
			Most amygdules are carbonate filled and white to beige colour. Unit is moderately																						
			chloritized. Numerous carbonate, carbonate/quartz, ± epidote veinlets cut the																	1					
			unit. Most are very irregular and vary from hairline to 4 or 5 cm thick, often																						
			asssociated with pillow selveges. Pyrite is notable in some of the veinlets from																	1					
			trace to 2% as disseminations and semi-massive patches. The carbonate seems																						
			to be carbonate as opposed to Fe carbonate as the veinlets are white as																						
			opposed to rusty oxidized after sitting exposed to the weather for an extended																	1					
			period of time. Some of the carbonate has a light pinkish tinge to it. Unit is lightly																						
			to moderately magnetic locally.																	1					
			Tr - 2+% PY locally in Cabonate veinlets, disseminated to semi-massive patches.										K403818	32.00	33.00	1.00	0.015								
			At 36.20 m - 6 cm thick carbonate, chlorite, epidote veinlet with Tr-1%																						
			disseminated Pyrite with brick red (hematite?) at 20 - 30 ° to core axis.																	1					
			Carbonate, ± quartz, tr epidote, and tr brick red hematite? locally. Tr - 2 % Pyrite										K403819	62.10	63.60	1.50	<0.005			1					
			locally as disseminations and semi-massive patches. Veinlets have variable																						
			angles to the core axis.																	1					
																				4					
			At 74.0 m to 75.40 m - Core missing from Box # 18. Total 1.4 m seems to me as																	1					
			If a sample was collected by person or persons unknown ???	I			I		<u> </u>				<u> </u>						 	──					
																				1					
				-			-													1					
								1		1															
								1	1	İ	1		1							t –					

Hole # BP12-01

Page 2 of 3

De	pth	Rock Type	Description	Struc	ct. core	e angle	s	Strain	Alterat	ion Chara	cteristics		Sample Assays								
From	To			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t				
											1	1									
100.00	170.00	Mafic Volcanic	Somewhat arbitrarily and gradationally there seems to be a change from the Pillowed Mafic																		
		MFLOW	Volcanic to a more massive Mafic Volcanic Flow. Basically the unit is as above described but																		
			without the pillow selveges quite as evident. The unit is fine to medium grained and massive.																		
			Pyrite seems to be more evident locally as disseminated grains or small masses. Locally																		
			amygdules are notable over short intervals. Irregular fractures are moderately chloritized some																		
			with carbonate ± epidote as well. Generally speaking, not as magnetic as the pillowed unit.																		
			Irregular veinlets and fractures are generally at moderate to high angles to the core axis.																		
			Carbonate is white carbonate not seeming to be Fe carbonate. Pyrite more common in the																		
			unit occurring as dissemonation of grains, blebs, and small masses associated with carbonate																		
			fracture / veinlet systems, and with carbonate as replacement in amygdules. Locally, where the																		
			unit is not epidotized the volcanics are light to moderately carbonatized.																		
			Grey / white carbonate veinlet in a 45° to core axis healed fracture / slip zone.										K403820	107.10	107.40	0.30	< 0.005				
			Epidote / sericite alteration associated. 2 - 3 % Pyrite as irregular masses in the veinlet.																		
			Notable increase in Pyrite content at about 120 m to the end of the hole. Pyrite occurs primarily																		
			as disseminated grains or small masses up to 2 cm diameter locally. Pyrite occurs as well																		
			associated with irregular carbonate veinlets and as semi-massive concentrations in what																		
			almost took like primary bands.																		
			At approximately 122 m or so the concentration of irregular carbonate, chlorite veinlets																		
			decreases somewhat.																		
			Unit Sampling:																		
			Carbonate, chlorite veinlet at high angle to the core axis. 15 - 20 % granular to										K403821	143.05	143.25	0.15	0.036				
			semi-massive pyrite.																		
			Olive coloured mafic volcanic with Tr-3% disseminated pyrite.										K403822	143.25	144.05	0.80	< 0.005				
-																					
-			Carbonate veinlet at high angle to the core axis. 5 - 7 % disseminated pyrite as grains and blebs.										K403823	144.05	144.30	0.25	0.034				
-																					
										1	1										

Hole # BP12-01

Page 3 of 3

De	pth	Rock Type	Description	Struct. core angles Strain Alteration Characteristics					Sample Assays												
From	То			s。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t				
			Olive coloured mafic volcanic with Trace - 2 % disseminated pyrite as grains and blebs.										K403824	144.30	144.75	0.45	< 0.005				
			High Au Standard # 17c										K403825	ORE/	AS AU STD #	17c	3.095				
			144.75 m to 145.85 m section notably banded at 50° to 60° to the core axis. Very strong pyrite										K403826	144.75	145.85	1.10	0.070				
			content to semi-massive for a 10 cm section at the bottom of the interval. Someone stole the																		
			section from 145.20 m to 145.65 m in the middle of the interval, likely the best part judging																		
			by the ends that were left. May have been a banded massive sulphide section??																		
			Carbonate/quartz veinlet with grey fine graimed mafic volcanic. Disseminated pyrite to										K403827	149.55	149.95	0.40	0.14				
			15 - 20 % locally. Upper contact at 50° to the core axis. Lower contact at 30° to the core axis.																		
			Carbonate/quartz/chlorite fracture zone at 30° to the core axis. Strong semi-massive pyrite for										K403828	153.35	154.75	1.40	0.032				
			about a 20 cm section of the core. Trace to 2 % pyrite for the balance of the interval.																		
			Carbonate/quartz/chlorite veinlet system at 30° to the core axis. ± 10 - 15 % disseminated										K403829	155.90	156.15	0.25	0.027				
			and semi.massive pyrite.																		
			From about 157 m or so to the bottom of the hole the unit becomes s bit coarser grained to																		
			medium grained with local finer grained sections. The unit becomes more strongly magnetic																		
			as well. May either be a coarsening of the flow or becoming more of a gabbroic unit.																		
170.00		EOH	End of Hole.																		

PLATINEX INC.

SHINING TREE PROJECT, ONTARIO -

Grid Coordinates:			<u>UTM - N</u>	AD83 - Zo	ne	<u>17</u>
		N:	5271331	E:	4	82984
Dip:	-55		Elevation:	404 m		
Azimuth:	0		Total Depth:	200 m		
Core Size:	NQ		Core Boxes:	47		

Down note Tests												
Туре	Depth		Dip	Azimuth								
EZ-SHOT	30		-55.5	1.1								
	60		-55.1	358.5								
	90		-54.9	0.4								
	120		-54.6	1.4								
	150		-54.3	4.0								
	180		-53.9	4.0								
Downhole azimuth readings have been corrected to true north by												

subtracting 10.5 degrees from the EZ shot instrument reading.

HOLE#: BP12-02

Pa	age	1	of	12
Date Started	1:	2012-03	-23 (D)	
Date Compl	eted:	2012-03	-24 (D)	
Claim#:		420797	1	
Contractor:		Laframb	oise Drilli	ng
Logged by: Typed by:		Dean R. Iain Trus	Cutting	Sampled by

IP Anomaly in the vicinity of the Beilby Lake Gold Occurrence. Target:

Sampled by: Bruce Cupskey

De	pth	Rock Type	Description	Struc	t. cor	e ang	les	Strain	Alterati	on Charac	teristic	S				Sample	Assays				
From	To			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t				
0.00	2.30	Overburden	Casing left in hole.																		
		OVB																			
2.30	14.25	Diabase	Medium to dark green grey mottled finely with grey. Medium to fine grained																		
		DIAB	equigranular and massive through the unit. Moderately to strongly																		
			magnetic through the unit. Rubbly at the top with short intervals of 'porous'																		
			sections with carbonate veinlets dissolved. As at 8.5 to 8.8m depth shot																		
			with a fine stockwork of carbonate, quartz carbonate plus or minus																		
			epidote veinlets from hairline to 0.75 cm thick. Greenish chloritic material																		
			along the walls in some of the thicker veinlets. A few veinlets noted																		
			on a 'preferred' orientation of plus or minus 35 degrees to the core axis.																		
			Only trace sulphides, pyrite noted occasionally. Lower unit contact is																		
			sharp at 55 degrees to core axis, xenolits of volcanics in diabase near																		
			the contact. But contact neither chilled or cooked No sampling in unit																		
																					1
14.25	25.80	Mafic Flow	Fine grained, equigranular, massive though heavily veined with a																		
		MFLOW	stockwork of fine carbonate, quartz carbonate, and epidote veinlets.																		
			Medium to dark green in colour. Rubbly in sections with rusty fractures,																		
			may be partially due to dissolution of some of the carbonate veinlets.																		
			Sparse disseminatated pyrite through the unit, as well as in association																		
			with some of the carbonate, quarts, and epidote veinlets. Some of the																		
			veinlets are fine "ladder type" breccia infills. Some of the carbonate																		
			has a pinkish tinge. Lower contact (30cm) rubbly and more on colour and																		
			grain size variance.																		
			Unit Sampling																		
			Rubbly section with poor recovery, Trace pyrite, Trace quartz carbonate										K403551	17.00	18.00	1.00	<0.005			'	
			veinlets.																	'	
			Trace pyrite locally with quartz carbonate veinlets							ļ			K403552	18.00	19.00	1.00	< 0.005			ļ'	
			I race pyrite with stronger quartz carbonate stockwork veinlets, pink										K403553	19.00	20.00	1.00	<0.005			<u> </u>	
			carbonate locally.										K402554	20.00	21.00	1.00	<0.00F	<u> </u>	 	'	-
			mace pyrite with light quartz carbonate verniets.	┢──┦									11403034	20.00	21.00	1.00	<0.005	┢───┤	┟───┦	i'	┢──┤
25.80	36.25	Gabbro	Maybe a coarse grained flow. Colour more of a medium grey green than																	'	
_0.00	50.20	GAB	the dark green of the mafic volcanics above. Coarser to medium grained							l											

BP12-02

Page 2 of 12

De	pth	Rock Type	Description	Struc	ct. cor	e angl	es	Strain	Alterati	on Chara	acteristi	ics				S	Sample A	ssays		
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			relatively equigranular throughout. Generally massive with no typical primary																	
			volcanic textures evident. Strong fine beige "leucoxene" xtals easily notable to																	
			several percent. 'Patchy' disseminated pyrite locally notable to 1-2% sulphides as																	
			grains or small masses. Fractures are generally sharp and filled with carbonate																	
			plus or minus chlorite and epidote. Stockwork veinlets are locally 'offset' up to a																	
			couple of centimetres. Towards the bottom of the unit, starts to develop a locally																	
			yellowish tinge from light sericite?																	
			31.75 - 32.60 Flooded quartz breccia zone with 60+% quartz fragments. One																	
			carbonatized/ sericitized moderately to strongly. Pyrite is strong 3-5% as granular																	
			masses. Quartz may have several phases as there are both white and grey																	
			versions with trace ammounts of tourmaline? Basically in strain noted in the walls																	
			of the veinlet, seems to be a brittle break. Seems to lie at 50 to 60 degree to																	
			core axis.																	
			Unit Sampling																	
			Irregular quartz carbonate veinlets with trace disseminated pyrite in veinlets and										K403555	26.00	27.00	1.00	< 0.005			
			rock.																	
			Irregular quartz carbonate veinlets with trace disseminated pyrite in veinlets and										K403556	27.00	28.00	1.00	< 0.005	-		
			rock.																	
			Irregular quartz carbonate veinlets with trace disseminated pyrite in veinlets and										K403557	28.00	29.00	1.00	< 0.005			
			rock.																	
			Irregular quartz carbonate veinlets with trace disseminated pyrite in veinlets and										K403558	29.00	30.00	1.00	< 0.005	-		
			rock.																	
			Irregular quartz carbonate veinlets with trace disseminated pyrite in veinlets and										K403559	30.00	31.00	1.00	< 0.005	-		
			rock.																	
			Irregular quartz carbonate veinlets with trace disseminated pyrite in veinlets and										K403560	31.00	31.75	0.75	0.009			
			rock.																	
			See description of quartz flood zone in unit.										K403561	31.75	32.60	0.85	0.036			
			As above the quartz flood zone.										K403562	32.60	33.60	1.00	0.005			
			As above the quartz flood zone.										K403563	33.60	34.60	1.00	0.008			
			As above the quartz flood zone.										K403564	34.60	35.45	0.85	0.007			
			As above the quartz flood zone.										K403565	35.45	36.25	0.80	0.009			

BP12-02

Page 3 of 12

De	pth	Rock Type	Description	Stru	ct. cor	e angl	es	Strain	Alterat	ion Char	acterist	ics				s	Sample A	ssays		
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
36.25	38.80	Mafic Volcanics	Sliver of mafic volcanics between the gabbro and felsic volcanics below. Basically					1												
		MFLOW	similar to the mafic volcanic unit above the gabbro except starting to show some																	
			sericite alteration (yellow/green) as patches along fractures. Some pink carbonate																	
			noted in some veinlets. Unit is medium to dark green in colour, fine to medium grained																	
			and essentially non-magnetic. Quartz carbonate and quartz carbonate breccia																	
			veinlets up to 2cm are notable. Broken in brittle fracture but not really foliated.																	
			Light to medium sericitization on fractures. Locally strong disseminated pyrite as																	
			grains or small masses up to 5%. Lower contact over 5cm, though relatively sharp																	
			with intermixed pieces, high angle to core axis. Locally some veinlets running at plus																	
			or minus 45 degrees to core axis, may or may not be significant.																	
			Unit Sampling																	
			Trace to 2% pyrite associated with fractures and quartz carbonate veinlets, locally										K403566	36.25	37.00	0.75	< 0.005			
			disseminated grains or small masses																	
			Trace to 2% pyrite associated with fractures and quartz carbonate veinlets, locally										K403567	37.00	38.00	1.00	0.006			
			disseminated grains or small masses																	
			Trace to 5% pyrite disseminated locally										K403568	38.00	38.80	0.80	0.018			
38.80	41.90	Felsic Volcanics	Fine to medium grained, yellow green (sericite) colour, massive and equigranular unit																	
		FFLOW	Relatively hard unit. Non-magnetic. Shot with stockwork of carbonate and quartz																	
			carbonate veinlets with pyrite associated. Sulphides (pyrite) usually along fractures																	
			or as small masses up to a cm or so along healed fractures. Do not believe this																	
			section is 'strained' more on a basis of irregular brittle fracture. Trace to 3%																	
			sulphides locally.																	
			Unit Sampling																	
			Trace to 3% pyrite locally with quartz carbonate veinlets and 'pods' or healed										K403569	38.80	40.35	1.55	< 0.005			
			fractures																	
			Trace to 3% pyrite locally with quartz carbonate veinlets and 'pods' or healed										K403570	40.35	41.90	1.55	< 0.005			
			fractures																	
41.90	43.15	Gabbro	Unit basically the same as the gabbro unit above (25.8m - 36.25m). 10cm upper																	
		GAB	contact a mix with the felsic volcanics, lower contact is sharp but irregular. Very																	
			strong disseminated pyrite as grains or small masses up to 5% or better locally.																	

BP12-02

Page 4 of 12

De	pth	Rock Type	Description	Stru	ct. cor	e angl	les	Strain	Alterati	on Chara	acterist	ics				:	Sample A	ssays		
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Becoming sericitized with a yellowish tint developing near the margins and along																	
			some of the fractures. Irregular quartz carbonate or carbonate veinlets in stockwork																	
			are common. Strong beige "leucoxene" specking is notable																	
			Unit Sampling																	
			Up to 5% disseminated pyrite and quartz carbonate or carbonate veinlets through										K403571	41.90	43.15	1.25	0.012			
			interval																	
43.15	85.80	Felsic Volcanic	Fine to medium grained, relatively massive equigranular unit. On the felsic end of																	
		FFLOW	the scale but not apparently as hard as a rhyolite, more dacitic - rhyolitic unit																	
			composition. Striking light to medium yellow-green colour. Moderately sericitized																	
			weaker to stronger locally. Fine stockwork of hairline to 2cm carbonate and quartz																	
			carbonate veinlets throughout. Looks as if shattered and healed. Unit can't be																	
			scratched with steel pin where less sericitized. Unit contains trace to 3% locally																	
			disseminated pyrite in blebs, disseminated xtals, associated with fractures and																	
			veinlets. Very little for primary volcanic textures. No notable strain in the unit, more																	
			brittle fracture likely. Unit is non-magnetic. This unit is likely the chargeability																	
			anomaly.																	
			Unit Sampling																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as grains, small										K403572	43.15	44.15	1.00	< 0.005			
			masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as grains, small										K403573	44.15	45.10	0.95	0.007			
			masses or with veinlets.																	
			Brecciated quartz carbonate veinlets with pink carbonate, white and grey quartz										K403574	45.10	46.35	1.25	0.014			
			trace to 3% granular pyrite.																	
			Low Au Standard # 65a										K403575	STD OR	EAS 65a	0.00	0.494			
			Brecciated quartz carbonate veinlets with pink carbonate, white and grey quartz										K403576	46.35	47.00	0.65	0.007			
			trace to 3% granular pyrite.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403577	47.00	48.50	1.50	0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403578	48.50	50.00	1.50	0.005			
			grains, small masses or with veinlets.																	
				1																

BP12-02

Page 5 of 12

De	pth	Rock Type	Description	Strue	ct. core	e angle	es	Strain	Alteratio	on Chara	cteristi	cs					Sample A	ssays		
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403579	50.00	51.50	1.50	< 0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403580	51.50	53.00	1.50	< 0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403581	53.00	54.50	1.50	< 0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403582	54.50	56.00	1.50	< 0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403583	56.00	57.50	1.50	< 0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403584	57.50	59.00	1.50	0.011			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403585	59.00	60.50	1.50	< 0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403586	60.50	62.00	1.50	0.008			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403587	62.00	63.50	1.50	0.01			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403588	63.50	65.00	1.50	0.012			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403589	65.00	66.50	1.50	0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlet stockwork with trace to 3% pyrite locally as disseminated										K403590	66.50	68.00	1.50	0.01			
			grains, small masses or with veinlets.																	
			74 to 85.8m Picking up patches of pinkish felsic volcanics, what is likely the																	
			original colour before sericitization to yellow green. Boundaries between the pink																	
			sections and yellow-green can be sharp or like 'clouds'. Cracks in the 'pink' felsic																	
			volcanics are often showing sericite alteration along them. Quartz carbonate																	
			veinlets with pyrite grans and masses permiate both types of felsics. Sulphide																	
			pyrite present at trace to 3% locally as disseminated grains or small masses through																	
			the unit and/or associated with the stockwork veins.																	

BP12-02

Page 6 of 12

De	pth	Rock Type	Description	Struc	ct. cor	e angl	es	Strain	Alteratio	on Chara	acterist	ics				:	Sample As	says		
From	To			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Unit Sampling Continued																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403591	68.00	69.50	1.50	0.018			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403592	69.50	71.00	1.50	0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403593	71.00	72.50	1.50	0.015			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403594	72.50	74.00	1.50	0.005			
			grains, small masses or with veinlets.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403595	74.00	75.50	1.50	0.005			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403596	75.50	77.00	1.50	0.011			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403597	77.00	78.50	1.50	0.006			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403598	78.50	80.00	1.50	< 0.005			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403599	80.00	81.50	1.50	< 0.005			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Au Standard # 61d										K403600	STD OR	EAS 61d	0.00	4.62			
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403601	81.50	83.00	1.50	0.008			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403602	83.00	84.50	1.50	0.009			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
			Quartz carbonate veinlets stockwork with trace to 3% pyrite locally as disseminated										K403603	84.50	85.80	1.30	0.008			
			grains, small masses or with veinlets. Starting to pick up some pink material.																	
85.80	137.20	Felsic Volcanics	Basically the same felsic volcanics as above described but the yellow-green																	
		FFLOW	sericite alteration is not as strongly developed, occurring more in patches over																	
			a few metres and along fractures in a more pickish version of the felsic volcanic																	
			unit. The unit is still fine to medium grained, very hard as a general rule. Non-																	
			magnetic. Still hosting a fine stockwork of quartz carbonate, carbonate, and sericite																	
			veinlets from hairline to several cm in thickness. Pyrite is variably present through																	
			the unit from trace to 3% locally as grains, small masses or associated with the		1															

BP12-02

Page 7 of 12

De	pth	Rock Type	Description	Struc	t. cor	e angle	es	Strain	Alteratio	on Chara	cteristi	cs					Sample A	ssays		
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			fracture / veinlet systems. Other than the odd breccia piece, not much notable for																	
			primary textures. The colour variance would believe to be a question of variable																	
			alteration overprinting, as opposed to a primary textural feature. Unit appears to be																	
			more 'brittle fractured' than strained.																	
			Unit Sampling																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403604	85.80	87.00	1.20	0.009			
			volcanic. Trace to 3% pyrite locally (rubbly locally)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403605	87.00	88.00	1.00	0.006			
			volcanic. Trace to 3% pyrite locally (rubbly locally)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403606	88.00	89.00	1.00	< 0.005			
			volcanic. Trace to 3% pyrite locally (rubbly locally)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403607	89.00	90.50	1.50	0.006			
			volcanic. Trace to 3% pyrite locally (rubbly section)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403608	90.50	92.00	1.50	0.005			
			volcanic. Trace to 3% pyrite locally (stronger quartz fracture system, breccia)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403609	92.00	93.50	1.50	0.012			
			volcanic. Trace to 3% pyrite locally (bottom third more pinkish)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403610	93.50	95.00	1.50	0.013			
			volcanic. Trace to 3% pyrite locally (top third more pinkish)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403611	95.00	96.50	1.50	0.006			
			volcanic. Trace to 3% pyrite locally (strong sericitization)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403612	96.50	98.00	1.50	0.01			
			volcanic. Trace to 3% pyrite locally (strong sericitization)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403613	98.00	99.50	1.50	0.009			
			volcanic. Trace to 3% pyrite locally (strong sericitization)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403614	99.50	100.50	1.00	0.008			
			volcanic. Trace to 3% pyrite locally (weaker sericitization along fractures)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403615	100.50	101.60	1.10	0.065			
			volcanic. Trace to 3% pyrite locally (pink volcanics with weak sericite alteration,																	
			chloritized on some of the fractures, high angle to core axis.)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403616	101.60	103.00	1.40	0.018			
			volcanic. Trace to 3% pyrite locally (weaker sericitization along fractures pink volc)																	

BP12-02

Page 8 of 12

De	pth	Rock Type	Description	Struc	t. core	e angle	S	Strain	Alteratio	on Chara	cteristi	cs					Sample A	ssays		
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403617	103.00	104.00	1.00	0.008			
			volcanic. Trace to 3% pyrite locally (weaker sericitization along fractures pink volc)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403618	104.00	105.50	1.50	0.009			
			volcanic. Trace to 3% pyrite locally (weaker sericitization along fractures pink volc)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403619	105.50	107.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (weaker sericitization along fractures pink volc)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403620	107.00	108.50	1.50	0.006			
			volcanic. Trace to 3% pyrite locally (Strong sericitization, olive green colour)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403621	108.50	110.00	1.50	0.016			
			volcanic. Trace to 3% pyrite locally (Strong sericitization, olive green colour)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403622	110.00	111.50	1.50	0.005			
			volcanic. Trace to 3% pyrite locally (Patchy sericitization along veinlets and fracture)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403623	111.50	113.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Patchy sericitization along veinlets and fracture)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403624	113.00	114.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Patchy sericitization along veinlets and fracture)																	
			Blank										K403625	BLA	NK	0.00	< 0.005			
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403626	114.50	116.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Patchy sericitization along veinlets and fracture)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403627	116.00	117.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Patchy sericitization along veinlets and fracture)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403628	117.50	119.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Patchy sericitization along veinlets and fracture)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403629	119.00	120.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Sericite increasing intensity)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403630	120.50	122.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (moderate and light sericite with fractures)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403631	122.00	123.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (moderate and light sericite with fractures)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403632	123.50	125.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (moderate to light sericite with fractures)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403633	125.00	126.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Light sericite with fractures)																	

BP12-02

Page 9 of 12

De	oth	Rock Type	Description	Strue	ct. cor	e angl	es	Strain	Alteratio	on Chara	cteristi	cs					Sample A	ssays		
From	To			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403634	126.50	128.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Light sericite with fractures)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403635	128.00	129.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (Light to moderate sericite with fractures)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403636	129.50	131.00	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (moderate to strong sericite with fractures and																	
			patches in pink felsic volcanics)																	
			Quartz carbonate or carbonate veinlets in stockwork with sericite in pinkish felsic										K403637	131.00	132.50	1.50	< 0.005			
			volcanic. Trace to 3% pyrite locally (moderate to strong sericite with fractures and																	
			patches in pink felsic volcanics)																	
			At about 132.5m unit starting to exibit some poorly to moderately developed strain																	
			at 45 to 55 degrees to the core axis. Rubbly zone with (kaolin (white powder) in																	
			fracture) at about 133.3 to 133.5m Some of the fragments have a bit of a quartz																	
			carbonate veinlets in them. No sulphides noted. Foliation drops off to not notable at																	
			about 137.2m at the same time as the yellow area sericization drops significantly in																	
			intensity.																	
			Unit Sampling																	
			Moderate to strong sericitation in felsic volcanics, trace to 1% pyrite as										K403638	132.50	134.00	1.50	< 0.005			
			disseminated grains, still stockwork of quartz carbonate.																	
			Moderate to strong sericitation in felsic volcanics, trace to 1% pyrite as										K403639	134.00	135.20	1.20	< 0.005			
			disseminated grains, still stockwork of quartz carbonate.																	
			Moderate to strong sericitation in felsic volcanics, trace to 1% pyrite as										K403640	135.20	136.20	1.00	< 0.005			
			disseminated grains, still stockwork of quartz carbonate.																	
			Moderate to strong sericitation in felsic volcanics, trace to 1% pyrite as										K403641	136.20	137.20	1.00	< 0.005			
			disseminated grains, still stockwork of quartz carbonate.																	
			Probable end of felsic volcanic unit.																	
137.20	150.00	Diabase	Massive, medium to dark grey green colour, equigranular for the most part. Very uniform																	
		DIAB	looking with magnetics varying from light to strong going trom top to bottom of the																	
			unit. Irregular stockwork of carbonate and epidote, specular hematite (locally)																	
			through the unit. Contacts relatively sharp at high angles to the core axis. Contacts																	
			more assigned on textural and colour differences between this unit and those																	
BP12-02

Page 10 of 12

De	oth	Rock Type	Description	Strue	ct. core	e angle	es	Strain	Alterati	on Chara	acteristi	ics					Sample A	ssays		
From	To			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			above and below. Only an occasional speck of pyrite noted locally through the unit																	
			Unit Sampling																	
			Upper contact sampling to close felsic unit samples										K403642	173.20	138.70	1.50	< 0.005			
			Lower contact sampling										K403643	148.50	150.00	1.50	< 0.005			
150.00	158.65	Gabbro	Unit similar to unit above but a little more felsic in composition, a bit lighter gey green																	
		GAB	colour, less magnetic (more spotty, to moderate), more quartz carbonate, carbonate																	
			veinlets notable and trace to 2% disseminated pyrite noted through the unit. Unit is																	
			medium to coarse grained, maybe 60/40 feldspar to mafics in modal composition.																	
			There are a few chloritic 'xenoliths' up to a couple cm in size floating around in the																	
			matrix. From 158m to the bottom of the unit there is some notable strain (foliation)																	
			lightly to moderately developed at 60 to 70 degrees to the core axis, more intense																	
			at the lower unit contact. Lower unit contact gradational over 10cm with small																	
			fragments of the unit below dragged into this unit. Very little notable for primary																	
			features. Irregular quartz carbonate, and carbonate veinlets are notable through																	
			the unit, only occasionally a speck of pyrite is noted in associaion.																	
			Unit Sampling																	
			Top contact with trace pyrite										K403644	150.00	151.50	1.50	< 0.005			
			15 cm thick pinish carbonate breccia veinlet with 2-4% disseminated pyrite, crosscut										K403645	156.00	156.50	0.50	< 0.005			
			by white carb veinlet, zone at around 60 degrees to core axis, white carb veinlet																	
			0.5cm, around 30 degrees to core axis.																	
			Quartz carbonate veinlets with trace pyrite, trace to 2% disseminated pyrite locally										K403646	156.50	158.00	1.50	< 0.005			
			Quartz carbonate veinlets with trace pyrite, trace to 2% disseminated pyrite locally										K403647	158.00	158.65	0.65	< 0.005			
158.65	171.75	Felsic to Inter-	Mixed pyroclastic unit with beds of lapilli tuff interbedded with ash tuff. Some of																	
		mediate pyro-	the lapilli fragments can be up to 10 to 15cm. Some of the ash beds can be up																	
		clastic unit	to 2m thick or so. Some of the lithic fragments are xtals or 'lathlike' and altered to a																	
		FTUFF	metallic' green like mariposite locally. The 'fabric' in the rock seems to be between																	
			50 to 60 degrees to core axis. Variable in colour from medium grey to beige mottled.																	
			Unit is non-magnetic. Unit variably carbonatized from weak to strong locally. Unit																	
			shot with quartz carbonate or carbonate veinlets. The entire unit is strongly pyritic													-				
			usually as disseminated grains or small masses or with fractures and veinlets,																	

BP12-02

Page 11 of 12

Dep	oth	Rock Type	Description	Struc	t. core	angl	es	Strain	Alteratio	on Chara	octeristi	CS	Sample Assays							
From	To			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			locally can be over 10 %.																	
			Unit Sampling																	
			Sericitized felsic pyroclastic, quartz carbonate veinlets with trace to 2% pyrite										K403648	158.65	159.95	1.30	0.005			
			Sericitized felsic pyroclastic, quartz carbonate veinlets with trace to 2% pyrite										K403649	159.95	160.75	0.80	0.009			
			Low Au Standard #65a										K403650	STD OR	EAS 65a	0.00	0.523			
			Sericitized felsic pyroclastic, quartz carbonate veinlets with trace to 2% pyrite										K403651	160.75	161.60	0.85	0.018			
			(10 cm pink carbonate veinlet with 3% pyrite at high angle to core axis)																	
			Ash with 3-5% disseminated pyrite										K403652	161.60	162.75	1.15	0.007			
			Ash with 5-10% disseminated pyrite										K403653	162.75	163.95	1.20	< 0.005			
			Ash with 5-10% disseminated pyrite										K403654	163.95	165.45	1.50	< 0.005			
			Lapilli tuff with 2-4% pyrite										K403655	165.45	166.45	1.00	< 0.005			
			Lapilli tuff with 2-4% pyrite										K403656	166.45	167.40	0.95	< 0.005			
			Ash with 5-10% disseminated pyrite										K403657	167.40	168.30	0.90	< 0.005			
			Lapilli tuff with 2-3% pyrite, flecks of fuchite										K403658	168.30	169.45	1.15	< 0.005			
			Lapilli tuff with 2-3% pyrite, flecks of fuchite (Fragments of 4cm carbonate veinlet)										K403659	169.45	170.60	1.15	< 0.005			
			Lapilli tuff with 2-3% pyrite, flecks of fuchite										K403660	170.60	171.75	1.15	< 0.005			
171.75	182.40	Intermediate	Medium to fine grained, medium to dark green in colour. Variably magnetic from																	
		to Mafic	weak to strong locally. Relatively hard unit. Unit is rubbly from 175.25 to 179.5m																	
		Volcanics	locally over sections, may be a fault influence but no gouge noted anywhere. Shot																	
		MFLOW	with fine carbonate veinlets but very little pyrite noted anywhere																	
			179.5 to 182m Stockwork of carbonate veinlets with red amorphous mineral																	
			relatively soft, scratches with steel needle, scratch powder red may be either																	
			hematite or cinnabar? (should look into closer) May have an orientation of 35																	
			degrees to core axis. There are small masses of pyrite associated with the veinlets.																	
			Really suspect hematite.																	
			Unit Sampling																	
			Contact with pyroclastics above										K403661	171.75	73.00	1.25	< 0.005			
			Stockwork, described above.										K403662	179.50	180.50	1.00	< 0.005			
182.40	185.15	Diabase	Medium green grey colour, medium to fine grained, equigranular and massive.																	
		DIAB	Essentially non-magnetic. Shot with irregular stockwork of carbonate/epidote																	

BP12-02

De	oth	Rock Type	Description	Struc	ct. cor	e ang	lles	s Strain Alteration Characteristics Sample Assays											
From	To			So	Fol	Flov	v Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t		
			veinlets from hairling to 0.5cm thick. Contacts sharp at 40 degrees to core axis																
			No sampling of unit																
185.15	195.10	Trachyte	Medium to fine grained, green with a pinkish tinge to almost black locally. Very hard																
		TRACH	and brittle fracturing. Fragments noted locally up to about a cm in size, chloritized.																
			Strongly magnetic throughout. Locally patches are epidotized as well as along some																
			of the fractures. Shot with stockwork of carb veinlets. Very little to no sulphides																
			noted anywhere.																
			Unit Sampling																
			Stockwork of carbonate veinlets, some pinkish tinges, trace pyrite										K403663	186.85	187.85	1.00	< 0.005		
			Stockwork breccia of carbonate veinlets, white with some chlorite, trace pyrite										K403664	194.50	195.00	0.50	< 0.005		
195.10	198.25	Diabase	Fine to medium grained, massive and equigranular. Medium to dark grey in colour.																
		DIAB	Moderately to strongly magnetic throughout. Occasional carbonate/ epidote veinlets.																
			Contacts are sharp, upper is at 40 degrees to core axis, lower is irregular with a																
			half cm thick chill. Overall non-descript as a unit.																
198.25	200.00	Trachyte	As above described 185.15 to 195.10m.																
		TRACH																	
200.00		EOH	End of Hole																

PLATINEX INC.

SHINING TREE PROJECT, ONTARIO -

Grid Coordinates:		<u>UTM - NAD83 - Zo</u>	one 17
		N: 5270976	E: 482983
Dip:	-45	Elevation: 401	m
Azimuth:	180	Total Depth: 152	m
Core Size:	NQ	Core Boxes: 36	3

	Down Hole	Tests	
Туре	Depth	Dip	Azimuth
EZ-Shot	30	-45.8	181.50
	60	-45.5	180.60
	90	-45.2	180.90
	120	-45.0	182.60
	150	-44.8	183.10

subtracting 10.5 degrees from the EZ shot instrument reading.

HOLE#: BP12-03

	Page	1	of	5
Date Star	ted:	March 24	, 2012 (N)	
Date Com	npleted:	March 25	, 2012 (D)	
Claim#:		4245864	(Collar) & 1	217520 (at Depth)
Contracto	or:	Laframbo	oise Drilling	9

Logged by: Dean R. Cutting, P.Geo. Sampled by: CXS

Target: IP Anomaly in the vicinity of the Beilby Lake Gold Occurrence.

_			Description					- ·											
De	pth	Rock Type	Description	Struc	ct. cor	e angl	es	Strain	Alterat	ion Charac	teristics					Samp	le Assays		
From	To			s。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t		
																			 L
0.00	1.50	Overburden	Casing left in the hole.																
		OVB																	
1.50	70.05	Gabbro	Very monotonous looking intrusive unit. Medium grey to grey-green colour.																
		GAB	Massive and generally equigranular, medium to fine grained locally. Moderately																
			to strongly magnetic throughout the unit. The top 5 m or so of the unit is quite																
			rubbly. No evidence noted of primary volcanic textures. Locally fine beige laths																
			of leucoxene notable as in the vicinity of 16 m. Locally the feldspar has a pinkish																
			tinge though most often whitish colour. Numerous carbonate stringers are																
			notable usually white or grey but occasionally with a pinkish tinge. Patchy light																
			epidote noted. Veinlets are irregularly oriented and from hairline to a few																
			centimetres thick. Occasional sections with disseminated pyrite as grains																
			through the unit. Unit lightly to moderately chloritized.																
			In the vicinity of 28 m, extensional fine "ladder" veinlets are noted associated																
			with a 2 cm thick carbonate veinlet.																
			Unit "fines" notably below 49.6 m, turns more greenish than grey and starts to																
			more resemble a mafic volcanic. No real contact noted, so likely a fining of the																
			intrusion.																
			51.90 m to 52.75 m is a healed up brecciated fracture zone at about 45° to the										K403830	51.90	52.75	0.85	0.179		
			core axis. Locally pinkish tinged but primarily grey-green in colour.																
			Carbonate/quartz flooding, trace hematite and 2% to 4% disseminated pyrite as																
			grains or small blebs.																
																			 L
70.05	74.40	Trachyte	Medium grained, massive and relatively equigranular though fines a bit locally.																 L
		TRACH	Typical trachyte unit. Greenish with a pinkish tinge in colour. Moderately	<u> </u>			I	ļ	I	<u> </u>	I	I							 I
			magnetic. Local irregular white to pink carbonate or carbonate/quartz veinlets	<u> </u>			I	ļ	I	<u> </u>	I	I							 L
		l	hairline to 4 mm thick. Upper contact at ±60° to core axis, relatively sharp with	-	I		I	I				I						l	 L
		l	pinkish carbonate veinlets associated. Lower contact again sharp with a fine	-	I		I	I				I						l	 ┣───
			"chill" of +5cm at +60° to the core axis, again associated with carbonate veinlets.	1	1	1					1								1

BP12-03

De	pth	Rock Type	Description	Struc	t. core	angle	es	Strain	Alterati	on Chara	acterist	tics				s	ample As	ssays		
From	To			s。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Trace disseminated granular pyrite locally.																	
			73.45 m to 73.55 m Very coarse grained pinkish-green mottled feldspar porphyry dyke.																	
			Contacts are at high angle to the core axis.																	
74.40	82.05	Gabbro	Massive, medium to fine grained equigranular intrusive looking unit as above described.																	
		GAB	Medium grey to grey-green colour. Moderately magnetic. Irregular fine white carbonate																	
			veinlets throughout. Trace disseminated granular pyrite locally. Fine beige leucoxene laths																	
			notable locally.																	
82.05	82.95	Felsite Dyke	Very strong sharp contacts notable, upper contact at 40° to the core axis and the lower																	
		FELS	contact at 60° to the core axis. Yellow-green with an orange tinge in colour. Fine grained with																	
			fine dark specks. Unit has altered the host rocks at upper and lower contacts for several																	
			centimetres from the contacts. Very hard. No sulphides noted.																	
82.95	83.45	Gabbro	Unit as above described but a bit finer grained. Core fractured for about 1 m from 83 m to 84																	
		GAB	m. Lower contact sharp at 85° to the core axis. Beige leucoxene laths are notable.																	
83.45	85.35	Mafic Flow	Typical massive , fine grained, medium green coloured unit. Seems to be a block stuffed																	
		MFLOW	between two dykes, or some sort of xenolith. May just be a finer phase of the gabbro, hard to																	
			tell. Essentially non-magnetic. Irregular carbonate veinlets are ubiquitous. No sulphides noted.																	
85.35	87.30	Feldspar Porphyry	Green, yellow, pink mottled colour. Medium to coarse grained locally with notable pink (as																	
		FP	well as white feldspars. Upper contact seems to be sharp though rubbly at likely a high angle																	
			to the core axis. Lower contact sharp but irregular. Unit is moderately chloritized. Strongly																	
			pyritic to 5% or more locally as disseminated grains or small masses. Unit is lightly																	
			carbonatized with irregular white carbonate veinlets.																	
			Pyrite up to 5% or better locally.										K403831	86.30	87.30	1.00	0.079			
87.30	96.00	Mafic Flow	Medium to dark green colour. Generally fine grained with subround carbonate filled																	
		MFLOW	amygdules notable locally. Moderately hard. Moderately chloritized. Irregular white carbonate																	
			veinlets continually present. Moderately locally carbonatized in the vicinity of carbonate																	
			veinlets. Trace disseminated pyrite grains locally with some concentration with carbonate																	

BP12-03

Page 3 of 5

De	pth	Rock Type	Description	Struc	ct. core	angles	Strain	Alterati	on Chara	cteristi	cs				5	Sample A	ssays		
From	To			S。	Fol	Flow Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			veinlets at the upper contact. Only weakly magnetic. Lower unit contact is sharp and				1	1									1		
			irregular seeming to be at a high angle to the core axis.																
			Trace to 2% pyrite in small stringers and as disseminated grains at the upper contact.									K403832	87.30	87.80	87.80	0.008			
96.00	115.50	Gabbro	Medium to dark grey-green colour. Medium to fine grained and generally quite equigranular.																
		GAB	Massive and competent looking unit. Moderately chloritized. Locally moderately magnetic.																
			Irregular carbonate veinlets, usually white or beige in colour are less than 1 cm thick.																
			There is a ±1cm thick chill margin at the upper contact. Occasional disseminated pyrite																
			grains.																
			110.05 m to 110.12 m Minor quartz/carbonate breccia veinlet at 50° to 60° to the core axis.																
			sharp walls with some metasomatism along the walls for a couple of millimetres. Maybe																
			a little grey chloritic slip in the center. Trace to 1% disseminated granular pyritein the veinlet.																
115.50	134.85	Felsic Volcanic	Upper contact with the gabbro is sharp but very irregular as an intrusive type contact. This																
		FFLOW	unit is green-olive to yellow-green in colour. Grey-green chloritic fractures are common.																
			Very hard and glassy in appearance. Very fine grained, in fact very little graining is visible at																
			all. Much of the moderately + sericitized unit (yellowish green) at the top of the interval has																
			up to 3 to 5% pyrite as disseminated grains and small masses locally. The unit seems to																
			have been shattered and healed with Carbonate/quartz veinlets. This unit is likely the IP																
			signature target for the drill hole.																
			Unit Sampling:																
			Trace to 3+% disseminated granular pyrite in sericitized rhyolite.									K403833	116.00	117.00	1.00	< 0.005			
			Trace to 3+% disseminated granular pyrite in sericitized rhyolite.									K403834	117.00	118.00	1.00	0.012			
			Trace to 3+% disseminated granular pyrite in sericitized rhyolite.									K403835	118.00	119.00	1.00	< 0.005			
			122.05 m to 122.15 m Minor intermediate-felsic dyke as below 134.85m to 138.40m. Seems																
			to be at a high angle to the core axis. Rhyolite walls seem to be a bit cooked.																

BP12-03

Page 4 of 5

De	pth	Rock Type	Description	Struct. core angles Strain Alteration Characteristics Sample Assays																
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Unit Sampling:	1						1										
			Trace to 3+% disseminated granular pyrite with some pyrite with the chlorite in fractures.										K403836	125.00	126.00	1.00	0.008			
			Trace to 3+% disseminated granular pyrite with some pyrite with the chlorite in fractures.										K403837	126.00	127.00	1.00	<0.005			
			128.95 m to 129.25 m Minor intermediate-felsic dyke as below 134.85m to 138.40m. Seems																	
			to be at a high angle to the core axis. Felsic volcanic walls seem to be a bit cooked.																	
			131.5 m to 134.85 m Unit seems to be shattered and healed with pinkish and white																	
			carbonate/quartz and greenish chlorite. Variable mottled colour. Intensity of the sericite																	
			alteration variable over short distances. Trace to 1% disseminated granular pyrite.																	
			Sample at the lower unit contact.										K403838	133.40	134.85	1.45	<0.005			
134.85	138.40	Int/Felsic Dyke	Medium to light green colour. Massive and fine grained with fine black specks. Specks may be																	
		I/FDYKE	chloritized crystals. Moderately hard. Non-magnetic. Local pinkish carbonate veinlets with the																	
			ubiquitous irregular white ones. Not particularly deformed, so likely a late feature. Upper																	
			contact sharp at 50° to the core axis. Lower contact sharp but irregular at low angle to the																	
			core axis. No sulphides noted.																	
138.40	145.80	Felsic Volcanic	Unit as above described 115.50 m to 134.85 m except more strongly shattered and healed																	
		FFLOW	with carbonate/quartz/chlorite veinlets. Trace disseminated granular pyrite locally through																	
			the unit. Some pyrite associated with the chloritic fractures. Non-magnetic. Moderately to																	
			strongly sericitized with a strong yellowish-green colour.																	
			Unit Sampling:																	
			Sericitized Felsic Volcanic with chloritic fractures and carbonate veinlets. Tr-2% pyrite locally.	<u> </u>									K403839	143.00	144.00	1.00	0.17			
				Į																
			Sericitized Felsic Volcanic with chloritic fractures and carbonate veinlets. Tr-2% pyrite locally.	<u> </u>									K403840	144.00	145.30	1.30	<0.005			I
				Į																
				Į																
				1																1

BP12-03

Page 5 of 5

De	pth	Rock Type	Description	Struct. core angles Strain Alteration Characteristics Sample Assays																
From	To			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
145.80	152.00	Gabbro	Medium to dark green-grey colour. Massive, medium to fine grained and equigranular intrusive																	
		GAB	unit. Moderately chloritized. Moderately hard. Lightly magnetic locally. Ubiquitous irregular																	
			white carbonate veinlets noted. Leucoxene specking (beige) notable locally. The top of the unit																	
			where in contact with the rhyolite 145.80 m to 147.90 m is chewed up badly and shot with																	
			white to pinkish carbonate veinlets. Looks as though some structural movement at one point																	
			on the contact "foliation", used loosely, seems to be at high angle to the core axis.																	
			Chloritization, epidotization, and carbonatization associated with veinlets near the contact.																	
			Moving away from the contact zone the unit appears more as a typical gabbroic unit with only																	
			occasional moderate angle 2 cm thick carbonate-chlorite slips to be noted. Only trace																	
			disseminated pyrite grains noted locally through the unit.																	
152.00		EOH	End of Hole.																	

PLATINEX INC.

SHINING TREE PROJECT, ONTARIO - McBRIDE PROSPECT

		<u>UTM - NAE</u>	083 - Zone 17
	N:	5271603	E: 483777
-45		Elevation:	379 m
180		Total Depth:	146.7 m
NQ		Core Boxes:	35
	-45 180 NQ	N: -45 180 NQ	UTM - NAE N: 5271603 -45 Elevation: 180 Total Depth: NQ Core Boxes:

Type	Depth	Dip	Azimuth
EZ-SHOT	30	-45.4	178.20
	60	-45.1	179.90
	90	-44.6	180.80
	120	-44.1	181.00

Downhole azimuth readings have been corrected to true north by subtracting 10.5 degrees from the EZ shot instrument reading.

HOLE#: MP12-01

Page	1 of		3
Date Started:	March 26, 20	012 (D)	
Date Completed:	March 27, 20)12 (D)	
Claim#:	1192177		
Contractor:	Laframbois	e Drilling	9
Logged by: Typed by:	Dean R. Cut Iain Trusler	ting	Sampled by:

Target: IP Anomaly in the vicinity of the McBride Gold Occurrence. Dean R. Cutting Sampled by: Bruce Cupskey lain Trusler

De	pth	Rock Type	Description	Struc	t. cor	e ang	les	Strain	Alterati	on Charac	teristics	s				Sample	e Assays			
From	To			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
0.00	2.20	Overburden	Casing left in hole																	
		OVB																		
2.20	146.70	Syenite to	Historically referred to as gabbro with pink feldspar or pink gabbro.																	
		Syenodiorite	Think interval of syenite to syenodiorite. Variable in colour locally from																	
		SYE	pinkish with green overtone to greenish with a pink overtone depending																	
			on the proportion of feldsar/ mafics or intensity of chloritization that is																	
			from the most part equigranular and massive with grain size from medium																	
			to fine grain. Unit is generally non magnetic with only very local intervals																	
			slightly to moderatly magnetic. Unit is hard throughout being lightly able to																	
			scratch with a steel scribe. Leucoxene spotting (beige, white, and green)																	
			is common throughout the unit and variable from trace - 3%. Set of																	
			carbonate veinlets disperesed through the unit at 35 degrees to the core																	
			axis, regular and easily notable, veinlets usually less than 1cm thick.																	
			There appears to be a set at 90 degrees to the 35 degree set (conjugate)																	
			There are also a series of stockwork irregulars. Carbonate veinlets are																	
			usually white but may locally have a pinkish tinge. Very little pyrite notable																	
			through the unit. Chloritization occurs along fractures and in patches																	
			through the unit variable in size from cm to m in extent.																	
			Unit Sampling																	
			Trace pyrite with pink white carb veinlets, high angle to core axis.										K403665	16.00	17.00	1.00	<0.005			
			Trace to 1% pyrite, fracture zone with moderate + epidote/chlorite,										K403666	17.00	17.60	0.60	< 0.005			
			carbonate veinlets.																	
			Trace pyrite with pink white carb veinlets, high angle to core axis.										K403667	17.60	18.60	1.00	0.183			
			White and pink carbonate veinlets at 35 degrees to core axis, no sulphides										K403668	23.00	24.50	1.50	0.046			
			White and pink carbonate veinlets at 35 degrees to core axis with light										K403669	24.50	26.00	1.50	0.019			1
			chloritization.																	
			Occasional carbonate veinlets, no sulphides										K403670	34.00	34.50	0.50	0.008			1
			10cm chloritic/epidote slip fracture zone at 30 degrees to core axis, ladder										K403671	34.50	35.00	0.50	<0.005			—
			fracture between the 2 walls. No sulphides										K402672	25.00	26.00	1.00	<0.005			<u> </u>
		┟─────┦	Occasional carbonate verniets, two supplides										K403672	36.00	37.00	1.00	<0.005			┣──┤
			spar and bematite no subbides	-									11403073	30.00	31.00	1.00	0.01			1
			Occasional carbonate veinlets with chlorite. No sulphides			-							K403674	37.00	38.00	1.00	< 0.005			

MP12-01

De	pth	Rock Type	Description	Struc	ct. cor	e angl	les	Strain	Alterati	on Chara	acteristi	cs				5	Sample As	says		
From	To			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			High Au Standard # 17c										K403675	STD OF	REAS 17c	0.00	2.69			
			Pink and white carbonate veinlets with chloritization, trace pyrite										K403676	43.50	44.00	0.50	< 0.005			
			7cm healed fracture zone 20 degrees to core axis, pink and white carbonate,										K403677	44.00	45.00	1.00	0.008			
			epidote/chlorite, trace pyrite.																	
			Pink and white carbonate veinlets with chloritization, trace pyrite										K403678	45.00	45.50	0.50	< 0.005			
			Chloritic syenite with carbonate veinlets, chloritic/epidote fracture at 80 degrees to										K403679	50.00	51.00	1.00	< 0.005			
			core axis, more intense strain at bottom of interval																	
			Strained zone with chloritic/epidote in fracture pink carbonate veinlet stockwork at										K403680	51.00	51.50	0.50	< 0.005			
			high angle to core axis, no sulphides																	
			Closure of zone to pinkish syenite, with occasional carbonate veinlets										K403681	51.50	52.50	1.00	< 0.005			
			Unit a bit strained at 75-85 degrees to core axis. Fractures with chlorite evident as																	
			well as carbonate veining from about 47.00 to 52.00.																	
			Carbonate veinlets in pink/green syenite, trace pyrite.										K403682	57.50	59.00	1.50	0.041			
			Carbonate veinlets in pink/green syenite, trace pyrite.										K403683	59.00	60.50	1.50	0.059			
			Rubbly fracture zone with late fracures, rough and coated with carbonate/chlorite																	
			from about 62.25 to 63.35																	
			Rubbly fracture zone with rough fracture chlorite/carbonate coating of most of the																	
			fractures. Unit appears to be "pitted" all though. The 'pitting' would appear to be																	
			"plucking" of the mafics along the cleavage a fracture planes on a fine crystal level																	
			scale. Short intervals occasionally through the unit much finer grained. These may																	
			be internal "chills" as the phases were being implaced. They tend to be fine grained																	
			and pink/reddish in colour. One very notable at 117.5m depth. From 68.35 to 72.00.																	
			Stockwork carbonate veinlets some pinkish, no sulphides.										K403684	84.75	86.25	1.50	0.036			
			12 cm pink/white carbonate veinlet breccia at 45 degrees to core axis, no sulphides										K403685	94.50	95.00	0.50	0.324			
			Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403686	95.00	96.50	1.50	0.043			
			lets, no sulphides																	
			Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403687	96.50	98.00	1.50	0.08			
			lets, no sulphides																	

MP12-01

Page 3 of 3

De	pth	Rock Type	Description	Struc	ct. cor	e angle	es	Strain	Alterati	ion Chara	acterist	ics				5	Sample A	ssays		
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403688	98.00	99.50	1.50	0.019			
			lets, no sulphides																	
			Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403689	99.50	101.00	1.50	0.017			
			lets, no sulphides																	
			Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403690	101.00	102.50	1.50	0.077			
			lets, low angle "green" carbonate veinlets 15-20 degrees to core axis, no sulphides.																	
			This section sampled on report of fine sulphides by JT, DRC did not note sulphides.																	
			Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403691	110.00	111.50	1.50	< 0.005			
			Variable angles 30,40,70 and stockwork general																	
			Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403692	111.50	113.00	1.50	0.022			
			Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403693	113.00	114.50	1.50	< 0.005			
			Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403694	114.50	116.00	1.50	< 0.005			
			Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403695	116.00	117.50	1.50	< 0.005			
			Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403696	117.50	119.00	1.50	0.01			
			Stockwork carbonate veinlets, pink and white, no sulphides.										K403697	131.00	132.50	1.50	< 0.005			
			Stockwork carbonate veinlets, pink and white, no sulphides.										K403698	132.50	134.00	1.50	0.012			
			Stockwork carbonate veinlets, pink and white, no sulphides.										K403699	139.75	140.50	0.75	< 0.005			
			Blank Standard										K403700	BL	ANK	0.00	0.005			
146.70		EOH	End of hole.																	
				I	I	I	I	I								I			 	
				_	I		I	I		I						ļ	I		 	
				<u> </u>	I	I	I	I		I						I	I		 	
				1											1	1				

PLATINEX INC.

SHINING TREE PROJECT, ONTARIO - CLARKE PROSPECT

Grid Coordinates:			<u>UTM - NAD</u>	83 - Zone 1	<u>7</u>
		N:	5267930	E:	482085
Dip:	-45		Elevation:	385 m	
Azimuth:	70		Total Depth:	200 m	
Core Size:	NQ		Core Boxes:	47	

	Down Ho	le les	ts	
Туре	Depth		Dip	Azimuth
EZ-SHOT	30		-43.8	71.1
	60		-43.5	72.3
	90		-43.2	74.4
	120		-42.9	74.7
	150		-42.7	75.0
	180		-42.6	77.8
Downhole azimuth	readings have b	een corr	ected to tr	ue north by

subtracting 10.5 degrees from the EZ shot instrument reading.

HOLE#: CPK12-01

Page	1 of	8
Date Started:	2012-03-27 (N)	
Date Completed:	2012-03-28 (N)	
Claim#:	4209215	
Contractor:	Laframboise Drilli	ing
Logged by: Typed by:	Dean R. Cutting lain Trusler	Sampled

Target: IP Anomaly in the vicinity of the Clarke Gold Occurrence. Sampled by: Bruce Cupskey

De	epth	Rock Type	Description	Struc	ct. cor	re angle	es	Strain	Alterati	ion Charac	teristic	s				Sample	e Assays				
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t				
0.00	5.80	Overburden	Casing left in hole																		
		OVB	-																		1
5.80	29.75	Mafic Flow	Medium to dark green colour, locally to beige/olive where L-M epidotized/																		1
		Pillowed	sericitized and or bleached. Medium to fine grained. Pillow selveges are																		
		PMFLOW	clearly notable often with chlorite/carbonate/epidote alteration associated.																		1
			Unit is basically non-magnetic. Impression that the top may be up hole																		1
			from possible amvodules at top of pillows. Carb veinlets are in a stock-	1																	
			work pattern with thicknesses from hairline to 6 cm or so, some of the																		1
			thicker veinlets are brecciated. There are quartz carbonate veinlets pre-	1																	
			sent as well often at high angles to c/a usually white to grev colour and	1																	
			possibly finely granular, trace pyrite with some of these. Trace to 1 %	1																	
			pyrite locally disseminated through the unit and associated the car-	1																	
			bonate or guartz carbonate veinlets usually as grains or small masses.	1																	
			Poorly deveoloped strain at 30 - 40 degrees to core axis notable locally.																		
			Unit Sampling	1																	
			Light green colour, carbonate veinlets, Trace pyrite	1									K403701	5.80	7.00	1.20	< 0.005				
			Light green colour, carbonate veinlets, Trace to 1% pyrite	1									K403702	7.00	8.50	1.50	< 0.005				
			Grev green colour, fine grained, coarse trace pyrite with Quartz Veinlets	1									K403703	8.50	10.00	1.50	< 0.005				
			Trace to 2% pvrite disseminated	1																	
			Grev colour, 5 cm quartz carbonate breccia veinlet at 90 degrees to core	1									K403704	10.00	11.00	1.00	0.007				
			axis, pink carbonate in parts, grev guartz veinlets, 2-3% disseminated																		
			pyrite grains																		
			Grey colour, grey quartz carbonate veinlets with 3-4% disseminated	1									K403705	11.00	12.00	1.00	< 0.005				
			pyrite in veins & wallrock																		
			Olive colour, grey quartz carbonate veinlets with trace - 1% disseminated										K403706	12.00	13.00	1.00	< 0.005				
			pyrite in veins & wallrock																		
			Olive Colour, grey quartz carbonate veinlets, trace pyrite										K403707	13.00	14.00	1.00	0.023				
			Olive Colour, grey quartz carbonate veinlets, trace pyrite	<u> </u>	I	I	I	I		I	ļ	I	K403708	14.00	15.50	1.50	< 0.005				+
	<u> </u>		Light to medium green, carbonate veinlets, trace pyrite	<u> </u>	I	I	I	I			<u> </u>	-	K403709	15.50	17.00	1.50	< 0.005				┨───┤
			Light to medium green, carbonate veinlets, trace pyrite	I	I	I	I				I		K403710	17.00	18.50	1.50	< 0.005				
			Light to medium green, carbonate veinlets, trace pyrite			I					I		K403711	18.50	20.00	1.50	<0.005				┨───┤
			Light to medium green, carbonate veinlets, trace pyrite										K403712	20.00	21.50	1.50	<0.005				╂───┤
			Indee to 2% pyrite disseminated Grey colour, 5 cm quartz carbonate breccia veinlet at 90 degrees to core axis, pirk carbonate in parts, grey quartz veinlets, 2-3% disseminated pyrite grains Grey colour, grey quartz carbonate veinlets with 3-4% disseminated pyrite in veins & wallrock Olive colour, grey quartz carbonate veinlets with trace - 1% disseminated pyrite in veins & wallrock Olive Colour, grey quartz carbonate veinlets, trace pyrite Ught to medium green, carbonate veinlets, trace pyrite Light to medium green, carbonate v										K403704 K403705 K403706 K403707 K403708 K403709 K403710 K403712 K403713	10.00 11.00 12.00 13.00 14.00 15.50 17.00 18.50 20.00 21.50	11.00 12.00 13.00 14.00 15.50 17.00 18.50 20.00 21.50 23.00	1.00 1.00 1.00 1.50 1.50 1.50 1.50 1.50	0.007 <0.005 <0.005 0.023 <0.005 <0.005 <0.005 <0.005 <0.005				

CPK12-01

De	oth	Rock Type	Description	Struc	ct. cor	e angle	es	Strain	Alteratio	on Chara	acteristi	ics				Sa	ample Ass	ays		
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			Medium to dark green grey, trace carbonate veinlets, trace pyrite.										K403714	23.00	24.50	1.50	< 0.005			
			Medium to dark green grey, trace carbonate veinlets, trace pyrite.										K403715	24.50	26.00	1.50	< 0.005			
			Medium to dark green grey, trace carbonate veinlets, trace pyrite.										K403716	26.00	27.50	1.50	< 0.005			
			Medium to dark green grey, trace carbonate veinlets, trace pyrite, feldspar										K403717	27.50	29.00	1.50	< 0.005			
			porphyritic.																	
			Medium to dark green grey, trace carbonate veinlets, trace pyrite, feldspar										K403718	29.00	29.75	0.75	< 0.005			
			porphyritic, contact with diabase below																	
29.75	51.45	Diabase	Classic Diabase Dyke. Medium to dark green/ grey colour. Fine on top and bottom																	
		DIAB	for about 2-3 metres both top and bottom. Equigranular and massives though rubbly																	
			with chlorite coated fractures through the unit over short intervals. Only very																	
			occasional carbonate/ epidote filled fractures up to a cm thick, essentially devoid of																	
			veining. Moderately chloritized throughout. Moderately to strongly magnetic through-																	
			out. Contacts are both sharp with a cm scale chill zone. Top contact is in a rubble,																	
			lower contact is sharp at 60 - 65 degrees to the core axis. Occasional masses																	
			(small) of pyrite disseminated through the unit. Late featureNo sampling in unit.																	
51.45	78.85	Mafic Flow	Unit basically as described from 5.80 to 29.75m above the diabase.																	
		Pillowed																		
		PMFLOW																		
			Unit Sampling																	
			Medium to dark green, stockwork carbonate veinlets +/- epidote, trace to 1% pyrite										K403719	51.45	53.00	1.55	< 0.005			
			locally.																	
			Medium to dark green, stockwork carbonate veinlets +/- epidote, trace to 1% pyrite										K403720	53.00	54.50	1.50	< 0.005			
			locally.																	
			Medium to dark green, stockwork carbonate veinlets +/- epidote, trace to 1% pyrite										K403721	54.50	56.00	1.50	< 0.005			
			locally.																	
			Medium to dark green, carbonate veinlet stockwork, trace to 1% disseminated pyrite										K403722	56.00	57.50	1.50	< 0.005			
			with veinlets and disseminated locally																	
			Medium to dark green, carbonate veinlet stockwork, trace to 1% disseminated pyrite										K403723	57.50	59.00	1.50	< 0.005			
			with veinlets and disseminated locally	<u> </u>																
			Medium to dark green, carbonate veinlet stockwork, trace to 1% disseminated pyrite										K403724	59.00	60.50	1.50	< 0.005			
			with veinlets and disseminated locally	<u> </u>																
			Low Au Standard # 65a										K403725	L STD OF	REAS 65a	0.00	0.525			

CPK12-01

Page 3 of 8

De	pth	Rock Type	Description	Stru	ct. cor	re angl	les	Strain	Alterati	on Chara	acterist	ics				5	Sample A	ssays		
From	To			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	1		
			Medium to dark green, more intense carbonate veinlets, 10cm grey/white quartz							1			K403726	60.50	62.00	1.50	0.031	1		
			veinlet at 40 degrees to core axis, trace - 3% pyrite																	
			Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets										K403727	62.00	63.50	1.50	< 0.005			
			and disseminated locally.																	
			Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets										K403728	63.50	65.00	1.50	< 0.005			
			and disseminated locally.																	
			Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets										K403729	65.00	66.50	1.50	0.006			
			and disseminated locally.																	
			Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets										K403730	66.50	68.00	1.50	< 0.005			
			and disseminated locally.																	
			Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets										K403731	68.00	69.50	1.50	< 0.005			
			and disseminated locally.																	
			Unit appears to have locally an interval with a bit more notable strain (foliation) with																	
			associated carbonate veinlets with pyrite from trace - 2% locally as grains or small																	
			masses. Z-fold at approximately 40 degrees to core axis. (73.00m to 75.00m)																	
78.85	83.15	Porphyritic	Porphyritic mafic flow or poorly sorted mafic ash flow. Medium to coarse grained																	
		Mafic Flow	mafic flow, medium green speckled colour. Massive and equigranular for the most																	
		MFLOWPOR	part with occasional anhedral feldspar masses up to 2-3mm size, about 1% at best																	
			Unit is non-magnetic. Unit is chloritized as the balance of the mafic volcanics. The																	
			unit is a bit coarser as a volcanic than most other volcanics observed to date in this																	
			hole. White/beige/grey, carbonate veinlets are present as elsewhere. Trace - 2%																	
			disseminated pyrite locally. Upper contact gradational over 10cm or so. Lower																	
			contact is sharp @ high angle to core axis and irregular																	
83.15	84.25	Feldspar	Intrusive unit with chills of +/- 1cm thick top and bottom. Brown/red.green colour at																	
		Porphyry	the top turning almost an olive at the bottom where fine grained matrix is bleached/																	
		FP	sericitized more intensely. Unit finer and changes colour at about 84m to bottom with		I	Į														
			a gradational internal contact over 1cm or so. Unit has a fine grained felsic matrix																	
			with phenocrysts of feldspar up to 2mm size, phenocrysts are beige to light green																	

CPK12-01

Page 4 of 8

De	pth	Rock Type	Description	Struc	Struct. core angles Strain S _o Fol Flow Vn Intens			Alterat	ion Char	acterist	ics					Sample A	ssays			
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
			colour and import a 'speckling' to the unit. Unit is non-magnetic. There are also dark																	
			phenocrysts of more mafic material about the same size. Lower seemingly intrusive																	
			style contact sharp and irregular, likely at high angle to core axis. Very few fine																	
			carbonate veinlets except where altered in the lower 25 cm. Trace pyrite, Chalco-																	
			pyrite, (?red sphalerite) in the quartz carbonate veinlets at the lower unit contact,																	
			at 10 degrees to core axis.																	
84.25	85.65	Mafic Flow	Remains of the unit from 78.85 to 83.15																	
		Porphyritic																		
		MFLOWPOR																		
85.65	108.00	Mafic Flow	Mafic to intermediate flows with occasional pillow selveges notable. Medium to fine																	
		Pillowed	grained relatively equigranular unit. Unit is medium green in colour. Unit is essentially																	
		PMFLOW	non-magnetic. Locally a bit of strain (foliation) lightly developed at about 70 degrees																	
			to core axis. Light to medium chloritized, epidotized, and carbonatized in the vicinity																	
			of the carbonate veinlet stockwork that is pervasive through the unit. Most of the																	
			carbonate veinlets are white to grey in colour most being less than 1cm thick but																	
			occasionally up to 15 cm as aggregates, epidote is sometimes associated as is																	
			quartz. Trace pyrite as grains or small masses are notable disseminated through																	
			the unit both in the wallrock and the veinlets. Upper unit contact gradual over 5cm																	
			or so, lower contact is an intrusive contact, sharp but irregular.																	
108.00	127.70	Diabase	Typical diabase unit as described 29.75 to 51.45m. Both upper and lower contacts																	
		DIAB	are a bit rubbly but the contacts are sharp with colour and grain size variations,																	
			Upper contact at +/- 20 degrees to core axis, Lower contact at 30 degrees to core axis.																	
			Almost like "serpentine" present along some of the fractures. Epidote and																	
			chloritic greenish colourNo sampling in unit.																	

CPK12-01

Page 5 of 8

De	pth	Rock Type	Description	Struct. core angles Strain Alteration Characteristics		Sample Assays														
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t			
127.70	147.15	Mafic Flow	Mafic to intermediate flows basically as above described from 85.65 to 108.00.																	
		MFLOW	There are only a couple of features through the unit that may be pillow selveges but																	
			not definitive. Unit basically massive and meduim to fine grained locally. In the coarse																	
			sections feldspar phenocrysts are notable as well as possible carbonate filled																	
			amygdules in the more fine sections. The feldspars are often irregular agglomerates of																	
			crystals up to several mm in size as opposed to well formed individual crystals. The unit																	
			is essentially non-magnetic. The unit, as most of the volcanics in the hole, has an																	
			irregular stockwork of fine carbonate veinlets from hairline to a cm in thickness. This																	
			unit (136.00 - 143.00), does however, in addition have several irregular quartz carb-																	
			onate veinlets up to +/- 10 cm in thickness. The veinlets are irregular masses of grey																	
			/waxy to cryptocrystalline quartz with beige/white masses of carbonate intermixed.																	
			Pyrite content as grains of small masses in disseminated from trace - 3% locally.																	
			Very interesting looking veining.																	
			Unit Sampling																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403763	127.70	128.80	1.10	< 0.005			
			grains or small masses																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403764	128.80	129.90	1.10	< 0.005			
			grains or small masses																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403765	129.90	131.00	1.10	< 0.005			
			grains or small masses																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403766	131.00	132.50	1.50	< 0.005			
			grains or small masses																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403767	132.50	134.00	1.50	< 0.005			
			grains or small masses																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403768	134.00	135.10	1.10	< 0.005			
			grains or small masses																	
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403769	135.10	136.30	1.20	< 0.005			
			grains or small masses																	

CPK12-01

Page 6 of 8

De	oth	Rock Type	Description	Struct. core angles		Strain	Alteration Characteristics			CS	Sample Assays										
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t				
			Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403770	136.30	137.50	1.20	< 0.005				
			grains or small masses. Strong quartz with trace to 1% pyrite as above noted																		
			35% of interval.																		
			As it approaches the unit contact with the coarse grained 'gabbro' below it starts to																		
			get short intervals of a metre or so of coarse grained and feldspar porphyritic like																		
			the gabbro. Maybe a bit of "interfingering" but compositions seem similar, grain size																		
			is more variable. From 141.00 to 143.00m the unit is a bit "strained and veined" at																		
			about 40 degrees to core axis. Carbonatized, epidotized with quartz carbonate vein-																		
			lets with up to 3% pyrite locally.																		
			Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite.										K403771	137.50	139.00	1.50	< 0.005				
			Low Au Standard #65a										K403775	STD OF	REAS 65	0.00	0.496				
147.15	167.70	Gabbro	(Could easily be a coarse mafic volcanic flow?) (TS+WR?) Medium to dark green																		
		GAB	colour. Medium to medium/coarse grained. Posesses irregular masses of feldspar																		
			(phenocrysts) up to 1.5cm diameter as subround shapes. Unit is quite massive																		
			and relatively equigranular with short intervals coarsening or fining. Has the appear-																		
			ance of an intrusive unit. Unit is non-magnetic throughout. Locally a bit strained as																		
			153.00 to 153.50 at 30 degrees to core axis, associated with carbonate veining and																		
			trace pyrite. Upper contact is sharp at 20 degrees to core axis, lower contact is																		
			sharp at 65 degrees to core axis. Stockwork of fine carbonate veinlets hairline to																		
			1 cm throughout. Seems to be a set of carbonate veinlets at 20 degrees to core																		
			axis. Trace pyrite disseminated through the unit as grains or small																		
			masses, one clot of 2-3cm diameter noted at 160.80m. Quartz carbonate veining more																		
			intense at the upper contact down to 150m or so, veining as described in the unit																		
			above, trace - 1% pyrite locally.						-												
									-												

CPK12-01

Page 7 of 8

Dep	oth	Rock Type	Description	Struct.	. core	angle	s	Strain	Alteratio	on Chara	cteristi	ics	Sample Assays								
From	To			So F	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t				
167.70	172.10	Mafic	Mafic fragmental unit, would probably call it a lapilli tuff. Fragments are clearly																		
		Fragmental	notable and variable in size from ash to 10cm or so. Unit is medium to light green and																		
		MTUFF	very mottled in appearance. Non-magnetic. Shot with carbonate or carbonate-																		
			quartz veinlets with trace - 1% pyrite locally. Unit is locally moderately strained in the coarse																		
			tuff areas at 60 to 75 degrees to core axis. Unit is lightly to moderately pervasively																		
			carbonatized.																		
172.10	180.75	Glomero	Diabase dike with a glomeroporphyritic core. The top 1.70m and bottom 2.80 m are																		
		Porphyritic	typical med to fine grained diabase. Uniform, massive, occasional carbonate/																		
		Diabase	epidote filled fractures or chloritic coated fracture rubble all that is notable. The																		
		DIAB	entire unit is moderately to strongly magnetic. The central core of the dyke is glomeropheric																		
			with large 0.5 to 3cm subround feldspathic masses like cherries in a pudding.																		
			These glomeropheric masses have been sausseritized to a striking green/yellow																		
			colour. This may be used as a sort of marker unit. Upper contact is sharp at 60 degrees																		
			to core axis, lower contact sharp at 35 degrees to core axisNo sampling in unit.																		
180.75	184.00	Mafic Flow	Mafic volcanic unit. Medium to fine grained. Medium to dark green mottled colour may																		
		Pillowed	be a pillowed unit. Carbonate/epidote veinlets are pervasive and irregularly oriented.																		
		PMFLOW	Patchy carbonatized/epidotized pervasive in and around the veinlets. Trace to 1%																		
			pyrite disseminated through the unit as grains or small masses.																		
184.00	185.10	Diabase	Fine grained magnetic diabase dike. Contacts are sharp upper contact at 60 degrees																		
		DIAB	to core axis, lower contact sharp at 60 degrees to core axis. Typical diabase unit																		

CPK12-01

Page 8 of 8

De	pth	Rock Type	Description	Stru	ct. cor	e ang	les	Strain	Alteratio	on Chara	acterist	tics	Sample Assays								
From	То			So	Fol	Flow	/ Vn	Intens	Туре	Intens	%QC\	/%Py	Sample	From	То	Width	Au g/t				
185.10	195.10	Mafic Flow	Mafic volcanic flow with pillow selveges notable locally. Variable fine to medium																		
		Pillowed	grained. Non-magnetic. Numerous quartz carbonate and carbonate veinlets in a																		
		PMFLOW	stockwork pattern. Trace - 1% disseminated pyrite locally through the unit. Moder-																		
			ately pervasively carbonatized through the unit. Maybe a bit of light strain locally at																		
			40 degrees to core axis.																		
			Irregular carbonate veinlets with trace to 1% pyrite locally.										K403812	192.50	194.00	1.50	< 0.005				
			Irregular carbonate veinlets with trace to 1% pyrite locally.										K403813	194.00	195.10	1.10	< 0.005				
195.10	195.55	Argillite	Argillite unit fine grained, dark grey/black to medium grey/green. Finely banded at																		
		ARG	60 degrees to core axis. Crenulated silica bands. Pyrite blebs along foliation/bedding																		
			planes. Dark section is slightly conductive. Locally carbonatized with a few carbonate																		
			veinlets through the unit. Overall 3-5% disseminated pyrite mostly with the																		
			fine grained dark section.																		
			Unit Sample										K403814	195.10	195.55	0.45	0.023				
195.55	200.00	Mafic Flow	As 185.10 to 195.10. Coarsening with local phenocryts from 198.1 to bottom of the																		
		Pillowed	hole. Locally a bit like the porphyritic gabbro (coarse flow?) at 147.15 to 167.70.																		
		PMFLOW	Light irregular carbonate veinlets throughout. Trace to 1% pyrite disseminated																		
			locally.																		
			Unit Sampling																		
			Tr to 1% disseminated pyrite.										K403815	195.55	197.00	1.45	< 0.005				
			Tr to 1% disseminated pyrite.										K403816	197.00	198.50	1.50	< 0.005				
200.00		EOH	End of Hole																		
					1																
					1		1														

Appendix II

Certificates of Analysis



Certificate of Analysis

Work Order: SU19451

Date: Apr 19, 2012

To: James Trusler Platinex Inc. 445 Apple Creek Blvd Markham ONTARIO L3R 9X7

P.O. No.	:	Batch: K403551-K403624
Project No.	:	-
No. Of Samples	:	74
Date Submitted	:	Apr 09, 2012
Report Comprises	:	Pages 1 to 3
		(Inclusive of Cover Sheet)

Distribution of unused material:

Store for 90 days.Instructions to follow: Comments:

Preparation of samples was performed at the SGS Sudbury site

Certified By

Bruce Robertson **Operations Manager**

SGS Minerals Services (Toronto) is accredited by Standards Council of Canada (SCC) and conforms to the requirements of ISO/IEC 17025 for specific tests as indicated on the scope of accreditation to be found at http://www.scc.ca/en/programs/lab/mineral.shtml

Report Footer:

L.N.R. = Listed not received = Not applicable

n.a.

1.S. = Insufficient Sample = No result

*INF = Composition of this sample makes detection impossible by this method

M after a result denotes ppb to ppm conversion, % denotes ppm to % conversion

Methods marked with an asterisk (e.g. *NAA08V) were subcontracted

Methods marked with the @ symbol (e.g. @AAS21E) denote accredited tests

This document is issued by the Company under its General Conditions of Service accessible at http://www.sgs.com/terms and conditions.htm. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

SGS

Final: SU19451 Order: Batch: K403551-K403624

Element	WtKg	Au
Method	WGH79	FAA313
Det.Lim.	0.001	5
Units	kg	ppb
K403551	1.242	<5
K403552	1.809	<5
K403553	2.295	<5
K403554	2.228	<5
K403555	2.111	<5
K403556	1.964	<5
K403557	2.198	<5
K403558	2.357	<5
K403559	2.216	<5
K403560	1.665	9
K403561	1.957	36
K403562	2.322	5
K403563	2.245	8
K403564	1.955	7
K403565	1.949	9
K403566	1.487	<5
K403567	2.163	6
K403568	1.975	.18
K403569	3.424	<5
K403570	3,284	<5
K403571	2.991	12
K403572	2.195	<5
K403573	1.758	7
K403574	2.755	14
K403575	0.055	494
K403576	1.496	. 7
K403577	3.197	5
K403578	3.048	5
K403579	3.387	<5
K403580	3.165	<5
K403581	3.163	<5
K403582	3.275	<5
K403583	3.249	<5
K403584	3.359	11
K403585	3.483	<5
K403586	3.344	8
K403587	3.259	10
K403588	3.255	12
K403589	3,076	5
K403590	3.378	10
K403591	2.992	18
K403592	3,232	5
K403593	3.275	15

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sqs.com/terms and conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

SGS

Final : SU19451 Order: Batch: K403551-K403624

Element	WtKg	Au
Method	WGH79	FAA313
Det.Lim.	0.001	5
Units	ĸg	jaqq
K403594	3.268	5
K403595	3.235	
K403596	3.225	11
K403597	3.290	6
K403598	3.311	<5
K403599	3.294	<5
K403600	0.055	4620
K403601	3.168	8
K403602	3.200	9
K403603	2.910	8
K403604	2,650	· 9
K403605	2.040	6
K403606	1.901	<5
K403607	2.889	6
K403608	3.219	5
K403609	3.105	12
K403610	3.210	13
K403611	3.166	6
K403612	3.142	10
K403613	3.319	9
K403614	2.212	8
K403615	2.474	65
K403616	2.913	18
K403617	2.104	8
K403618	3.112	9
K403619	3.368	<5
K403620	3.358	6
K403621	3.318	16
K403622	3.356	5
K403623	3.221	<5
K403624	3.237	<5
*Rep K403579		<5]
*Rep K403613		10
en en fan it en een een een eerste eerste kerste eerste eerste eerste eerste eerste eerste eerste eerste eerste	a ara ana ara marana faranar ara ana ana ana an	·

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sgs.com/terms_and_conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

22

SG

Certificate of Analysis

Work Order: SU19452

Apr 17, 2012 Date:

To: James Trusler Platinex Inc. 445 Apple Creek Blvd Markham ONTARIO L3R 9X7

P.O. No.	:	Batch: K403625-K403664
Project No.	:	-
No. Of Samples	:	40
Date Submitted	:	Apr 09, 2012
Report Comprises	:	Pages 1 to 2
		(Inclusive of Cover Sheet)

Distribution of unused material:

Store for 90 days.Instructions to follow: Comments:

Preparation of samples was performed at the SGS Sudbury site

Certified By

Bruce Robertson Operations Manager

SGS Minerals Services (Toronto) is accredited by Standards Council of Canada (SCC) and conforms to the requirements of ISO/IEC 17025 for specific tests as indicated on the scope of accreditation to be found at http://www.scc.ca/en/programs/lab/mineral.shtml

Report Footer:

L.N.R. = Listed not received n.a. = Not applicable

= Insufficient Sample LS. = No result

*INF. = Composition of this sample makes detection impossible by this method

M after a result denotes ppb to ppm conversion, % denotes ppm to % conversion

Methods marked with an asterisk (e.g. *NAA08V) were subcontracted

Methods marked with the @ symbol (e.g. @AAS21E) denote accredited tests

This document is issued by the Company under its General Conditions of Service accessible at http://www.sgs.com/terms_and_conditions.htm. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law,

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com



Final : SU19452 Order: Batch: K403625-K403664

Floment	WłKa	Διι
Method	WGH79	FAA313
Det.I im.	0.001	5
Units	kg	ppb
K403625	0.735	<5
K403626	3.566	<5
K403627	3.308	<5
K403628	3.136	<5
K403629	3.164	<5
K403630	3.317	<5
K403631	. 3.179	<5
K403632	2,998	<5
K403633	3.214	<5
K403634	3.282	<5
K403635	3,366	<5
K403636	3.180	<5
K403637	2.699	<5
K403638	3.666	<5
K403639	2.697	<5
K403640	2.253	<5
K403641	2.317	<5
K403642	3.460	<5
K403643	3.446	<5
K403644	3.481	<5
K403645	1.295	<5
K403646	3.463	<5
K403647	1,555	<5
K403648	3.048	5
K403649	1.845	9
K403650	0.067	523
K403651	1,972	18
K403652	2.411	7
K403653	2.706	<5
K403654	3.326	<5
K403655	2,230	<5
K403656	2.012	<5
K403657	1.983	<5
K403658	2.483	<5
K403659	2.269	<5
K403660	2.446	<5
K403661	2.833	<5
K403662	2.204	<5
K403663	2.306	<5
K403664	1.176	<5
*Rep K403643		<5

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sgs.com/terms_and_conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com



Certificate of Analysis

Work Order: SU19656

Date: May 03, 2012

To: James Trusler Platinex Inc. 445 Apple Creek Blvd Markham ONTARIO L3R 9X7

> P.O. No. : Batch: B Project No. : -No. Of Samples : 36 Date Submitted : Apr 20, 2012 Report Comprises : Pages 1 to 2 (Inclusive of Cover Sheet)

Distribution of unused material: STORE: Comments:

Preparation of samples was performed at the SGS Sudbury site

Certified Bv :

Bruce Robertson Operations Manager

SGS Minerals Services (Toronto) is accredited by Standards Council of Canada (SCC) and conforms to the requirements of ISO/IEC 17025 for specific tests as indicated on the scope of accreditation to be found at http://www.scc.ca/en/programs/lab/mineral.shtml

Report Footer:

L.N.R. = Listed not received n.a. = Not applicable

ble

I.S. = Insufficient Sample -- = No result

*INF = Composition of this sample makes detection impossible by this method

M after a result denotes ppb to ppm conversion, % denotes ppm to % conversion

Methods marked with an asterisk (e.g. *NAA08V) were subcontracted

Methods marked with the @ symbol (e.g. @AAS21E) denote accredited tests

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sqs.com/terms_and_conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com



Final : SU19656 Order: Batch: B

Element	WtKg	Au
Method	WGH79	FAA313
Det.Lim.	0.001	5
Units	kgį	ppb
K403665	2.150	<5
K403666	1.401	<5
K403667	2.453	183
K403668	3.259	46
K403669	3.243	19
K403670	1.149	8
K403671	1.260	<5
K403672	2,306	<5
K403673	2.084	10
K403674	2.152	<5
K403675	0.062	2690
K403676	1.067	<5
K403677	2.172	8
K403678	1.166	<5
K403679	2.153	<5
K403680	1.080	<5
K403681	2.014	<5
K403682	3.197	41
K403683	3.378	59
K403684	3,362	36
K403685	1.008	324
K403686	3.217	43
K403687	3.285	80
K403688	3.290	19
K403689	3.178	17
K403690	3.281	77
K403691	3.245	<5
K403692	3.311	22
K403693	3.391	<5
K403694	3.022	<5
K403695	· 3.385	<5
K403696	3.388	10
K403697	3.230	<5
K403698	3.651	12
K403699	1.707	<5
K403700	0.755	5
*Rep K403668		45

This document is issued by the Company under its General Conditions of Service accessible at http://www.sgs.com/terms_and_conditions.htm. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

Member of the SGS Group (Société Générale de Surveillance)

SGS

Certificate of Analysis

Work Order: SU19657

Date: May 03, 2012

To: James Trusler Platinex Inc. 445 Apple Creek Blvd Markham ONTARIO L3R 9X7

P.O. No.	:	Batch: C
Project No.	:	-
No. Of Samples	:	31
Date Submitted	:	Apr 20, 2012
Report Comprises	:	Pages 1 to 2
		(Inclusive of Cover Sheet)

Distribution of unused material: STORE: Comments:

Preparation of samples was performed at the SGS Sudbury site

Certified By

Bruce Robertson Operations Manager

SGS Minerals Services (Toronto) is accredited by Standards Council of Canada (SCC) and conforms to the requirements of ISO/IEC 17025 for specific tests as indicated on the scope of accreditation to be found at http://www.scc.ca/en/programs/lab/mineral.shtml

Report Footer:

L.N.R. = Listed not received n.a. = Not applicable I.S. = Insufficient Sample -- = No result

*INF = Composition of this sample makes detection impossible by this method

M after a result denotes ppb to ppm conversion, % denotes ppm to % conversion

Methods marked with an asterisk (e.g. *NAA08V) were subcontracted

Methods marked with the @ symbol (e.g. @AAS21E) denote accredited tests

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sgs.com/terms_and_conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

Member of the SGS Group (Société Générale de Surveillance)



Final: SU19657 Order: Batch: C

Element	WtKg	Au
Method	WGH79	FAA313
Det.Lim.	0.001	5
Units	kg	ppb
K403701	2.734	<5
K403702	3.309	<5
K403703	3.311	<5
K403704	2.171	7
K403705	2.292	<5
K403706	2.303	<5
K403707	2.283	23
K403708	3.592	<5
K403709	3.242	<5
K403710	3.442	<5
K403711	3.654	<5
K403712	3.508	<5
K403713	3.261	<5
K403714	3.621	<5
K403715	3.063	<5
K403716	3.428	<5
K403717	3.272	<5
K403718	1.700	<5
K403719	3.655	<5
K403720	3.610	<5
K403721	3.679	<5
K403722	3.702	<5
K403723	3.722	<5
K403724	3.573	<5
K403725	0.063	525
K403726	3.418	31
K403727	3.506	<5
K403728	3.363	<5
K403729	3,905	6
K403730	3.417	<5
K403731	3.884	<5
*Rep K403712		<5

Page 2 of 2

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sgs.com/terms_and_conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the Client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

Certificate of Analysis

Work Order: SU19926

Date: May 22, 2012

To: James Trusler Platinex Inc. 445 Apple Creek Blvd Markham ONTARIO L3R 9X7

P.O. No. : Batch: D Project No. : -No. Of Samples : 16 Date Submitted : May 14, 2012 Report Comprises : Pages 1 to 2 (Inclusive of Cover Sheet)

Distribution of unused material: STORE:

Comments:

Preparation of samples was performed at the SGS Sudbury site

Certified By :

Bruce Robertson Operations Manager

SGS Minerals Services (Toronto) is accredited by Standards Council of Canada (SCC) and conforms to the requirements of ISO/IEC 17025 for specific tests as indicated on the scope of accreditation to be found at http://www.scc.ca/en/programs/lab/mineral.shtml

Report Footer:

L.N.R. = Listed not received n.a. = Not applicable I.S. = Insufficient Sample - = No result

*INF = Composition of this sample makes detection impossible by this method

M after a result denotes ppb to ppm conversion, % denotes ppm to % conversion

Methods marked with an asterisk (e.g. *NAA08V) were subcontracted

Methods marked with the @ symbol (e.g. @AAS21E) denote accredited tests

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sgs.com/terms_and_conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com

SGS

Final: SU19926 Order: Batch: D

Element	WtKg	Au
Method	WGH79	FAA313
Det.Lim.	0,001	5
Units	кg	add
K403763	2.710	<5
K403764	2,818	<5
K403765	2.609	<5
K403766	3.727	<5
K403767	3.709	<5
K403768	2.687	<5
K403769	2.802	<5
K403770	2.802	<5
K403771	3.639	<5
K403775	0.063	496
K403812	3.280	<5
K403812DUP	<0.001	<5
K403813	2,380	<5
K403814	0.958	23
K403815	3.699	<5
K403816	3.807	<5
*Rep K403770		<5

This document is issued by the Company under its General Conditions of Service accessible at <u>http://www.sgs.com/terms and conditions.htm</u>. Attention is drawn to the limitation of liability, indemnification and jurisdiction issues defined therein.

WARNING: The sample(s) to which the findings recorded herein (the "Findings") relate was (were) drawn and / or provided by the Client or by a third party acting at the Client's direction. The Findings constitute no warranty of the sample's representativity of the goods and strictly relate to the sample(s). The Company accepts no liability with regard to the origin or source from which the sample(s) is/are said to be extracted. The findings report on the samples provided by the client and are not intended for commercial or contractual settlement purposes. Any unauthorized alteration, forgery or falsification of the content or appearance of this document is unlawful and offenders may be prosecuted to the fullest extent of the law.

SGS Canada Inc. Mineral Services 1885 Leslie Street Toronto ON t(416) 445-5755 f(416) 445-4152 www.ca.sgs.com



1046 Gorham Street Thunder Bay, ON Canada P7B 5X5 Tel: (807) 626-1630 Fax: (807) 622-7571 www.accurassay.com assay@accurassay.com

Monday, November 28, 2016

Final Certificate

Platinex Inc Suite 807 20 Williams Roe Blvd Newmarket, ON, Can L3Y 5V6 Ph#: (416) 565-5616 Email: drcutting@yahoo.ca, jtrusler@platinex.com

Date Received: 11/25/2016 Date Completed: 11/28/2016 Job #: 201655646 Reference: Core RUSH Sample #: 23

Acc #	Client ID	Au g/t (ppm)	
100500	K403818	0.015	
100501	K403819	<0.005	
100502	K403820	<0.005	
100503	K403821	0.036	
100504	K403822	<0.005	
100505	K403823	0.034	
100506	K403824	<0.005	
100507	K403825	3.095	
100508	K403826	0.070	
100509	K403827	0.016	
100510	K403827 Dup	0.012	
100511	K403828	0.032	
100512	K403829	0.027	
100513	K403830	0.179	
100514	K403831	0.079	
100515	K403832	0.008	
100516	K403833	<0.005	
100517	K403834	0.012	
100518	K403835	<0.005	
100519	K403836	0.008	
100520	K403837	<0.005	
100521	K403837 Dup	<0.005	
100522	K403838	<0.005	
100523	K403839	0.017	
100524	K403840	<0.005	

APPLIED SCOPES: ALP1, ALFA1

ch Supervisor

Validated By:

Guillaume Bouffard, Lab

Certified By: Mever hele Nathalie Neve atory Manager - Rouyn Lat

Authorized By:

Derek Demianiuk, VP Quality

The results included on this report relate only to the items tested.

The Certificate of Analysis should not be reproduced except in full, without the written approval of the laboratory.



1046 Gorham Street Thunder Bay, ON Canada P7B 5X5 Tel: (807) 626-1630 Fax: (807) 622-7571 www.accurassay.com assay@accurassay.com

Monday, November 28, 2016

Final Certificate

Platinex Inc Suite 807 20 Williams Roe Blvd Newmarket, ON, Can L3Y 5V6 Ph#: (416) 565-5616 Email: drcutting@yahoo.ca, jtrusler@platinex.com

Control Standards

QC Type

Element

Au

QC Performance (ppm)

0.580

Job #: 201655646 Reference: Core RUSH Sample #: 23

Date Received: 11/25/2016

Date Completed: 11/28/2016

Std Dev (ppm) 0.050

APPLIED SCOPES: ALP1, ALFA1

Validated By:

Guillaume Bouffard, Lab Tech Supervisor

Certified By: lever Nathalie Neveau

Authorized By:

Mean (ppm)

0.640

Derek Demianiuk, VP Quality

The results included on this report relate only to the items tested. The Certificate of Analysis should not be reproduced except in full, without the written approval of the laboratory.

PLATINEX INC. SHINING TREE PROJECT, ONTARIO - CLARKE PROSPECT

Grid Coordinates:	UTM Nad	83 Zone 17	
LAT/LONG Coordinates:		N: 5267930	E: 482085
Dip:	-45	Elevation:	385
Azimuth:	70	Total Depth:	200 m
Core Size:	NQ		

Туре	Depth		Dip	Azimuth
EZ-SHOT	30		-43.8	71.1
	60		-43.5	72.3
	90		-43.2	74.4
	120		-42.9	74.7
	150		-42.7	75.0
	180		-42.6	77.8
Downhole azimuth subtracting 10.5 d	n readings have egrees from the	e been co e EZ shot	rrected to t instrumen	true north by t reading.

HOLE#: CPK12-01

Page	1	of	11
Date Started:	March 27	, 2012	
Date Completed:	March 28	, 2012	
Claim#:	4209215	5	
Contractor:	Laframb	oise Drillir	ng
Logged by:	D. Cutting	g	Sample

Target:

De	pth	Rock Type	Description	Struc	Struct. core angles			Strain	Alteratio	teration Characteristics		;	Sample Assays					
From	То			S₀	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
0.00	5.80	Overburden	Casing left in hole															
		OVB																
5.80	29.75	Mafic Flow	Medium to dark green colour, locally to beige/olive where L-M epidotized/															
		Pillowed	sericitized and or bleached. Medium to fine grained. Pillow selveges are															
		PMFLOW	clearly notable often with chlorite/carbonate/epidote alteration associated.															
			Unit is basically non-magnetic. Impression that the top may be up hole															
			from possible amygdules at top of pulnes?. Carb veinlets are in a stock-															
			work pattern with thicknesses from hairline to 6 cm or so, some of the															
			thicker veinlets are brecciated. There are quartz carbonate veinlets pre-															
			sent as well often at highs to core axis usually white to grey colour and															
			possibly finely grained?, trace pyrite with some quartz then trace - 1 to 6															
			pyrite locally disseminated through the unit and associated with it. Car-															
			bonate and quartz carbonate veinlets usually as grains or small masses.															
			Poorly deveoloped strain at 30 - 40 degrees to core axis notable locally.															
			Unit Sampling															
		K403701	Light green colour, carbonate veinlets, Trace pyrite										K403701	5.80	7.00	1.20	<5	
		K403702	Light green colour, carbonate veinlets, Trace to 1% pyrite										K403702	7.00	8.50	1.50	<5	
		K403703	Grey green colour, fine grained, coarse trace pyrite with Quartz Veinlets										K403703	8.50	10.00	1.50	<5	
			Trace to 2% pyrite disseminated															
		K403704	Grey colour, 5 cm quartz carbonate breccia veinlet at 90 degrees to core										K403704	10.00	11.00	1.00	7	
			axis, pink carbonate in parts, grey quartz veinlets, 2-3% disseminated															

CPK12-01

De	pth	Rock Type	Description	Strue	Struct. core angles			Strain	Alterati	on Chara	acterist	ics				S	ample Ass	says
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		K403714	Medium to dark green grey, trace carbonate veinlets, trace pyrite.										K403714	23.00	24.50	1.50	<5	
		K403715	Medium to dark green grey, trace carbonate veinlets, trace pyrite.										K403715	24.50	26.00	1.50	<5	
		K403716	Medium to dark green grey, trace carbonate veinlets, trace pyrite.										K403716	26.00	27.50	1.50	<5	
		K403717	Medium to dark green grey, trace carbonate veinlets, trace pyrite feldspar										K403717	27.50	29.00	1.50	<5	
			porphyritic.															
		K403718	Medium to dark green grey, trace carbonate veinlets, trace pyrite feldspar										K403718	29.00	29.75	0.75	<5	
			porphyritic, contact with diabase below															
29.75	51.45	Diabase	Classic Diabase Dyke. Medium to dark green/ grey colour. Fine on top and bottom															
		DIAB	for about 2-3 metres top and bottom. Equigranular and massives though nubby															
			with chlorite crated fractures through the unit over short intervals. Only very															
			occasional carbonate/ epidote filled fractures up to a cm thick, essentially devoid of															
			veining. Moderately chloritized throughout. Moderately to strongly magnetic through-															
			out. Contacts are both sharp with a cm scale chill zone. Top contact is in a rubble,															
			lower contact is sharp at 60 - 65 degrees to the core axis. Occasional masses															
			(small) of pyrite disseminated through the unit, Late featureNo sampling in unit.															
		Mafic Flow	Unit basically as described from 5.80 to 29.75m above the diabase.															
		Pillowed																
51.45	78.85	PMFLOW																
			Unit Sampling															
		K403719	Medium to dark green, stockwork carbonate veinlets +/- epidote, trace to 1% pyrite										K403719	51.45	53.00	1.55	<5	
			locally.															
		K403720	Medium to dark green, stockwork carbonate veinlets +/- epidote, trace to 1% pyrite										K403720	53.00	54.50	1.50	<5	
			locally.															
		K403721	Medium to dark green, stockwork carbonate veinlets +/- epidote, trace to 1% pyrite										K403721	54.50	56.00	1.50	<5	
			locally.															
		K403722	Medium to dark green, carbonate veinlet stockwork, trace to 1% disseminated pyrite										K403722	56.00	57.50	1.50	<5	
			with veinlets and disseminated locally															
		K403723	Medium to dark green, carbonate veinlet stockwork, trace to 1% disseminated pyrite										K403723	57.50	59.00	1.50	<5	
			with veinlets and disseminated locally															
		K403724	Medium to dark green, carbonate veinlet stockwork, trace to 1% disseminated pyrite										K403724	59.00	60.50	1.50	<5	

CPK12-01

Page 3 of 11

De	pth	Rock Type	Description	Struct. core angles			Strain	Alterat	ion Chara	cterist	ics				S	ample As	says
From	То			S。	S _o Fol Flow Vn I		Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		K403726	Medium to dark green, more intense carbonate veinlets, 10cm grey/white quartz									K403726	60.50	62.00	1.50	31	
			veinlet at 40 degrees to core axis, trace - 3% pyrite														
		K403727	Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets									K403727	62.00	63.50	1.50	<5	
			and desseminated locally.														
		K403728	Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets									K403728	63.50	65.00	1.50	<5	
			and desseminated locally.														
		K403729	Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets									K403729	65.00	66.50	1.50	6	
			and desseminated locally.														
		K403730	Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets									K403730	66.50	68.00	1.50	<5	
			and desseminated locally.														
		K403731	Medium to dark green, carbonate veinlet stockwork, trace - 1% pyrite with veinlets									K403731	68.00	69.50	1.50	<5	
			and desseminated locally.														
73.00	75.00		Unit appreas to have locally an interval with a bit more notable strain (foliation) with														
			associated carbonate veinlets with pyrite from trace - 2% locally as grains or small														
			masses. Z-fold at approximately 40 degrees to core axis.														
																	1
		K403732	Medium to dark green, carbonate veinlets, trace to 1% disseminated pyrite locally									K403732	69.50	71.00	1.50		
			with veinlets and in the wallrock														
		K403733	Medium to dark green, carbonate veinlets, trace to 1% disseminated pyrite locally									K403733	71.00	72.00	1.00		
			with veinlets and in the wallrock														
		K403734	Medium to dark green, carbonate veinlets, trace to 1% disseminated pyrite locally									K403734	72.00	73.00	1.00		
			with veinlets and in the wallrock														
		K403735	Please see description above for intersection 73.00 to 75.00m.									K403735	73.00	74.00	1.00		
		K403736	Please see description above for intersection 73.00 to 75.00m.									K403736	74.00	75.00	1.00		
		K403737	Medium green colour, white/grey carbonate veinlet stockwork, trace pyrite and									K403737	75.00	76.00	1.00		1
			disseminated with veinlets and desseminated.														
		K403738	Medium green colour, white/grey carbonate veinlet stockwork, trace pyrite and									K403738	76.00	77.00	1.00		
			disseminated with veinlets and desseminated.														
		K403739	Medium green colour, white/grey carbonate veinlet stockwork, trace pyrite and									K403739	77.00	78.00	1.00		
			disseminated with veinlets and desseminated.														
PLATINEX INC. SHINING TREE PROJECT, ONTARIO

CPK12-01

Page 4 of 11

De	pth	Rock Type	Description	Struc	ct. core	e angles	s	Strain	Alteratio	on Chara	cterist	ics					Sample As	ssays
From	То			S。	Fol	Flow Vr	n Ir	ntens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		K403740	Medium green colour, white/grey carbonate veinlet stockwork, trace pyrite and	1									K403740	78.00	78.85	0.85		
			disseminated with veinlets and desseminated.															
78.85	83.15	Porphyritic	Porphyritic mafic flow or poorly sorted mafic ash flow. Medium to coarse grained															
		Mafic Flow	mafic flow, medium green speckled colour. Massive and equigranular for the most															
		MFLOWPOR	part with occasional anhedral feldspar masses up to 2-3mm size, about 1% at best															
			Unit is non-magnetic. Unit is chloritized as the balance of the mafic volcanics. The															
			unit is a bit corser as a volcanic than most other volcanics observed to date in this															
			hole. White/beige/grey, carbonate veinlets are present as elsewhere. Trace - 2%															
			disseminated pyrite locally. Upper contact gradational over 10cm or so. Lower															
			contact is sharp @ high angle to core axis and irregular															
			Unit Sampling															
		K403741	Medium to light green, Cg?, carbonate veinlet stockwork with trace -1% diss-										K403741	78.85	80.30	0.45		
			eminated pyrite with veinlets and disseminated.															
		K403742	Medium to light green, Cg?, carbonate veinlet stockwork with trace -1% diss-										K403742	80.30	81.75	1.45		
			eminated pyrite with veinlets and disseminated.															
		K403743	Medium to light green, Cg?, carbonate veinlet stockwork with trace -1% diss-										K403743	81.75	83.15	1.40		
			eminated pyrite with veinlets and disseminated.															
83.15	84.25	Feldspar	Intrusive unit with chills of +/- 1cm thick top and bottom. Brown/red.green colour at															
		Porphyry	the top turning almost an olive at the bottom where fine grained matrix is bleached/															
		FP	Sericitized more intensely. Unit finer and changes colour at about 84m to bottom with															
			a gredational interval contact over 1cm or so. Unit has a fine grained felsic matrix															
			with phenocrysts of feldspar up to 2mm size, phenocrysts are beige to light green															
			colour and import a 'spackling' to the unit. Unit is non-magnetic. There are also dark															
			phenocrysts of more mafic material about the same size. Lower seaming intrusive															
			style contact sharp and irregular, likely at high angle to core axis. Very little fine															
			carbonate veinlets except where altered in the lower 25 cm. Trace pyrite, Chalco-															
			pyrite, (?red sphalerite) in the quartz carbonate veinlets at the lower unit contact,															
			at 10 degrees to core axis.															
		K403744	Unit Sample										K403744	83.15	84.25	1.10		

Page 5 of 11

De	pth	Rock Type	Description	Struc	ct. cor	e angl	es	Strain	Alterati	on Chara	acterist	ics					Sample A	ssays
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
84.25	85.65	Mafic Flow	Remains of the unit from 78.85 to 83.15															
		Porphyritic																
		MFLOWPOR																
		K403745	Unit Sample - Trace pyrite disseminated and with carbonate veinlets										K403745	84.25	85.65	1.40		
85.65	108.00	Mafic Flow	Mafic to intermediate flows with occasional pillow selveges notable. Medium to fine															
		Pillowed	grained relatively equigranular unit. Unit is medium green in colour. Unit is essentially															
		PMFLOW	non-magnetic. Locally a bit of strain (foliation) lightly developed at about 70 degrees															
			to core axis. Light to medium chloritized, epidotized, and carbonatized in the vicinity															
			of the carbonate veinlets stockwork that is pervasive through the unit. Most of the															
			carbonate veinlets are white to grey in colour most bein less than 1cm thick but															
			occasionally up to 15 cm as aggregates, epidote is sometimes associated as is															
			quartz. Trace pyrite as grains or small masses are notable disseminated through															
			the unit both in the wallrock and the veinlets. Upper unit contact gradual over 5cm															
			or so, lower contact is an intrusion contact sharp but irregular.															
			Unit Sampling															
		K403746	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403746	85.65	86.75	1.10		
			disseminated pyrite locally, ??															
		K403747	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403747	86.75	87.90	1.15		
			disseminated pyrite locally, ??															
		K403748	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403748	87.90	89.00	1.10		
			disseminated pyrite locally, ?? Carbonate up?															
		K403749	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403749	89.00	90.50	1.50		
			disseminated pyrite locally, ??															
		K403750	High Au Standard #61a										K403750	STAN	DARD	0.00		
		K403751	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403751	90.50	92.00	1.50		
			disseminated pyrite locally, ??															
		K403752	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403752	92.00	93.50	1.50		
			disseminated pyrite locally, ??	I	I	I	I			<u> </u>								<u> </u>
		K403753	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403753	93.50	95.00	1.50		
			disseminated pyrite locally, ??	I	I	I	I			<u> </u>								<u> </u>
		K403754	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403754	95.00	96.50	1.50		
			disseminated pyrite locally, ??															

Page 6 of 11

De	pth	Rock Type	Description	Struc	ct. cor	e angle	es	Strain	Alterati	on Chara	acterist	ics					Sample A	ssays
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		K403755	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403755	96.50	98.00	1.50		
			disseminated pyrite locally, ?? Carbonate up ?															
		K403756	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403756	98.00	99.50	1.50		
			disseminated pyrite locally, ?? Carbonate up ?															
		K403757	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403757	99.50	101.00	1.50		
			disseminated pyrite locally, ?? Carbonate up ?															
		K403758	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403758	101.00	102.50	1.50		
			disseminated pyrite locally, ?? Carbonate up ?															
		K403759	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403759	102.50	104.00	1.50		
			disseminated pyrite locally, ??															
		K403760	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403760	104.00	105.50	1.50		
			disseminated pyrite locally, ??															
		K403761	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403761	105.50	107.00	1.50		
			disseminated pyrite locally, ??															
		K403762	Medium green, carbonate stockwork veinlets with epidote and quartz, trace - 1%										K403762	107.00	108.00	1.00		
			disseminated pyrite locally, ??															
108.00	127.70	Diabase	Typical diabase unit as described 29.75 to 51.45m. Both upper and lower contacts															
		DIAB	are a bit rubbly but the contacts are sharp with colour and grain size variations,															
			unconformity at +/- 20 degrees to core axis, loaclly at 30 degrees to core axis.															
			Almost like "serpentine" growing along some of the fractures. Epidote and															
			chloritic colourNo sampling in unit.															
127.70	147.15	Mafic Flow	Mafic to intermediate flows basically as above described from 85.65 to 108.00.															
		MFLOW	There are only a couple of features through the unit that may be pillow selveges but															
			not definitive. Unit basically massive and meduim to fine grained locally. In the coarse															
			sections feldspar phenocrysts are notable as well as possible carbonate filled															
			amygdules in the more fine sections. The feldspar are often irregular agglomerate of															
			xtah? something several mm in size as opposed to well formed xtah?. The unit is															
			essentially non-magnetic. The unit as most of the volcanics in the hole, has an						ļ									
			irregular stockwork of fine carbonate veinlets from hairline to a cm in thickness. This															
			unit (136.00 - 143.00) does however in addition have several irregular quartz carb-															
			onate veinlets up to +/- 10 cm in thickness. The veinlets are irregular masses of grey															
			/waxy to crypto? crystalline quartz with beige/white masses of carbonate itermixed.															

Page 7 of 11

De	pth	Rock Type	Description	Struc	ct. cor	e angl	es	Strain	Alterati	on Chara	cterist	ics					Sample A	Assays
From	То			So	Fol	Flow	٧n	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
			Pyrite contens as grains of small masses in disseminated from trace - 3% locally.															
			Very interesting veining :)															
			Unit Sampling															
		K403763	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403763	127.70	128.80	1.10		
			grains or small masses															
		K403764	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403764	128.80	129.90	1.10		
			grains or small masses															
		K403765	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403765	129.90	131.00	1.10		
			grains or small masses															
		K403766	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403766	131.00	132.50	1.50		
			grains or small masses															
		K403767	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403767	132.50	134.00	1.50		
			grains or small masses															
		K403768	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403768	134.00	135.10	1.10		
			grains or small masses															
		K403769	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403769	135.10	136.30	1.20		
			grains or small masses															
		K403770	Medium green, carbonate stockwork veinlets, trace and disseminated pyrite as										K403770	136.30	137.50	1.20		
			grains or small masses. Strong quartz with trace to 1% pyrite as above noted															
			35% of interval :)															
			As it approaches the unit contact with the coarse grained 'gabbro' below it starts to															
			get short intervals of a metre or so of coarse grained and feldspar porphyritic like															
			the gabbro. Maybe a bit of "interfingering" but compositions seem similar, grain size															
			is more variable. From 141.00 to 143.00m the unit is a bit "strained and woven?" at															
			about 40 degrees to core axis. Carbonatized, epidotized with quartz carbonate vein-															
			lets with up to 3% pyrite locally.															
		K403771	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite.										K403771	137.50	139.00	1.50		
		K403772	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite.										K403772	139.00	140.00	1.00		
		K403773	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite. Picking up intensity										K403773	140.00	141.00	1.00		
			of quartz veinlets with pyrite.															
																		1

Page 8 of 11

De	pth	Rock Type	Description	Stru	ct. cor	e angl	es	Strain	Alterati	ion Chara	acterist	ics					Sample A	Assays
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		K403774	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite. Picking up intensity										K403774	141.00	142.00	1.00		
			of quartz veinlets with pyrite.															
		K403775	Blank Standard										K403775	STAN	DARD	0.00		
		K403776	Trace carbonate veinlets in porphyritic volcanic unit, 2% pyrite. Picking up intensity										K403776	142.00	143.00	1.00		
			of quartz veinlets with pyrite.															
		K403777	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite.										K403777	143.00	144.50	1.50		
		K403778	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite.										K403778	144.50	146.00	1.50		
		K403779	Trace carbonate veinlets in porphyritic volcanic unit, trace pyrite.										K403779	146.00	147.15	1.15		
147.15	167.70	Gabbro	(Could easily be a coarse mafic volcanic flow?) (TS+WR?) Medium to dark green															
		GAB	colour. Medium to medium/coarse grained. Posesses irregular masses of feldspar															
			(phenocrysts) up to 1.5cm 0?? As irregular subround shapes. Unit is quite massive															
			and relatively equigranular with short intervals coarsening or fining. Has the appear-															
			ance of an intrusive unit. Unit is non-magnetic throughout. Locally a bit strained as															
			153.00 to 153.50 at 30 degrees to core axis, associated with carbonate veining and															
			trace pyrite. Upper contact is sharp at 20 degrees to core axis, lower contact is															
			sharp at 65 degrees to core axis. Stockwork of fine carbonate veinlets is hairline to															
			1 cm throughout. Seems to be a set of carbonate veinlets at 20 degrees to core															
			axis. Trace sulphie pyrite as disseminated through the unit as grains or small															
			masses, one clow of 2-3cm 0?? Noted at 160.80m. Quartz carbonate veining more															
			intense at the upper contact down to 150m or so, veining as described in the unit															
			above, trace - 1% pyrite locally.															
			Unit Sampling															
		K403780	Quartz carbonate veinlets with trace to 1% pyrite locally.										K403780	147.15	148.35	1.20		
		K403781	Quartz carbonate veinlets with trace to 1% pyrite locally.										K403781	148.35	149.55	1.20		
		K403782	Quartz carbonate veinlets with trace to 1% pyrite locally.										K403782	149.55	150.75	1.20		
		K403783	Light carbonate veinlets with trace pyrite locally.										K403783	150.75	152.00	1.25		
		K403784	Light carbonate veinlets with trace pyrite locally.										K403784	152.00	153.00	1.00		
		K403785	20 degrees (something) with carbonate veinlets and trace pyrite										K403785	153.00	153.50	0.50		
		K403786	Light carbonate veinlets with trace pyrite locally.						ļ				K403786	153.50	155.00	1.50		l
		K403787	Light carbonate veinlets with trace pyrite locally.	1	I				ļ				K403787	155.00	156.50	1.50		I
		K403788	Light carbonate veinlets with trace pyrite locally.						ļ				K403788	156.50	158.00	1.50		
		K403789	Light carbonate veinlets with trace pyrite locally.										K403789	158.00	159.30	1.30		

PLATINEX INC. SHINING TREE PROJECT, ONTARIO

CPK12-01

Page 9 of 11

De	pth	Rock Type	Description	Strue	ct. cor	e angl	es	Strain	Alterati	on Chara	acterist	ics					Sample A	ssays
From	То	21		So	Fol	Flow	Vn	Intens	Type	Intens	%QCV	%Py	Sample	From	То	Width	Aug/t	
		K403790	Light carbonate veinlets with trace pyrite locally.									Ĺ	K403790	159.30	160.50	1.20		
		K403791	Light carbonate veinlets with trace pyrite locally with pyrite blebs										K403791	160.50	161.00	0.50		
		K403792	Light carbonate veinlets with trace pyrite locally.										K403792	161.00	162.50	1.50		
		K403793	Light carbonate veinlets with trace pyrite locally.										K403793	162.50	164.00	1.50		
		K403794	Light carbonate veinlets with trace pyrite locally.										K403794	164.00	165.50	1.50		
		K403795	Light carbonate veinlets with trace pyrite locally.										K403795	165.50	167.00	1.50		
		K403796	Light carbonate veinlets with trace pyrite locally.										K403796	167.00	167.70	0.70		
167.70	172.10	Mafic	Mafic fragmented unit, would probably call it a lapilli tuff. Fragments are clearly															
		Fragmented	notable and variable in size from ash to 10cm or so. Unit is medium to light green and															
		MTUFF	very mottled in appearance. Non-magnetic. Shot? With carbonate or carbonate															
			quartz veinlets with trace - 1% pyrite locally. Unit is locally not strained in the coarse															
			tuff areas at 60 to 75 degrees to core axis. Unit is lightly to moderatly pervasively															
			Carbonatized.															
			Unit Sampling															
		K403797	Quartz carbonate or Cixedity? With trace - 1% disseminated pyrite										K403797	167.70	168.80	1.10		
		K403798	Quartz carbonate or Cixedity? With trace - 1% disseminated pyrite										K403798	168.80	169.70	0.90		
		K403799	Quartz carbonate or Cixedity? With trace - 1% disseminated pyrite										K403799	169.70	170.50	0.80		
		K403800	Low Au Standard #65a										K403800	STAN	DARD	0.00		
		K403801	Quartz carbonate or Cixedity? With trace - 1% disseminated pyrite										K403801	170.50	171.30	0.80		
		K403802	Quartz carbonate or Cixedity? With trace - 1% disseminated pyrite										K403802	171.30	172.10	0.80		
172.10	180.75	Glomero	Diabase dike with a glomeroporphyritic core. The top 1.70m and bottom 2.80 m are															
		Porphyritic	typical med to fine grained diabase. Uniform, massive, occasional carbonate/															
		DIAB	epidote filled fractures or chloritic coated fracture rubble all that is notable. The															
			entire unit is mal-style? Magnetic. The central core of the dyke is glomeropheric															
			with large 0.5 to 3cm subround feldspathic masses like cherries in a pudding.															
			There glomeropheric masses have been sausseritized to a strikey green/yellow															
			colour. This may be used as a sort of mark unit. Unconformity is sharp at 60 degrees															
			to core axis, L/c sharp at 35 degrees to core axisNo sampling in unit.															
												1						

Page 10 of 11

De	pth	Rock Type	Description	Struc	t. cor	e angle	es	Strain	Alterati	on Chara	acterist	ics					Sample A	ssays
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
180.75	184.00	Mafic Flow	Mafic volcanic unit. Medium to fine grained. Medium to dark green muttled colour may					1	1	1								
		Pillowed	be a pillowed unit. Carbonate/epidote veinlets are pervasive and irregularly oriented.															
		PMFLOW	Patchy carbonatized/epidotized pervasive in and around the veinlets. Trace to 1%															
			pyrite disseminated through the unit as grains or small masses.															
			Unit Sampling															
		K403803	Irregular carbonate veinlets with trace to 1% pyrite locally.										K403803	180.75	182.00	1.25		
		K403804	Irregular carbonate veinlets with trace to 1% pyrite locally.										K403804	182.00	183.00	1.00		
		K403805	Irregular carbonate veinlets with trace to 1% pyrite locally.										K403805	183.00	184.00	1.00		
184.00	185.10	Diabase	Fine grained magnetic diabase dike. Contacts are sharp upper contact at 60 degrees															
		DIAB	to core axis, lower contact sharp at 60 degrees to core axis. Typical diabase unit															
		K403806	Unit Sample - sampled for continuity										K403806	184.00	185.10	1.10		
185.10	195.10	Mafic Flow	Mafic volcanic flow with pillow selveges notable locally. Variable fine to medium															<u> </u>
		Pillowed	grained. Non-magnetic. Numerous quartz carbonate and carbonate veinlets in a															<u> </u>
		PMFLOW	stock work pattern. Trace - 1% disseminated pyrite locally through the unit. Moder-															<u> </u>
			ately pervasively carbonatized through the unit. Maybe a bit of light strain locally at															<u> </u>
			40 degrees to core axis.															<u> </u>
			Unit Sampling															<u> </u>
		K403807											K403807	185.10	186.50	1.40		<u> </u>
		K403808											K403808	186.50	188.00	1.50		<u> </u>
		K403809											K403809	188.00	189.50	1.50		<u> </u>
		K403810											K403810	189.50	191.00	1.50		<u> </u>
		K403811											K403811	191.00	192.50	1.50		<u> </u>
		K403812											K403812	192.50	194.00	1.50		<u> </u>
		K403813											K403813	194.00	195.10	1.10		<u> </u>
																		<u> </u>
195.10	195.55	Argillite	Argillite unit fine grained, dark grey/black to medium grey/green. Finely banded at															
		ARG	60 degrees to core axis. Crenulated silica bands. Pyrite blebs along foliation/bedding															
			Planer. Dark section is conductive a bit. Locally carbonatized and a few carbonate															
			veinlets through the unit. Overall 3-5% disseminated pyrite mostly with the fg dark															I

PLATINEX INC. SHINING TREE PROJECT, ONTARIO

CPK12-01

Page 11 of 11

De	pth	Rock Type	Description	Struc	t. cor	e angl	es	Strain	Alterati	on Chara	cterist	ics					Sample A	Assays
From	То			So	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		continued	section.	1														
		K403814	Unit Sample										K403814	195.10	195.55	0.45		
195.55	200.00	Mafic Flow	As 185.10 to 195.10. Coarsening with local phenocryts from 198.1 to bottom of the															
		Pillowed	hole. Locally a bit like the porphyritic gabbro (coarse flow?) at 147.15 to 167.70.															
		PMFLOW	Light irregular carbonate veinlets throughout. Trace to 1% pyrite disseminated															
			locally.															
			Unit Sampling															
		K403815											K403815	195.55	197.00	1.45		
		K403816											K403816	197.00	198.50	1.50		
		K403817											K403817	198.50	200.00	1.50		
200.00		EOH	End of Hole															
		-																
				1	1	l	l			1		l						
					Ĩ													
				1	1	l	l			1		l						
				1	1													

PLATINEX INC. SHINING TREE PROJECT, ONTARIO - MCBRIDE PROSPECT

Grid Coordinates:		UTM Nad	83 Zone 17
LAT/LONG Coordinates:		N: 5271603	E: 483777
Dip:	-45	Elevation:	379
Azimuth:	180	Total Depth:	146.7 m
Core Size:	NQ		

	Down H	ole Tes	sts	
Туре	Depth		Dip	Azimuth
EZ-SHOT	30		-45.4	178.20
	60		-45.1	179.90
	90		-44.6	180.80
	120		-44.1	181.00
Downhole azimuth subtracting 10.5 d	n readings have egrees from the	e been co e EZ shot	rrected to t instrumen	true north by it reading.

HOLE#: MP12-01

Page	1	of	3
Date Started:	March 26	, 2012	
Date Completed:	March 27	, 2012	
Claim#:	1192177		
Contractor:	Laframbo	oise Drillin	g
Logged by:	D. Cutting	I	Sample

Target:

De	pth	Rock Type	Description	Struc	ct. cor	e ang	les	Strain	Alterati	ion Charac	teristics	5				Sampl	e Assays	
From	То			S₀	Fol	Flow	٧n	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
0.00	3.00	Overburden	Casing left in hole															
		OVB																
2.20	146.70	Syenite to	Historically referred to as gabbro or pink feldspar or pink gabbro.															
		Syenodiorite	Think interval of syenite to syenodiorite. Variable in colour locally from															
		SYE	pinkish with green overtone to greenish with a pink overtone depending															
			on the proportion of feldsar/ matus? Of intensity of chloritization that is															
			from the most part equigranular and massive with grain size from medium															
			to fine grain. Unit is generally non magnetic with very intervals being															
			slight to moderatly magnetic. Unit is hard throughout being light, able to															
			scratch with a steel scribe. Leucoxene spotting (beige, white, and green)															
			is common throughout the unit and variable from trace - 3%. Set of															
			carbonic veinlets disperesed through the unit at 35 degrees to the core															
			axis, regular and easily notable, veinlets usually less than 1cm thick.															
			There appears to be a set at 90 degrees to the 35 degree set (conjugate)															
			There are also a series of stockwork irregulars. Carbonic veinlets are															
			usually white but may locally have a pinkish tinge. Very little pyrite notable															
			through the unit. Chloritization occurs along fractures and in patches															
			through the unit variable in size from cm to m in extent.															
			Unit Sampling															
		K403665	Trace pyrite with pink white carb veinlets, high angle to core axis.										K403665	16.00	17.00	1.00	<5	
		K403666	Trace to 1% pyrite, fracture zone with veinlets? epidote/chlorite,										K403666	17.00	17.60	0.60	<5	

MP12-01

Page 2 of 3

De	pth	Rock Type	Description	Struc	t. core	e angle	es	Strain	Alterati	on Chara	cterist	ics				s	ample As	ssays
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	%Py	Sample	From	То	Width	Au g/t	
		K403675	High Au Standard # 17c	1									K403675	STAN	DARD	0.00	2690	
		K403676	Pink and white carbonate veinlets with chloritization, trace pyrite										K403676	43.50	44.00	0.50	<5	
		K403677	7cm healed fracture zone 20 degrees to core axis, pink and white carbonate,							1			K403677	44.00	45.00	1.00	8	
			epidote/chlorite, trace pyrite.															
		K403678	Pink and white carbonate veinlets with chloritization, trace pyrite										K403678	45.00	45.50	0.50	<5	
		K403679	Chloritic syenite with carbonate veinlets, chloritic/epidote fracture at 80 degrees to										K403679	50.00	51.00	1.00	<5	
			core axis, more intense strain at bottom of interval															
		K403680	Strained zome with chloritic/epidote in fracture pink carbonate veinlet stockwork at										K403680	51.00	51.50	0.50	<5	
			high angle to core axis, no sulphides															
		K403681	Closure of zone to pinkish syenite, with occasional carbonate veinlets										K403681	51.50	52.50	1.00	<5	
47.00	52.00		Unit a bit strained at 75-85 degrees to core axis. Fractures with chlorite evident as															
			well as carbonate veining.															
		K403682	Carbonate veinlets in pink/green syenite, trace pyrite.										K403682	57.50	59.00	1.50	41	
		K403683	Carbonate veinlets in pink/green syenite, trace pyrite.										K403683	59.00	60.50	1.50	59	
62.25	63.35		Rubbly fracture zone with late fracures, rough and coated with carbonate/chlorite															
			from about 62.25 to 63.35															
68.35	72.00		Rubbly fracture zone with rough fracture chlorite/carbonate coating of most of the															
			fractures. Unit appears to be "pitted" all though the 'pitting' would appear to be															
			"plucking" of the mafics along the cleavage a fracture planes on a fine xtad? Label															
			scale. Short intervals occasionally through the unit much finer grained. These may															
			be intervals "chills" as the placers were being replaced. They tend to be fine grained															
			and pink/reddish in colour. One very notable at 117.5m depth.															
		K403684	Stockwork carbonate veinlets some pinker, no sulphides										K403684	84.75	86.25	1.50	36	
		K403685	12 cm pink/white carbonate veinlet breccia at 45 degrees to core axis, no sulphides										K403685	94.50	95.00	0.50	324	
		K403686	Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403686	95.00	96.50	1.50	43	
			lets, no sulphides															
		K403687	Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-							I			K403687	96.50	98.00	1.50	80	
			lets, no sulphides															

PLATINEX INC. SHINING TREE PROJECT, ONTARIO

MP12-01

Page 3 of 3

De	pth	Rock Type	Description	Struc	t. cor	e angle	es	Strain	Alterati	ion Chara	cteristi	cs				5	Sample A	ssays
From	То			S。	Fol	Flow	Vn	Intens	Туре	Intens	%QCV	∕%Py	Sample	From	То	Width	Au g/t	
		K403688	Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-					1					K403688	98.00	99.50	1.50	19	
			lets, no sulphides															
		K403689	Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403689	99.50	101.00	1.50	17	
			lets, no sulphides															
		K403690	Pink/white carbonate veinlet stockwork, a bit chloritized/bleached around the vein-										K403690	101.00	102.50	1.50	77	
			lets, low angle "green" carbonate veinlets 15-20 degrees to core axis, no sulphides.															
			This section sampled on report of fine sulphides by JT, DRC did no note sulphides.															
		K403691	Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403691	110.00	111.50	1.50	<5	
			Variable angles 30,40,70 and stockwork general															
		K403692	Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403692	111.50	113.00	1.50	22	
		K403693	Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403693	113.00	114.50	1.50	<5	
		K403694	Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403694	114.50	116.00	1.50	<5	
		K403695	Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403695	116.00	117.50	1.50	<5	
		K403696	Stockwork carbonate veinlets pink/white, chlorite associated locally, no sulphides										K403696	117.50	119.00	1.50	10	
		K403697	Stockwork carbonate veinlets, pink and white, no sulphides.										K403697	131.00	132.50	1.50	<5	
		K403698	Stockwork carbonate veinlets, pink and white, no sulphides.										K403698	132.50	134.00	1.50	12	
		K403699	Stockwork carbonate veinlets, pink and white, no sulphides.										K403699	139.75	140.50	0.75	<5	
		K403700	Blank Standard										K403700	STAN	DARD	0.00	5	
		EOH	End of hole															
																	<u> </u>	
				_	I	I	I					I					<u> </u>	
								ļ								<u> </u>	<u> </u>	
				_	I	I	I										<u> </u>	\vdash
				_													<u> </u>	
																	<u> </u>	
				1		1			1	1	1	1					1 '	1

DIAMOND DRILLHOLE

collar:

INCLINATION:

PAGE 1 OF

LOCATION:

· .

•

AZIMUTH:

1

LENGTH:

STARTED:	March 31,2012	PM
FINISHED:	April 4,2012	PM

BP12-02.

LOGGED BY: DR Cuiting

REMARKS:

Split by Bruce Cupskey

NTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
Fram (m)	То (тт)		% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
0.0	2.3m	Casing - Querburden 3n Casing in the hole									1
		· · · · · · · · · · · · · · · · · · ·									
2.3m	14.25	DIAR - Diabase					ų				
	<u> </u>	- Medin to dork orcen are mottled finding gress. Mediun to fine grained									i
	<u> </u>	continguiller and messive through threat. Moderately to storch			·	<u> </u>				I	
		magnetic through the wint. Rubbly at the top we shall interest of									
L		Porous sections with carb vender dissofred as at 8,5 to 8.8m depth						<u> </u>	<u> </u>		
	1	Shot-with a fini, stockwork of Carb, OK + ep venilate from hairline			<u> </u>				ļ	ļ	
	<u></u>	to 0.75 on thick. Greench chlority material along the wolld in some of the						L			
	<u>.</u>	thicken weinlets. A Semventete noted as a "presenced" orienteting il				[<u> </u>	[
,		33° to the K. Conty trace supplies of nord accompany			<u> </u>			<u> </u>		<u> </u>	
	<u> </u>	have whit contact is sharp al 55° to Yay Xenoliths of idicance					<u>.</u>	<u> </u>		Į	
	<u> </u>	in Diabase near the contact but contact nethin chilled a collect					<u> </u>	L	Ļ	Ļ	.
	<u>.</u>				<u>[</u>	<u> </u>	-		Į	<u> </u>	
	Ì	No Samphus warranted in the unit						<u> </u>	<u>.</u>	<u> </u>	
	1			· .		<u> </u>	<u> </u>	·	ļ		
	ļ					·····		· · ·	<u> </u>		
14.25	25.80	MFlaw Mafic Valcanie, Flaw			<u> </u>	<u> </u>		ļ	<u> </u>	<u> </u>	
							<u>[</u>		<u> </u>	<u> </u>	
		Fine grained, equananular, massive thank heavily yeined with a			<u></u>	ļ	<u> </u>		<u> </u>		
		stockwerks of Die C. OK + en venters. Melto de areas in chour.			1	<u> </u>	<u> </u>		<u>.</u>		
		Rubble a certionic will much fractures, was be naturally due to				1	<u>i</u>				
		discouter of sm. of the cash warrets, Sparse due on through the mut				Į	1		-	<u> </u>	
	<u> </u>	as well as a association but some of the carb/ Q leh verility son it									i
		the vendets and fine "ladden trat" precise infills, Some of the carbonale									
L	•										

. `

PAGE 2 OF

NTERS	ECTION	DESCRIPTION	SAMPLE					ASSA	rs	
From (m)	To (m)	(30 am)	% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)		
		has a publish tinger howen contact vubbly and more on color and								
		grain Size vorsance :					<u> </u>			
		Unit Sampling:								
						<u></u>				
				1/11 0 000	1-1-1	1				<u></u>
		Kuth section in peo recovery in pry Tr O/cutets	1	K403551	14.00	10.00				
	-	Tet a locale we Ok white	<u> </u>	K. HA2551	18.00	19.00	· · · ·			
			+		10.04	1				
·····		Tr+p, wy strong Q/c stockwork would's, public car's locally	· · · · · · · · · · · · · · · · · · ·	K463553	19.00	20.00				
	· ·			Viloo EE21	20.00	1.00				
·		The any , we list all were		1445227	20.00	<u></u>				
25,80	36,25	Gabbro - Maybe a cource grained flow	<u> </u>						<u> </u>	
			<u> </u>							
		Lolor may of a medin open green then the dark green of the								
		may understate above loage grained to madium grained relatively	+							
		equipmenting Throughoute Leneral Mascule Lats hat piel								
		where a charter of the second of "Patil di a the local had								
		to 1-2% subbly of acoust of small necked Francis and							·····	
		Servally show and filled will Con + chi + ep. Stock werk.			<u>-</u>		100	77		
		verwheter and locally "offset" in to a comple of cu. Toward to both a to u	let stan	the to de	elma a	ellangly	Les A.Fr	m lick	4 serie	?
				3	đ i		e	/		
		31.75-32.60 Flooded Sta breccia Zone up 60+ " Oto Snaqueuk							<u> </u>	
		one contranstized Scriptized material to strangly Buticis strong	<u></u>							
		3-51 % as Grander masses. At may have several phases as there	<u> </u>							
		are bill white veryon a and are versions my to a Throughing?	1		· · · · · · · · · · · ·	<u> </u>			[

INTERSECTION	DESCRIPTION	SAMPLE % Sulphides Number From To						ASS	AYS	
From To (m) (m)		% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
	Basical in strain noted in the valls of the veilet. seems to be a									
	prottle preak. Seen to lie at 50 +0 50 - 6 %.					· .				
	Unit Sampling									
									<u> </u>	
						· · ·	· · · · ·			ļ
	Freques Q/c vets 10/ That down by in vets and rock		K403555	26.00	27.00					
			Vuelle	<u> </u>		<u> </u>				
			K46506	27.90	29.00					
	¥/		VIIOSET	10 -0	100 00	<u> </u>				
			5405777	20.00	29.00	1				
			112/38	00 -0	20.00					
			5705270	<u>L9.00</u>	30.00	1				
			VUZERA	7.00	01-0					
			1.000	5 0,774	DL.VV	1				
	<u>k</u>		VUSSLA	21 00	2110	1				
			<u> 200,000</u>	<u></u>		1				
	See description of Oto Place Tox is for		K463561	21 az	32.60	1				
	<u></u>			, 22 (1				1
	At the the Oto Flood Too		(403562	32.60	33.60	4				
·····						ľ				- 1
-	Α		1463563	33.60	3460					
<u> </u>						1				
	ł		KH03564	34,60	35.45	I				
	zł		K403565	35.K	3625					
	1			·····	9. 9. 9.					
				**********************	3					



NTERS	ECTION	DESCRIPTION		4	SAMPLE				ASS	AYS	
From	To		% Sulphides	Number	From	To	Width	Au (mb)	1	1	
	<u> </u>				<u>fib</u>	<u>, (19</u>		رميس			
26.15	3800	Masic Notania									
-0,22	20.00										
	<u> </u>	Strend M. C. Blance Labor 18 a Hun and State March Labor									
	- ·	Provents Sinde to the Mill will along to got the sale of the birds									{
		Car saves when (Yally and the hursely and the start of the hursely and the hu		·							
	[Some seniest where it some line which are the day and				<u> </u>					
	1	in the End and and established and established and and				[
	\$	DE passific milete in la 2 cm an notable Rate in buttle fraction				<u>.</u>	1				
		his not really Shicked L-M songitization of fractioner locally					1				
		store dia a a access a small reference with 5% forchy laver coute the			····						
		are San Hard relation share with intermited direct, bird & to the					1				
		Locally some weilete to make at # 45% & man a way not be cievificant-				······································	1				
·····							1				
							ļ				
		Und-Sameline				[!				
							1				
							ļ				
		TV-2% my associated of Fraction and Q/C office, local due around a shall musice		K403566	36.25	37.00	1				194 <u>3</u> -
							l				
		- įł		K403567	37.00	38.00					
										·	
		Tr-SK - dui local		K463568	38.00	33.30	ł				
Ī											
			T								
							1				
							1				
					*****		1				
							1				

DIAMOND DRILLHOLE

BP12-02

NTER	SECTION	DESCRIPTION	SAMPLE				ASS	AYS			
From	To (m)		% Sulphides	Number	From	To	Width	Au (mb)			
					<u>X:7</u>	<u> </u>	<u><u> </u></u>	/			
38.90	41.90	Felsic Volcanics								İ	
		- medium			·					İ	
		Fine Agrained, vellow meen color, massive and equiprenula with					<u></u>			l	
		Strikure Sorieite yellows/areen colo. Relativel hast wit. Non-Mecutic				,				l	
	1	Shahatt shokwark & Constand Q/K where we an asconiated. Stalide									
		and which along frictures a se shall masses upt a and so claus hected		****							
		Encotures. Dande seliese this sectors is "strained" may on a basis & inecular									
	<u>[</u>	bottle fracture. Tr-31 subplider localh.						·			
<u> </u>	<u>.</u>										
	·						1				
		Unit Samplin									
	ļ										
							<u>.</u>				
		Tr-3% on locally up O/C ulets and 'sodis' on healed fractures		K463569	38.80	40.35	<u> </u>				
						i					
	-	X		K403570	40.35	41.90	}				
				***							10-1
							<u>[</u>	 			
41.90	43.15	Gablico					<u> </u>				
		Vint basically as 25.80 to 36.25. 10 an upper contact a mix with the					<u>.</u>				
		Selsic volcanics, lower contact is sharp but irregular, Ver, strong due						· · · · · · · · · · · · · · · · · · ·			-
		an as grains on small massers up to 5% a better back. Recommy seriestized		******							
		with a yellowish that developmen new the margine and along sou of the			***		ļ				
		fractures. Incarda QCA Carp results in stadewerk and comment		······································			ļ				
		Stron heren " Encorem " spectrue is notable						 			
				• • • • • • • • • • • • • • • • • • •			<u>.</u>				
-	-	horo 5% due on + Q/C or C could thank interroad		K403571	41.90	43.15	<u>.</u>				
	1										

.

INTERS	ECTION	DESCRIPTION	SAMPLE				ASS	AYS	<u> </u>		
From (m)	To		% Sulphides	Number	From	To	Width	Au (moh)			
					<u> </u>	(11)	(11)	μογ			
1215	OFan	Falsic Valering (Flairs)		******							
7242	0.000	Letsuc Michael i and		******					į		
	<u> </u>	Fire to making appined and had a serie of an income la serie									
		Date Elsa and while call bland a south as la loss									
	<u> </u>	which the docking the sector at not appressed or have been									
	:	trigation more marchine - rayoure what comports , stricting light								······	
		when to strange breed over. moderates services and									
		Cab Of we late the days hake as if shathered and healed									
		That south he sentend if cheel and the last southing									
		Wast contains Tan 8+2/ today of in black due stab accountering					1				
		Fondance + weilake. Non little the miner internet textures the windle									
		stacion de mix man hill. Providen Well Unit is non-monster					1				
		This with a like chose childs, anough a									
			1				1				
							1				
							1				
						<u>.</u>					
							1		·····		
·		Hrit Scholine				<u>.</u>	1				
·											
					4 						
		Ale what shale and Ton 3% as break a straight will be		Vibar 17	40.15	4415	1				-
		are our strangere white property as grains to small masses a we want		<u>*******</u>	15.12	<u> </u>	1				
		**		14-25-12	4415	4510					
* D ("# 		DIVISIO	<u> </u>		1				
		Runter N/ 11 - Db - T 2%		NUN2 KTES	LK IN	111-25	<u>.</u>				
		Drecciation 0/10 vites with prime care, white 2 gray unit In- 3 bana py	>	VIL-25-25		<u>79.22</u>	1				
		Kow ITA 4 BJA	- <i>P</i>	Vilon =11	111 12-	Maa	1				
		58 	<u> </u> [<u>K7V32+6</u>	10.35	7+.00					[
							1				ļ

PAGE 7 OF

INTERS	SECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From	To (m)		% Sulphides	Number	From	To (m)	Width (m)	Au (mb)			
		Unit Samphus Continued			<u> </u>	<u>.</u>					
		D/C veinly shockwork w/Tr-3%+ on locally 95 Aproins. Small masses on w/ veinlyts		K463571	47.00	48.50					
		1		K403578	48.50	50.00					
				K403 579	50.00	51.50					
	-	14		Viloran	ELA	52.00	<u> </u>			<u></u>	<u> </u>
		×1		<u>15405200</u>	21.50	25.00					
			<u> </u>	142581	53.00	51/00	[
				10.000	- • • • •		<u> </u>				
	-	η		K403582	54.50	56.00					
	<u></u>	Ŋ		1443 583	55.00	57.50	ļ				
		μ <u>ι</u>		K46358+	57,50	59.00	<u> </u>				
				1410-05	All marks						4
		N		<u>K465 585</u>	57.00	69,50					
<u> </u>		<i>k</i> 1	<u> </u>	WI. OKRI	LA Ca	12 00					
				00 01	3¥• • 9	96.98					
• • • • • • • • • • • • • • • • • • •	-	h		Kin3581	61.m	61.50					-
	<u>.</u>						İ				
<u></u>	· · ·	и	<u> </u>	K%3588	63.50	65.00					
						[I				
		Ŋ		K4135999	65.00	\$6.50					
agy ana Bayyperin Mart Pro s											
raak t (2002 - 1888 - 1 88	~	η		1403590	66.50	68.00					

DIAMOND DRILLHOLE #

÷

BP12-02

PAGE OF

NTERS	SECTION	DESCRIPTION (M) A 1 2 202	1		SAMPLE				ASS	AYS	
From	To	Second load of 16 Boxes on the Tabler stat to process	% Sulphides	Number	From	To	Width	Au (aph)			
							<u></u>		-		
		74m to Br. 30 m. Dickin up Datcher is sentice felcie draw what									
		is likel the pricinal com before Servicitizati to relian green									
		Randance between the pinte centra and yellow green can be sharp									
	<u> </u>	on the clouds' Cracks in the pink' felse whereas are shown soriath				ļ.					
	Į	alteration address then. Of weights up of growing processing permission both									<u></u>
L		types of Selsich. Sulphule by precent of To-38 locally as disc graves								[
		of sullingsors through the inter and for associated will the stock work	· · · · · · · · · · · · · · · · · · ·			1			<u> </u>		
		<u>Versee</u>		****				······			
	<u> </u>				·····		1				
		10000 Sampling Continues	•	••••••••••••••••••••••••••••••	:						
		Plans late deducte ultra 3% - louboas due and small march and 20 late	4	K463591	12.00	1950					
		C C C C C C C C C C C C C C C C C C C									
	· ····	γ		K463592	69.50	¥.90					
- 437-00											,
	, <u> </u>	t ₁		<u> </u>	71.00	72.50					
					<u>.</u>						
L		V)		K463594	72.50	२५ ००	<u> </u>				
	<u></u>			×	~ /					ļ	
		V/ Statin ball of son put when	4	K403545	74.00	<u> </u>			r	<u> </u>	
				16: 05Q1	7000						
				<u>K403 278</u>	-530	44.00					
		рания и правити и правити и правити и правити и правити и правити и правити и правити и правити и правити и пра		in oral	7.4 ~	at a					
		V 1 T		<u>0405217</u>	<u>++.00</u>	+2.24					
		h Vi		1.100 KDQ	10 m					<u> </u>	
				KWSJU	<u> </u>	1 24.35					
		'n		1462,599	80.00	<u>%) (n</u>				 	
		h Vl h		<u>K463 598</u> K463 599	73.SD 30.D	89,00 89,00					

MIERS	SECTION	DESCRIPTION			AMPLE				ASS/	IYS	
From (m)	To (m)		% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
		An Standard # 61d		K463600						*****	
	-	н і		K403601	81.50	83.00					
		۵ ۲		K463600	83.00	9450			Ī	1	
									Í		
		4		K403603	8450	85.90					
	Ī								li li		
85.80	137.20	Felsic Valcanics							į		
	1								1	1	
		Basical the came felsic volcanics as above described but the								Į	
		Yellow-green servicity alteraty is not as stranch developed according					}				
		may in noticity and alance Fractures in a may produce descended									
		the Solici interest which. The word is still first to medus provided, were								1	
		hard as a sourced really. Now-Magnetic, Still harden a fine archiverte								-	
		of DC/CERP/SE Sender De mainline to several an in Abirlanera									
		Rail - marchil proceed thank it with a Tay 32 beell as	<u>†</u>								
		are shall march a president with the Darker liverely area	1						Ī	1	
		They the Ald process nice not much patrille & power triteres.									
		The color up in a fallene to be a question of which although					1				
····		presence of the sourced in a primar beckered Deater. Why prover					l				
	(Laber m. Chille Darkenel Him Chroniel	1								
		<u>, an all there. Sold the there were in the second s</u>		Î			<u></u>		İ		
		Unit Sandura :	<u> </u>								
							<u></u>				
		Ola Cille and the soule of the Dick T. 27 - 1. (Right back)		KURKOU	85,90	87,00	<u>.</u>				
		WILS - WERT IN SHIRMARY WIT SETURE IN FILME IN FRINK IN TO A FILMENT IN THE SHEAP STATE IN	1	In correct of the second second							
		y I v Y		KU1 Lne	64.00	88 m			l		
		/	<u> </u>	<u>500 5010</u>	1.00	VV, ~~	i				
				Y ilas Lal	and and	<i>a</i> a					******
!	-		1.	N792 600	79.00	U100	1	1			

PAGE) OF

NTER	SECTION	DE	SCRIPTION			SAMPLE				ASS/	<u>AYS</u>]
From (m)	To (m)		· · ·	% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
	-	12 12	(Rully Serta)		K463607	89.00	90.50					
			,									
		ž	(Strange Oto Finder agel, Breecon)		1403608	90.50	92.00					
		Λ	(Bitan Ine pukal)		K403601	.92.00	43.50					
<u>`</u>	<u> </u>			<u> </u>	Vilalin	00.0-	06-14				<u>.</u> 	
		79	[Jop 13 mar pulad]		<u>16705010</u>	75,50	43.00	-				
		<i>s</i> ?	(4- cair h		1452 411	95 00	96.50			ļ	. <u></u>	
	*					12.00					i	
		ų	(String Servicitie)		K403 612	96.50	93.00	1				
		4	(Strang Sericity)		18463613	98.00	99.50					
								ļ				
		<i>b</i>	Weater Servit alou Frankere	l	K463614	99.50	100.50					
								[
	*****	2	(Tink Volume of weak sea altering)		593015	100.50	101.60	<u> </u>			1	
				<u>_</u>	Verslil	6160	102.00	 				
		*2	Westing Society along trading pulle vole]		<u>6165810</u>		103.00					
		"L	/ n	<u>_</u>	K43611	103.00	104 00					
¢(. 		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,						1				1
*****		V	/ *	1	K463618	104.00	105.50					
an (a an an an an an an an an an an an an an												
		h			K403619	105.50	104.00					
				<u> </u>	ļ			<u> </u>				
	1027	h	(Strong Serverty, Obus garcola)		<u>K403620</u>	107.00	109.50	<u> </u>				
					1			<u> </u>				
	~	k	()	_	840362)	108.50	110,00					
									1			į

PAGE // OF

INTERS	SECTION	DESCRIPTION	······		5	AMPLE				ASS	YS	
From (m)	То (тт)		%8	Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
	-	QLOC Vlets in charlingthe 10/ service in anken felor vacano The 3% to 10	zelh		K403622	10.00	111.50					
		(Ritch Servit along 1	lets and faster 1			· · · · ·						
		<u> </u>			K403623	111.50	3.00					· · ·
									· ·			
					K403624	113.00	114.50	[
ļ			BLANK		<u>K403625</u>							j
		<i>v</i> e			<u>K403626</u>	14.50	116.00					<u></u>
		4			VILONA	11/ ~~	Ma	-				
					<u>NTU-524</u>	110.00				1		
·		ъ			KURLOQ	11250	119.00					1 tor
<u> </u>	<u> </u>				1-103 02 0					Í	- Au	<u></u>
<u>}</u>	-	n Serieto inr	marin interaction		K4036A	19.00	120.50	1				A
										İ		
	-	Mal Se	with affections		K.403630	120.50	12.00					V
	1		Pric FV					ł	5 - A	ł	April	3,20
		¥			(423631	122.00	23.50				,	
								ļ				
	-	k (N-LS.	with up fred in)]	K¥63632	123.50	125.00	<u> </u>	ļ			-
		Rule -	FV (<u>.</u>				
	-	i (L Souch	-7- Frank mink FV]		<u> </u>	125.00	126.57					
			~~~~~~	i	As a las	3.00						
		И.		577 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	403634	126.50	23.0					
		$\rho$	2		VU 61194	1977	1/0 ~-					,
		L-M Serve	Luy best in Kull FM		KHOS655	25.00	107, 20					
					11.0101	No -	101	[ [				
		n (M-5 Saul	-44 mile carter -	<u></u>	510565b	(29.30	151.00	1				
		<u>by 2</u>	WK FU . I				<u>.</u>					
							i	1				í

PAGE 2 OF

NTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From (m)	To (m)		% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
										ļ	
		h (N.S. Sencite usknoten + petetas		K403637	131.00	132.50					
		the Parke FU									
		pullocter?									
		At about 132.50 n wit starting to exhibit some and developed straine			-						
		at 45-55° to the % - Rubbly Zne with Kappin (while goude) on facture ) at									
		about 133.30 to 133.50m. Some of the fragments have a bit of Ok whet						-			
		in them no sulphide noted. Fairetin drops of t not notable at									
		about 137.20 m at the same time as the vellow over sericitization					<u>.</u>				
		drops significantly is intrasity									
							<u>.</u>				
		Modente to ston socia in FU TA-18 and as graine, still sheke and g alcolots		<u>K403638</u>	32.50	34.00					
							ļ				· · · · · · · · · · · · · · · · · · ·
	1	b.		K403 639	134.00	135.20	<b>.</b>				
							<u></u>				
	. —	<u>n</u>		K403 640	135.20	136.20	ļ				
				1	1.0.1					ļ	
		<u>h</u>		<u> </u>	136.20	137.20					
							[			ļļ	ļ
		Probable and of FV units					<u> </u>				
		1		• 			<u></u>				
										l	
							ļ				
	<i>.</i>			······			; ;			<u> </u>	ļ
				, ,			İ. İ.				
							[				
										<u></u>	<u> </u>



INTERSECTION	DESCRIPTION		<u> </u>	SAMPLE				ASS	AYS	<u> </u>
From To (m) (m)	Start Documenting Boxes 33 -> 47 at the both of the hole.	% Suphides	Number	From	To	Width (m)	Au (nob)			
				<u> </u>		<u> </u>				
137.20150-00	Dabase					1	1			
		1				<u> </u>				
	Massing media to date and orean colon, encionanda for themat had						[			
	Very uniform lody w/ macherline variable for light to stame going								l	
	For top to bottom of the wint. Inequila stack werk of contrasting endote								· · · · · · · · · · · · · · · · · · ·	1
	a speciale hereiting (Bocall) through the wrist- Contacts relatively sharp		ļ			<u> </u>	ļ			<u></u>
	at high & to the cove axin. Contracte nume accigned on textured and				<u> </u>	<u>.</u>			ļ	<u> </u>
	call differences between this with and three above and below.	<u> </u>		*****				ļ	<u> </u>	<u> </u>
	(My an occasional quit of patr what brack though through the	<u> </u>						<u> </u>		<u> </u>
							<u> </u>			
	1 Unit Samolin						<u> </u>	<u> </u>		Floral 7
			WL-PRIO	1000-	100 Ja	1		<u> </u>	<u> </u>	<u> </u>
	Mpper Contact Scarfing to Close taki with scarfier		KIOS 642	131.20	130.70					
			11-96-10	VID 00	1600			<u> </u>		
	house and the second		<u>CTVUTA</u>	1110-31	100.00	1				<u> </u>
			1					<b>{</b> }		
50 m 150 he	$\Omega_{\rm abb}$					1		1	<u> </u>	<u> </u>
DAVA LEAVE	<u> </u>	-				1	1	1		<u>.</u>
	Ust Canta de met abour la L a little van Relei in Concerbra	1		-	1	1	1	1	1	1
	a but lights and and tall less manuface (non south to understa).	•				1	1	1		
	my Off C weiters notable and Thave to 2% due to noted		1	1	1					
	Have the will be the second of the second of the fille shows					l				
	to partice up and al a production. There are a few throut a year attend to be					1				
	a could can in stran glosity proved in the installed From 158 m to the						1			
	lation of the white there is some matable strand (Solicita) lichtly to						-	I		1
·····	moderates developed of 60-30° +. H. V. mu, interce of H.					-	1			
	Dise where the the light where the hand at and the	1	······································				1			

PAGE 4 OF

INTERS	SECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From (m)	To (m)		% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
		with small fragments of the will be bro progred who this with. Very									
		little not able for primary features, Irregula Q/C and C what's are notable									
	<u>.</u>	through the wait out, accelerally a speek of printy is noted in assauter.		, , ,							
		Unit Sauph at he Contacts			•				-		
	1										
		Top contact up Top		<u>K403 644</u>	150.00	151.50					
						Į					
ļ	<u>.</u>	156 There puter cadonale breeces where up 24th damps that by white		1403 65	56.00	15%.50	• •			l	
		Carb Mile The at 2 60 to 1/2 what carb what 0.50 a 30 16 1/2	<b>_</b>	ļ							
					1		2				
<u> </u>	<u> </u>	We where w/ to py, Tr-2% dry by healt,		<u>Ktizz 675.</u>	156.30	158.00			·	I	[]
ļ				ast a week	1			-		<u> </u>	<b></b>
	ļ		2	<u>5103 687</u>	158.00	159.65	<u> </u>			<u>[]</u>	<u> </u>
ļ				ļ							
	[			ļ		<u> </u>					
100 100			ļ				<u> </u>				<u> </u>
158.65	F1.75	telsic to Interreducto Knoclestic Unit					ļ	: 2.		ļ	
		see a line a line a line a line a line a line a line a line a line a line a line a line a line a line a line a	<u>]</u>	<u>.</u>	1						
·		Milel production will be bedog ispliciting introbuilded of all telt			) 		1				[·
		Sou of the paper sugarante Carle up to 10 to 15 to thick. Some		[ [							[
		of the Est weeks some be my to have the or san some of the Har Engrande								<u> </u>	
	<u> </u>	air ytals an lattellite and altered to a matellinger file manpaper tody	<u> </u>	<u> </u>							
· · · · ·		The Fibry is the well seens to 3 between 20 to 10 to 16, Whill									
· 		is cale to make grang to being matthed. Nink is now-magniture					1				
		Mate namably constantized for weak to stary bally . Mut shit will	<b>.</b>	<u></u>			; ;				
		On real of Ped when The ensure we is strong prophic would be don		<u>.</u>							
		oppras in call massing in with Suching and Finding backy can be	ļ				 				; ;
		our 10 1/2 .				<b>!</b>					
ŕ				l						<u> </u>	

PAGE 5 OF

NTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From	To		% Sulphides	Number	From	To (m)	Width	Au (mb)			
	1				<u> </u>	kin i					
		Unit Samalific									
		The second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second								i	
		Servitory filszanka Of white w To-28 a		X463648	158.65	159.95					
		33		K46364A	159.95	160.75					
		Standord # 65a		K463650							
	-	( bu pake about 1 3+ 2 g at high (+ 2)		K463651	160.75	161.60					
	<u>.</u>										
		Hish w/ 3-5+2 due fr	 	X403652	161,60	162.75					
				111-1-0		100-					
		5-10% durp, in ash		K403755	16275	163.45					
		· / · · · · · · · · · · · · · · · · · ·		Vilolai	100	18 <b>-</b> 16 - 0					
		33		157852057	163.75	162.35					<u>i</u>
				VIA	le n-	11,100				( <b> </b>	
		Lapity 124 W 2418		<u>1705 000</u>	107.75	106.00					
				VURLES	ILI US	1224					
			<u></u>	N-W-Kalta	100.12	101.14					
		5-10% AL AL A DEL		K463 6-1	14.40	16830					
	_	LANK TILL 2-3° A Flack & Fight		K403653	168.30	169.45					
	-	Frasmerke à 4a Cerbulut-1		K463659	169.45	170.60					
		jî ji		K493660	170-60	171.75					
				ć							
İ											
											[

NTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From (m)	To (m)		% Sulphides	Number	From (m)	To (m)	Width (m)	Au (ppb)			
FIR	182.40	Interected to Maga Volcances			-						
An an an an an an an an an an an an an an											
		Medin to fine grained, Medin to dark oreen in color, Variabil									
		macushi for weak to stome locally. Richard had unt.									
		Wit is mult for 125.25 to 179.50 local and continue man be a fault									
		influence but no goinge noted guistion. Shet with fine contracte veniets									
		last way little water and an where .									
		177.50 -> 182.0 Succede of cab wendets with red anophan response			·····					<u>[</u>	
		relativel, sold, sucher investige sinder party red		· · · · ·		*				Ļ	
		why we also hereafter on Classe San? Should lode who class								ļ	
		May have an arientetia of 350 - 5%. There are smell				ļ				<b>[</b> ]	
		masser of py associated with the wenderter Really suspect hereotet									
		У I I I									
										<b></b>	
		Unit Samplin ?								Į	
										<b></b>	
		Contact w/ Repodation		K46366	71.75	173.00					
					3					<u> </u>	
	-	Stockwerk decirised above		662	<u>79.50</u>	160,50				<b>[</b> ]	
										<u> </u>	
											а. — ^С
										<u>[]</u>	
						 				<u>                                     </u>	
						ļ				ļ	
										ļ	
ľ										<u> </u>	
										ļ	

PAGE 170F

NTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From	To		% Sulphides	Number	From	To (m)	Width	Au	i		
			·			(11)					
10211	iarie	Distance									
101-40	107.15	Deases.					••••••				
		Mai and A i Dan days days									
		Theam open open com, mean to time grained, equigranular and									
		massive. Essentially non-magnety, shot with megular stockwalk				·					
	· · · · · · · · · · · · · · · · · · ·	Of carb/ep reinters the hairline to 0.5 on thicle. Contacts Sharp				-					
		at 40 \$ c/a							-		
									· · · · ·		
			·								
	-				·····						<u></u>
					·····			<b> </b>			<u> </u>
								l		ļ	
185.15	145.10	Irachyty war			·····			ļ	·.		
		to We				· · · · · · · · · · · · · · · · · · ·					
		Med to fine grand, green with a punked tinget locally. Very	<b>_</b>								
		hard and hattle fracturing. Fragments noted locally up to									
		about a cn is size, chloritized. Strongly magnetic throughout		•••••				[			
		heally patches are epidotized as well as along some of						<b> </b>		<u> </u>	
,		the frectioner. Shot with stack work of cond weinherts. yen, little to									•
		no subtide noted anywhere						 		<b></b>	
- 19										<u> </u>	
								l	·.		
		Unit Sandurs'								<u> </u>	
i	1										
		Stock werk & cook would be some northead times , to pa		KH03663	186.85	187.85					
				***********************	······	Tie House		1			
		State a log of the Will Way of the to a	1	K40366U	194.50	195.00		1		İ	
		- TARMAR WAS ALL AND MAN JOUR CHAR AND IN KA	1		· · · · · · · · · · · · · · · · · · ·						
					•••••			1	i	<u> </u>	
			<u> </u>		L					ليسبب	<u>i</u>

NTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From	То		% Sulphides	Number	From	То	Width	Au			
(m)	(m)				(m)	(111)	(m)	(ppo) : 			
	100.0										
195.10	M'0.25	Diabase					· · · · · · · · · · · · · · · · · · ·				
								<u> </u>			
		Fire to median argument, massive and equisicanular. Med to								ļ	
		dark and in Kalon. Moderatch to strongly magnetic									
		through S.L. Orraicional consten veillets, Contacts are share									
		user at 40° 1 %. love means with a 1/2 cm thick					-				
		chill. Querall non-descript as a with					·				
	1	0									-
100.25	200.00	Treenty									
		De abound receibed 18515+ 19510m									
					••••••						
2000		EUN OF 1/01E			······			· · ·			
200,00	EOH	END OF HULE									
		Filling and Areas a Marca De Daivil 122								·	
·····		Finish logging and Sampling Marking @ 4115 FM MPTHY, CAR			·						
		5									
		1/1 0 1 1 0 1 5									
		K40355 > K403664 indusive = 114 Samples to be Collected									
							-				
					*****				· .		
· · · ·									······		
			I.:I			. <u> </u>		;		i	

### **DIAMOND DRILLHOLE #**

PAGE	OF
------	----

INTERS	ECTION	DESCRIPTION			SAMPLE				ASS	AYS	
From (m)	To (m)		% Sulphides	Number	From	To (m)	Width (m)	Au (pob)			
									l		
									Ì		
									ł		
	ļ							1			
	ļ										
							<u>.</u>				
				·			ļ				
							<u> </u>				
							Ļ				
							ļ	·			
				*****		- -	<u> </u>				
				******			<u> </u>				
				******************		·	ļ				
· · · · · · · · · · · · · · · · · · ·	· · · ·										
							ļ				
							<u> </u>		<u> </u>		
:											
							<b> </b>				
							1				
				*			ļ	<u> </u>			
							1	<b>.</b>			
			ļ				<b>!</b>	Į	* * *		
								ļ			
							<u>t</u>	<u> </u>			: 

## SUMMARY DIAMOND DRILL CORE LOG

DRILL HOLE NUMBER: BP12-03

PAGE OF

FROM (m)	TO (m)	ROCK TYPE	SUMMARY DESCRIPTION
0.00	1.50	DB	Casing
V100	11.50		( in the second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second second se
150	+70		Califica Pacciple Masic Valrania ?
1:20	I for		Has 5 is with a bacelle lightly recording, med to
			100 cm is view with an a walking texture
			- Of shines show sole at he drie to Se shull
			To 12 de an
			Net 1 an Pg
			at + 19 h Rubble sect a chart lu
			mit tim typen sarry where the
			- Sa 10 mill - KI 11 HG
			- on ofthe venters are ported w/ neutres
			Dil colling
			Tercy epiaste locary
			- 20 - Juli mall Of an act (50)
			ZON LEDOLE VENUEL SE PILICES (SM)
			Mon Inps Journaling.
			Starts to the a sut cour the trade sold
			SITT & Breakful trattin Low hearth wy
			33. In al theating heratite and 2-4 bain (12
			prost as grains and in the Otz
	-1.1	- 15	Dul d 1
flm	44.40	Trachyto	twish strongh magnetic, strongy tragmented
			Ula in real due 1/C @ 60° to %a and days
			y with president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president president
			In Abotilis would almost be an FP.
			TV-18 dr. es locally

## SUMMARY DIAMOND DRILL CORE LOG

DRILL HOLE NUMBER: BP12-03

PAGE ____ OF ____

FROM (m)	TO (m)	ROCK TYPE	SUMMARY DESCRIPTION
	()		
	015 2	0	
74110	K. Bole	2017	Med. to fin con Mcfillacuitor fine Gabbro
71.40	FOUL		Tiching to the year of and upsand
			The Case Usinets
	17	C-	OFF SAT DIS/ 1 - has welled
		~	000 - The Allors hav streng versely
			Acd story by to SI locally and and small storing
			May be as FP
			1 Lou(oku Soh
			Fine graund and equigrander hot & neurone por
			All with below
			Tuberelle Contact w/ will
			Trodocato
A			
		A	
115.6	134.85		Felsic volcands
			Light yellewbrees color very fine grand
			Shattened and shat with Sing stodework a Q/C willet
		2	baiding a bit widen the dim a and in the vendete.
		N/	Storn & all the both ne. He Parket to 3-5% Locals
		1//	man py to be connected by connect
134,85	1384		Totoridate / Felsie Du
1- 1.02			manadans prosec equ
122.4	NER		Ele III I II I
120.1	TJ.TJ.		1990 brand any gode but with strong weing
		V	and g but min chattened
0/=0	1.10 0-		
175.0	149.05		Sto dework Cars verster in Strated blenning chloritored
			Trpy
VIDOF	TP2 0		
175.85	15600		hasash
122	Fend		
156.0	120H		