STRATHY TOWNSHIP REPORT NO. 23

This file contains work performed by Goldfields Mining on claim:

T.44793	Hole #	1	Feb/60
		2	Feb/60
		3	Mar/60



GOLDFIELDS MINING
CORPORATION

Scale 1" = 200'

NET LAKE PROPERTY

STRATHY TWP.


Blazed to Ash
Quartz
Baseline Dwgs. 1349, 1355 & 1416
RECEIVED
MAY 13 1960
TEMISKAMING
MINING DIVISION
Note:
All measurements by stadia.
N
Next
1053'
Fd Picket
8+00S
132°20'
140°17'30"
145°15'30"
163°40'
172°33'
T44793
355'
D.D.H. #1 (F.S for #2)
(approx.)
635'
677'
675'
1235'
D.D.H #2
F.S.
Picket
D.D.H #3
1 to 2 400'
Fd Cl. Post
3-T44793
Sketch Showing
Location
D.D.H. 1, 2 & 3
NET LAKE
TOWNSHIP STRATHY
Scale: 1 inch = 200 feet

DIAMOND DRILL RECORD

PROPERTY Net Lake, Strathy Twp., Ontario. Claim T44793 HOLE NO.

SHEET NUMBER 1 SECTION FROM TO STARTED February 18, 1960

LATITUDE Station 11 00S DATUM COMPLETED February 23, 1960

DEPARTURE Line 8 75W BEARING ULTIMATE DEPTH 364'

ELEVATION Lake Level DIP collar -90°;180'-88°10';360'-89° PROPOSED DEPTH

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES		
0-75	Casing; lake bottom at 40'; overburden-mud, sand, gravel, boulders to 75'							
75.0-107.0	Andesite - fine grained, dense, greenish-grey, uniform texture and color; weakly chloritized and carbonatized; core is blocky and partly broken up, which was probably the reason why the hole was jammed with sand between 80.0-100.0 and had to be reamed. Frequent thin white or pink carbonate stringers at high angle to core. Fine disseminated pyrite throughout. Frequent irregular stringers and patches of fine grained pyrite as fracture fillings and replacement of the host rock.							
	30% massive patches and stringers of pyrite, little pyrrhotite and traces of chalcopyrite	7477	79.5	81.0				
	25% mineralization as above	7478	90.8	93.0				

NORTHERN MINER FORM 505 REV: '54

DRILLED BY *Inspiration Mining & Development Co. Ltd.* SIGNED Radmila Subotincic

DIAMOND DRILL RECORD

PROPERTY..... HOLE NO.....

SHEET NUMBER 2 SECTION FROM..... TO..... STARTED.....

LATITUDE..... DATUM..... COMPLETED.....

DEPARTURE..... BEARING..... ULTIMATE DEPTH.....

ELEVATION..... DIP..... PROPOSED DEPTH.....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES			
	91.5 - strong chloritic slip at 20 to core								
	99.5 - 107.0 decrease in chlorite alteratin, weakly brecciated and bleached along irregular fractures.								
	30% massive patches and stringers	7479	101.0	102.0					
107.0-112	Diorite dike - fine grained, medium grey, heavily carbonatized; sparse specks of disseminated pyrite. Both contacts weakly brecciated and scheared at 60° to core.								
112.0-119.2	Andesite-similar to section 75.0-107.0. Weakly brecciated and bleached along irregular fractures; weakly carbonatized; frequent thin white carbonate stringers at high angle to core. Fine disseminated pyrite throughout; scattered irregular pyrite stringers, patches, seams. 112.2-113.3, 117.5-118.7 estimated 10% fine grained pyrite with minor pyrrhotite, and sparse trace of chalcopyrite.								

NORTHERN MINER FORM 505 REV. '51

DRILLED BY..... SIGNED.....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 3 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES		
119.2-120.9	Diorite dike - as from 107.0 - 112.0. Contacts at 70° to core.							
X	119.3 - 1/4" Cb. stringer at 65° to core with pyrite crystals.							
120.9-128.5	Andesite - similar to section 112.0 - 119.2; frequent thin white Cb. stringers. Fine disseminated pyrite through; minor veinlets, patches and stringers of pyrite.							
	121.5-122.0 estimated 15% of pyrite							
	122.9-123.5 10% pink Cb. stringers as breccia cement, network of fine grained pyrite veinlets crosscutting Cb. stringers. Few specks of chalcopyrite noticed.							
128.5-138.0	Porphyritic andesite; arbitrary contacts; widely scattered greyish-white subrounded "phenocrysts" of feldspars set in fine grained grey matrix similar to the preceding andesite. "Phenocrysts" are il defined and aggregates of feldspar grains rather than individual crystals. Scattered thin white Cb stringers at high angles to core. Minor fine disseminated pyrite; occassional irregular pyrite seams, stringers or patches.							

DRILLED BY ..... SIGNED .....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 4 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES			
138.0-144.0	Andesite - fine grained, dense, medium grey-greenish, weakly carbonatized; light silicification from 138.0-142.0; fair chloritization from 142.0-144.0. Disseminated pyrite throughout; occassional irregular pyrite patch, stringer or seam.								
	139.7 - 14" Cb. stringer with crystals of pyrite at 40° to core.								
144.0-152.0	Diorite dike - as from 107.0-112.0; vague contact at 144.0 at high angle to core; lower contact obscured by broken core; sparse disseminated pyrite								
X	At 145/0-1/2" Cb. veinlet at high angle to core with pyrite inside and outside of the walls. At 146.0 - thin Cb. veinlet at high angle to core with coarse pyrite crystals.								
152.0-159.6	Andesite - as from 138.0-144.0. Fine disseminated pyrite throughout, at 152.2 153.0 and 154.2 stringers and irregular patches of fine pyrite with trace of chalcopyrite.								
	152.9 - 1/4" pink Cb. stringer with grains of pyrite and specks of chalcopyrite.								

DRILLED BY ..... SIGNED .....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 5 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES
	157.1-157.3 massive pyrite stringer with Cb. along the walls at 60° to core.					
159.6-168.0	Porphyritic andesite, as from 128.5-138.0; arbitrary contacts. Minor fine grained disseminated pyrite.					
	162.6-163.0 5% massive pyrite, minor pyrrhotite and trace of chalcopyrite as breccia cement					
	163.0-163.7 several narrow irregular pyrite stringers, minor patchy quartz material and some bleaching.					
	163.7-164.2 60% fine pyrite, minor pyrrhotite and Cb. material at high angle to core.					
	166.8-1 1/2" pyrite cemented breccia zone at 50° to core.					
168.0-197.0	Andesite - as from 138.0-144.0; moderate brecciation throughout, especially from 169.0-172.0 and 181.0-197.0. Prominent bleaching along irregular fractures, of rock fragments and as patches and streaks at moderate to low angles to core. Occassional thin discontinuous Cb. stringer at moderate angle to core. Abundant irregular streaks stringers, fracture					

DRILLED BY ..... SIGNED .....

A10

DIAMOND DRILL RECORD

PROPERTY..... HOLE NO.....

SHEET NUMBER 6 SECTION FROM..... TO..... STARTED.....

LATITUDE..... DATUM..... COMPLETED.....

DEPARTURE..... BEARING..... ULTIMATE DEPTH.....

ELEVATION..... DIP..... PROPOSED DEPTH.....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES			
	fillings and patchy replacement of the host rock by pyrite and pyrrhotite. Occassionally patchy Cb. associated with sulphides. More frequent traces of chalcopyrite. Important is the increase in pyrrhotite content, which may in places be 50% of the sulphides present.								
197.0-237.0	Andesite - as from 138.0-144.0. More or less massive. From 197.0-198.0 and 199.3-203.4 pillow structures-fine grained rims (rather vague) and coarse grained interior marked by small aggregates of dark green minerals set in lighter colored matrix. Frequent thin Cb. stringers at high angles to core. Numerous irregular pyrite and pyrrhotite and trace of chalcopyrite.stringers, streaks, irregular patches and masses, especially in slightly bleached and fractured sections such as from 198.0-198.5, 206.5-207.5, 211.7-212.6 and 230.6-237.0. Pyrrhotite content increasing.								
X	25% pyrite and pyrrhotite in brecciated and fractured section	7480	220.0	221.6					

NORTHERN MINER FORM 505 REV: '54

DRILLED BY..... SIGNED.....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 7 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES
	20% pyrite and pyrrhotite, minor Cb.	7481	225.6	226.6		
237.0-242.5	Diorite dike, as from 107.0-112.0. Fine grained contacts at high angle to core.					
242.5-248.0	Andesite, medium greyish-green, fine grained, weakly carbonatized, fair to moderate brecciation and bleaching; crisscrossed with heavy stingers and irregular patches of pyrrhotite, minor pyrite. Very sparse chalcopyrite as specks and whisps often associated with patchy Cb. material. Estimated 20% sulphides present. 247.2-248.0 core badly broken up.					
		7482	243.0	247.2		
248.0-266.0	Andesite- medium grey, fine grained, weakly carbonatized. Fine disseminated pyrite Frequent irregular fractures and short weakly brecciated and bleached sections criss-crossed and filled in by pyrrhotite and minor pyrite. Frequent irregular pyrrhotite stringers and patches. Sparse trace of chalcopyrite and pyrrhotite content increasing.					
	Estimated 15% sulphides	7483	248.0	253.0		

NORTHERN MINER FORM 505 REV: '51

DRILLED BY ..... SIGNED .....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 8 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES
	Estimated 10% sulphides	7484	253.0	256.6		
	262.6-262.8 patchy light yellowish-green Epidote alteration					
266.0-295.0	Andesite breccia - medium grey-greenish fine grained matrix and lighter grey angular fragments. Contacts indistinct. This rock has been strongly brecciated and its original character obscured. Moderate bleaching mainly of rock fragments. Weak chlorite alteration mainly of the matrix. Chlorite often present in and around fractures. Heavy pyrrhotite mineralization often along two sets of intersecting fractures at low or moderate angle. Pyrrhotite fills and replaces the wall rock of these fractures; it also occurs as dissemination or massive replacement patches. Little pyrite associated with pyrrhotite. Occassionally specks of chalcopyrite are present in pyrrhotite, but most often chalcopyrite occurs as thin seams or whisps associated with carbonate crosscutting pyrrhotite, which suggests that it is later than pyrrhotite. Few such seams are best					

DRILLED BY ..... SIGNED .....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 9 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES			
	shown from 290.0-291.0. Estimated 20-25% sulphides.								
		7485	268.0	273.0					
		7486	273.0	278.0					
		7487	278.0	283.0					
		7488	283.0	288.0					
		7489	288.0	293.0					
295.0-333.0	Andesite (?) - medium grey, medium to coarse grained. Abundant darker green chloritized mafic aggregates (1 to 4 mm) in lighter grey fine grained matrix. Weakly carbonatized. Matrix is quite light colored and the rock is more acidic in composition than normal andesite - possibly it is dacite. Minor and erratic dissemination of pyrrhotite and pyrite. Scattered stringers and irregular patchy replacements of pyrrhotite, sometimes associate with little chlorite. Pyrite content varies from low to moderate, and pyrite grain often surrounded by phyorrite.								

NORTHERN MINER FORM 565 REV: 1-54

DRILLED BY ..... SIGNED .....

A14

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 10 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES
	310.0-312.5 heavy patches and stringers of irregular Cb.-Quartz, minor chlorite, and minor blebs and crystals of pyrrhotite and pyrite.					
	320.0-333.0 increase in pyrrhotite mineralization; rock is weakly brecciated, fractured and bleached. Pyrrhotite occurs as stringers, patches and dissemination; little pyrite associated with pyrrhotite. At 324.0, 324.5 and 324.6 small whisps of chalcopyrite associated with Cb. stringers cutting across pyrrhotite. At 324.6 1/4" Cb. stringer at high angle to core with blebs and specks of chalcopyrite in and outside the vein. Estimated 15% to 20% sulphides.					
		7490	320.0	324.0		
		7491	324.0	329.0		
		7492	329.0	332.5		
333.0-364.0	Porphyritic andesite- similar to section 128.5-138.0. Contacts gradational and arbitrary (feldspar aggregates appear for the first time at about 320.0					

NORTHERN MINER FORM 506 REV: /54

DRILLED BY ..... SIGNED .....

A15

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 11 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES		
	but are very scarse and erratic). Weakly carbonatized. Very little disseminated pyrrhotite, and pyrite. Occassional pyrrhotite, (pyrite) stringers, veinlets or irregular patches, often associated with little chlorite.							
	333.0-337.0 weakly brecciated, fair pyrrhotite in patches of stringers. Minor pyrite usually surrounded by pyrrhotite.							
	339.5-342.0 weak brecciation and bleaching with scattered pyrrhotite seams and patches.							
	350.5-350.7 80% pyrrhotite veinlet with specks of chalcopyrite							
	352.7 whisp of chalcopyrite associated with small Cb. patch in massive pyrrhotite veinlet.							
	354.0-355.0 8" ground core.							
	360.4-364.0 moderate brecciation, some bleaching, scattered heavy pyrrhotite patches, seams and breccia cement.							
	363.5-1/2" Cb. stringer at 30 to core with chalcopyrite along and outside the walls							

DRILLED BY ..... SIGNED .....

DIAMOND DRILL RECORD

PROPERTY ..... HOLE NO. .....

SHEET NUMBER 12 SECTION FROM ..... TO ..... STARTED .....

LATITUDE ..... DATUM ..... COMPLETED .....

DEPARTURE ..... BEARING ..... ULTIMATE DEPTH .....

ELEVATION ..... DIP ..... PROPOSED DEPTH .....

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	ASSAY VALUES			
	within massive pyrrhotite.								
364.0	End of hole.								

NORTHERN MINER FORM 509 REV. 54

DRILLED BY ..... SIGNED .....

A(17)

DIAMOND DRILL RECORD

PROPERTY [illegible] **HOLE NO.** 2

SHEET NUMBER _____ SECTION FROM _____ TO _____ STARTED [illegible]

LATITUDE [illegible] 1447 43 DATUM [illegible] COMPLETED [illegible]

DEPARTURE [illegible] BEARING [illegible] ULTIMATE DEPTH 693'

ELEVATION [illegible] DIP -45° (300'-46°) (630'-44°) PROPOSED DEPTH 696'

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	GOLD $	SLUDGE GOLD $		
0 - 88	Casing (90° H2O, mod sand, gravel, boulders)						
88 - 410	ANDESITE						
	fine grained, light to dark grey, occ. greenish due to chlorite mineralization, occ. veinlets and stringers of pink-white carbonate sometimes with qtz, sparse qtz. veins, sparse cpy. associated with carbonate veins.						
	Veins at 40° to 50° to core with many erratic, sparse fine pyr, primary py.						
	109.5, 118, 132.5 - 133.5 carb. veins 40 - 50° to core						
	143.2 - 144.5 breccia appearance.						
	217 - 3/4 strong chloride with shearing.						
	230.5 - 232.5 qtz. vein (erratic)						
410 - 421.5	DIABASE DIKE						
	Greenish white med. grained at 65° to core						
	413 - 414.5 qtz.-carb.-diabase breccia						
	420 post dike qtz. - pink carb. vein (1")						
	421 post dike qtz. - pink carb. vein (4")						
	with massive fine pyr, sparse cpy.						

N.M.P., TORONTO—STOCK FORM No. 501 REV. 12/51

DRILLED BY [illegible] SIGNED [illegible]

DIAMOND DRILL RECORD

PROPERTY [illegible] HOLE NO. 3

SHEET NUMBER _____ SECTION FROM _____ TO _____ STARTED 24-3-60

LATITUDE Line 940 W DATUM Claim #44791 COMPLETED 4-3-60

DEPARTURE 1430 S BEARING 325° ULTIMATE DEPTH 637'

ELEVATION Lake Level DIP -45° (200'-43°) (600'-44°) PROPOSED DEPTH 650'

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	Cu [illegible]	[illegible]	Au	Ag
481.8 - 618	ANDESITE						
	As from 88 - 418 but slightly more chlorite with ph -						
	wh carb. veins at 50° to core.						
	MINERALIZATION						
	520 - 575						
	Mineralized zone of 5% - 25% pyrr. trace to one per cent						
	cpy. in siliceous andesitic matrix, except 550 - 553.8						
	in sheared chloritic matrix and 540.1 to 544.6 in						
	graphitic matrix. All zones tested positive Mi.						
	520 - 533.8	1	3.8	T	T	N	0.18
	533.8 - 534.8	2	1.0	0.16	T	N	0.07
	534.8 - 537.8	3	3.0	T	T	N	0.04
	537.8 - 539.8	4	1.9	1.00	T	N	0.03
	539.8 - 543	5	3.8	0.00	T	N	0.00
	543 - 546.8	6	3.8	0.04	T	N	0.10
	546.8 - 548.1	7	4.9	0.06	T	N	0.16
	548.1 - 544	8	3.9	0.00	T	N	0.04
	544 - 548.6	9	4.6	0.00	T	N	0.06

N.M.P., TORONTO—STOCK FORM No. 501 REV. 12/51

DRILLED BY _____ SIGNED _____

A19

DIAMOND DRILL RECORD

PROPERTY [illegible] LAKE, STRATHY TWP., ONTARIO HOLE NO. 2

SHEET NUMBER _____ SECTION FROM _____ TO _____ STARTED 24-3-60

LATITUDE 1100 S DATUM Claim #44723 COMPLETED 4-3-60

DEPARTURE 1420 E BEARING 325° ULTIMATE DEPTH 637'

ELEVATION Lake Level DIP -45° (300'-46°) (620'-44°) PROPOSED DEPTH 636'

DEPTH FEET	FORMATION	SAMPLE NO.	WIDTH OF SAMPLE	Oz. per ton	[illegible]	Au	Ag
	548.6 - 552.6	10	4.0	0.12	T	T	0.03
	552.6 - 557.6	11	5.0			T	0.16
	557.6 - 562.6	12	5.0			T	0.08
	562.6 - 567.6	13	5.0			T	0.10
612 - 615	LAMPRO PORPH DIKE						
	Greyish black, coarse grained with prominent biotite						
	mica. Dike at 40° to core.						
615 - 637	ARKOSITE						
	As prev. described (83 - 410) (421.6 - 612)						
637	END OF HOLE						
	Lost Core						
	113 - 120 (4) 120 - 122 (1) 227 - 234 (6)						
	264 - 268 (3) 272 - 276 (1) 276 - 283 (4.5)						
	283 - 289 (1.5) 402 - 408 (1) 486 - 495 (1.5)						
	495 - 500 (5) 562 - 571 (2) 571 - 573 (1)						
	Blocky Core Recovery						
	168 - 244 30%						
	244 - 312 30%						
	437 - 499 25%						

N.M.P., TORONTO—STOCK FORM NO. 501 REV. 12/51

DRILLED BY Inspiration N. & D. Co. Ltd.

SIGNED [illegible]

DIAMOND DRILL RECORD

PROPERTY [illegible] LAKE, STRATHY TOWNSHIP, ONTARIO **HOLE NO.** 3

SHEET NUMBER 1	SECTION FROM ____ TO ____	STARTED March 5, 1960
LATITUDE L[illegible] [illegible]00'	DATUM Claim #44793	COMPLETED March 14, 1960
DEPARTURE [illegible] 1400' [illegible]	BEARING 323°	ULTIMATE DEPTH [illegible]
ELEVATION Lake Level	DIP -45° ([illegible]' -51°) ([illegible]' -40°)	PROPOSED DEPTH 400

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	GOLD $	SLUDGE GOLD $		
0 - [illegible]	[illegible] ([illegible]' [illegible]), mud, sand, gravel, boulders)						
[illegible] - 2[illegible].[illegible]	Andesite						
	light gray when siliceous, greenish due to presence of chlorite, generally dark and fine grained. General presence of qtz-carb veins and stringers. The veins vary from 30° to 60° to core with erratic stringers. Cpy. occ. present with veining, sparse pyrr. and py.						
	7[illegible].5 - [illegible]1.[illegible] qtz-carb, chloritic sheared, siliceous andesitic matrix bearing sparse cpy., cpy., pyrr., py. carb. vein [illegible]0° to core vug veining and oxidation evident.						
	[illegible].5 and [illegible]7 carb-quartz veining (1/2") at 30° to core.						
2[illegible].[illegible] - [illegible].7	[illegible] Dike						
	Greenish white with post dike qtz carb vein (1/2")						

N.M.P., TORONTO—STOCK FORM NO. 501 REV. 12/51

DRILLED BY Inspiration [illegible] Co. Ltd.

SIGNED G. Ross

DIAMOND DRILL RECORD

PROPERTY _____ HOLE NO. _____

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	GOLD $	SLUDGE GOLD $		
	[illegible] at 30° to core. Fine at 65° - 90° to core. fine grained.						
[illegible]	[illegible]						
	[illegible] 1% cpy in qtz-carb siliceous andesitic matrix.						
	317.1 and 31[illegible] qtz carb veins (1/4") at 60° to core. (369.5) (377.5) (400) (408) sparse cpy. evident.						
	411 and 416 sparse cpy. pyrr. in qtz carb veins (1/2") at 60° to core.						
	419.5 barren 4" qtz-carb vein.						
	450 - 45[illegible] sparse cpy. in dark andesite.						
	452.2 (1/2") and 452.6 (3") qtz carb veins 60° to core.						
453.1 - 47[illegible]	[illegible]						
	453.1 - 4[illegible] greenish gray diorite fresh appearance with red feldspar specks.						
	456.3 - 45[illegible] greenish white diabase med. grained.						
	45[illegible] - 47[illegible] siliceous sheared andesite.						

N.M.P., TORONTO—STOCK FORM NO. 501 REV. 12/51

DRILLED BY _____ SIGNED _____

DIAMOND DRILL RECORD

PROPERTY _____ HOLE NO. _____

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	GOLD $	SLUDGE GOLD $	Au	
	474.5 - 478.5 diorite as above. med. grained also.						
478.5 - 694.3	Andesite as from (44 - 246.4)(246.7 - 456.3)						
	466 qtz-carb vein at 40° to core with erratic stringers and red feldspar mineralization.						
	(531 - 535) (537 - 540) and minor zones at depth are composed of dense to fine fine grained andesite with strong epidote alteration.						
	Mineralization						
	568 - 574 2% pyrr. sparse py., cpy., with evident layered mineral lineation at 30° - 35° to core from 570 to 572. Siliceous andesitic matrix.	25	6.0			T	
	634 - 663						
	mineralized zone of 5% to 35% pyrr. mainly in massive						

N.M.P., TORONTO—STOCK FORM No. 501 REV. 12/51

DRILLED BY _____ SIGNED _____

A25

DIAMOND DRILL RECORD

PROPERTY _____ HOLE NO. _____

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	[illegible]	[illegible]	Au		Ag.
	short sections in a siliceous andesitic matrix bearing sparse py. and negligible to 1% cpy. Trace of graphite evident.							
	634 - 637	14	3.0	0.06	T	T		0.20
	637 - 643	15	6.0	0.03	T	N		0.04
	643 - 645	16	2.0	0.25	0.01	N		0.04
	645 - 646.7	17	1.7	0.11	T =	N		0.03
	646.7 - 650	18	3.3	0.00	0.01	N		0.00
	650 - 652.5	19	2.5	0.31	T=	N		0.03
	652.5 - 655	20	2.5	0.25	T	N		N
	655 - 660	21	5.0	0.13	0.01	N		T
	660 - 665	22	5.0	0.10	0.02	N		T
	665 - 670.3	23	5.3	0.23	0.03	N		0.16
	670.3 - 675.3	24	5.0	0.03	0.01	N		0.22
	646.7 - 650 Diorite dike. med. grained. greyish white.							
604.3 - 606	Diorite dike - gray and med. grained.							

N.M.P., TORONTO—STOCK FORM No. 501 REV. 12/51

DRILLED BY _____ SIGNED _____

A24

DIAMOND DRILL RECORD

PROPERTY _____ **HOLE NO.** _____

SHEET NUMBER 5 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE No.	WIDTH OF SAMPLE	GOLD $	SLUDGE GOLD $		
695.6 - 699	Andesite						
	as before (66 - 265.6) (266.7 - 430.3) (470.5 - 694.5)						
699	END OF HOLE						

N.M.P., TORONTO—STOCK FORM No. 501 REV. 12/51

DRILLED BY _____ SIGNED G. Cass for J. A. Cass



Sketch Showing
Location
D.D.H. 1, 2 & 3
NET LAKE
TOWNSHIP STRATHY
Scale: 1 inch = 200 feet