



Diamond Drilling

Township of TYRRELL

Report No: 16

Work performed by: Getty Mines Limited

| Claim Nº | Hole Nº | Footage | Date | Note |
|----------|---------|---------|---------|------|
| L 398873 | J-75-9 | 409.01 | June/75 | (1) |
| | J-75-10 | 145,51 | July/75 | (1) |

Notes:

(1) #102-76

GETTY MINES, LIMITED

Hole Number

J-75-9

Page 1 of 2

DRILL HOLE LOG

| Property. Juby Location. Tyrrell Township, Ontario | Core Size. AQ Elev. Collar. 185° Bearing45° | Starting Date JUNE 6, 1975 Completion Date. JUNE 8, 1975 |
|--|---|--|
| Grid. 'A' Latitude. 17 + 25N Departure. 0+ 00 | Dip45° Length. 409' Horiz. Trace. 289 Vert. Trace. 209 | Logged by. Donald C. Cameron |

| Dip Tests | | | | | | | | |
|---|-------|--------|--|--|--|--|--|--|
| | Angle | | | | | | | |
| Depth | Read | Actual | | | | | | |
| Collar | | 450 | | | | | | |
| 50' | : | 42° | | | | | | |
| 3001 | | 41°30 | | | | | | |
| · | | | | | | | | |
| · • • • • • • • • • • • • • • • • • • • | | | | | | | | |

| 77.01 | то | TO DESCRIPTION | SAMPLE | FOOTAGE | | CORE | ASSAY | | | | | |
|-------|-------|----------------|--|---------|------------|-------|----------|---------|----------|----------|----------|--------------|
| FROM | | | NUMBER | FROM | TO | LGTH. | | | | | | |
| 0 | 18.0' | OVERBURDEN: | Clay, sand, boulders | | | | | | | | | |
| | | | | | | | | | | | | |
| 18.01 | 51.5' | INTERBEDDED G | REYWACKE & CHERTY SEDIMENTS: | | | | | | | | | |
| | · | | The grey wacke is medium to fine grained | | | | | | | | | |
| | | | and dark grey in color. Generally massive | | | | | | | | | 1 |
| | | | and with very minor specks of pyrite. The | | | | | | | <u> </u> | | |
| | · | | cherty sediments are generally light brown | | | | | | | | | • |
| | | | to buff in color and very fine grained. | | | | | | <u> </u> | | <u> </u> | |
| | | | Light sericitic alteration and with some | | | | | | | <u> </u> | <u> </u> | |
| | | | chloritization. Locally fractured on a | | | | | | | | | |
| | | | small scale with narrow (less than 5 mm | | | | | | | | | |
| | | | thick) quartz veins irregularly intersecting | | | | | | | <u> </u> | | |
| | | | the sedimentary units. Bedding at 45 to | | | | | | | | | |
| | | | core axis. | | | | <u> </u> | | | <u> </u> | - | - |
| | | | | | | | | | | | | 1 |
| 51.5' | 59.71 | RHYOLITE: | The unit is light brown in color, slightly | | | | | | | | ļ | |
| | | | porphyritic with rounded and diffused | | | | | | | <u> </u> | | |
| | 50.00 | | phenocrysts of quartz to 2 mm. Contact | | | | | 1 | <u> </u> | | | |
| | | | with the preceding unit is at 70° to core axis | | | | | | | | | |
| | | | The unit is massive and with very minor | | Para Maria | | | | | | | <u> </u> |
| | | | fine grained specks of pyrite (less than | | | | | | | | <u> </u> | |
| | | | 0.1%). From 52.3 to 53.7 a quartz vein | | | | | | | | | 1 |
| | | | with green specks of fuchite to about 20%, | | | | | | | | 1 | |
| | | | | · | | | | | | | <u> </u> | |
| | | | | | | | | | | | | |
| | | | | | | | | | | | 1 | |
| | | | | | | | | <u></u> | | <u> </u> | | |


GETTY MINES, LIMITED


Hole Number


7-75-9

DRILL HOLE LOG

| , | | DRILL HOLE L | <i></i> | | | | | | | | |
|--------------|-------------|---|----------|----------|---------|--------------------|-------|-----|-------|----------|----|
| ED 014 | | | SAMPLE | f | FOOTAGE | | CORE | | | ASSAY | |
| FROM | TO | DESCRIPTION | NUMBER | FROM | TO | LGTH Au oz. Ag oz. | | | | | |
| | | intersects core axis at 45°. No significan | | | | | | | | | |
| | | mineralization associated with the quartz | | | | | | | | | |
| | | vein. | | | | | | | | | |
| | | | | | | | | | | | |
| 59.71 | 93. 0' | INTERBEDDED GREYWACKE AND CHERTY SEDIMENTS: | - | | | | | | | | |
| | | As above from 18.0 to 51.5. | | | | | | | | | |
| | | | | | | | | | | | |
| 93.01 | 95.41 | FRACTURE ZONE: | | | | | | | | | |
| | | Chert and quartz veining in the fractures. | No | | | | | | | | |
| | | mineralization associated with the fractur | ng. | | | | | | | | |
| | | | | | | | | | | | |
| 95.41 | 345. 2' | GREYWACKE & CHERTY SEDIMENTS: | | | | | | | | | 3. |
| | | Generally the greywacke is confined to the | J-6853 | 232, 5 | 234.0 | 1.5 | 0.005 | | | | |
| | | upper portion of the unit (to about 115.5). | J-6854 | 238.0 | 240.5 | 2.5 | 0.02 | | · | | |
| | | The rest of the unit is essentially cherty | J-6855 | 306.4 | 310.0 | 3.6 | 0.005 | | | | |
| | | sediments. From 115.5 to 117.6 a green | | | | | | | | | |
| | | and black quartz vein with fuchite and gra- | hitic | <u> </u> | | | | | | 1 | |
| | · | containing about 20% disseminated pyrite. | | | | | | | | | |
| | | The bottom foot of this short section is al | | | | | | | | | |
| | | graphite. The cherty sediments are fine | | | | | | | | | |
| | | grained and light grey to brown in color. | | | | | | | | | |
| | | | | | | | | 100 | | | |
| 345. 21 | 348. 7' | QUARTZ: Green quartz vein with intense shearing a | d J-6856 | 345. 2 | 348.7 | 3, 5 | 0.005 | | | | |
| | 14.1 | heavy development of fuchite alteration. | | | | | | | | <u> </u> | |
| | | Disseminated and stringer pyrite to about | 20%. | | | | | | 1 1 1 | | |
| | | | | | | | | | | | |
| 348.71 | 409.01 | CHERTY SEDIMENTS: | | | | | | | | | |
| · | | As above in the previous units but general | у | | | | | | | | |
| | | no greywacke. From 355.8-358.1 the uni | | | | | | | | | |
| | | is brecciated with a graphite matrix and | | | | | | | | | |
| | | 75-80% angular fragments of the cherty | | | | | | | | | |
| | | sediments. | | | | | | | | | |
| 409.01 | | END OF HOLE | | | | | | | | 1 | |
| | | | | | | | | | | | |
| | | | . , | | | | | | | | |







Milner Twp. - M.236

PAT CR

MI

