



Geological Report on the Property of

GLEN AUDEN RESOURCES LIMITED

Maisonville and Grenfell Townships
Larder Lake Mining Division
District of Timiskaming

RECEIVED

NOV 15 1985

MINING LANDS SECTION

by

Nadia Cairra, B.Sc.

*Qual
2.6239.*

Robert S. Middleton Exploration Services Inc.
P.O. Box 1637 Timmins, Ontario
October 2, 1985

P4N 7W8



42A01NE0044 2.8629 MAISONVILLE

010C

TABL

SUMMARY i

INTRODUCTION 1

 Location, Access and Facilities 1

 Property 2

 Topography and Vegetation 6

HISTORY OF EXPLORATION 7

REGIONAL GEOLOGY 15

PROPERTY GEOLOGY 17

TABLE OF LITHOLOGIES 21

STRUCTURAL GEOLOGY 39

ROCK ALTERATION AND METAMORPHISM 42

ECONOMIC GEOLOGY 44

 Detailed Geology Areas 1 - 17 61

CONCLUSIONS 85

RECOMMENDATIONS 86

REFERENCES 90

CERTIFICATION

APPENDICES

LIST OF TABLES

1. Table of Lithologies
2. Description, Location, Results of Rock Samples

LIST OF FIGURES

Figure 1	Location Map		1" = 150 miles
Figure 2	Claim Index Map - Maisonville		1" = 1/2 mile
Figure 2b	Claim Index Map - Grenfell Twp.		1" = 1/2 mile
Figure 3	Previous Work		1" = 1/2 mile
Figure 4	Geology of Maisonville Twp. Map 2215		1" = 1/2 mile
Figure 5	General Geology		
Figure 6	Property Geology	sheet 1	(1:4000)
Figure 7	Property Geology	sheet 2	(1:4000)
Figure 8	Property Geology	sheet 3	(1:4000)
Figure 9	Area 1 Map	Detailed Geology	(1: 100)
Figure 10	Area 2 Map	Detailed Geology	(1: 100)
Figure 11	Area 4 Map	Detailed Geology	(1: 100)
Figure 12	Area 5 Map	Detailed Geology	(1: 100)
Figure 13	Area 6 Map	Detailed Geology	(1: 100)

SUMMARY


Glen Auden Resources hold a 122 claim property in Maisonville and Grenfell Townships, 12 miles northwest of the town of Kirkland Lake, Ontario. A number of gold showings and a former small gold producer is located within the property boundaries. The host rocks are iron-rich and iron-poor tholeiitic basalts and syenites with similar geological settings as certain gold deposits in the Kirkland Lake, Matachewan, Matheson and Timmins areas. Several areas of mineralization exist on the property within cherty sulphide facies exhalative horizons, north-south trending shear zones and within quartz-carbonate stringer zones within the iron-rich tholeiitic basalts and the syenites. The majority of these zones are good targets for large tonnage gold deposits.

INTRODUCTION

A reconnaissance geological mapping and sampling program was carried out from July 21 to September 11, 1985 on the Glen Auden Resources, Maisonville-Grenfell township property discovered several auriferous zones of mineralization. This report describes the 122 claim property held by the joint venture and is based on the geological mapping and a review of previous work records at the Resident Geologists office in Kirkland Lake. The property, which is located in the Larder Lake Mining Division of Ontario, encompasses a number of gold showings as well as a former gold producer. This is the first time, however, that the majority of the area has been assembled as one land package thereby allowing for a more comprehensive exploration program. In addition, the known gold showings are not zones of conductivity and therefore previous airborne surveys by the Ontario Geological Survey (1979) and others have not attracted much exploration effort. Weak conductors which may reflect graphitic horizons or structural (fault) zones do occur which may be related indirectly to gold mineralization. The geological mapping program has outlined several target areas for second stage exploration consisting of linecutting, induced polarization surveying, extensive trenching, and diamond drilling.

Location, Access and Facilities

The property is located in the south-west portion of



PROVINCE OF ONTARIO

REVISIONS	GEOLOGICAL ENGINEERING SERVICES Frank P. Tagliamonte P.Eng.		
	for	GLEN AUDEN RESOURCES LIMITED	
	Title	LOCATION MAP	
		FIG. 1	
	Date: JULY 1985	Scale: 1"=60mi.	N.T.S.:
	Drawn:	Approved:	File:

Maisonville Township and virtually all of the land under Lake Sese kinika in Grenfell Township. Some land is also held in Concession VII of Grenfell Township. The general area covered by the largest part of the property would be portions of Con. I, II, III and IV, Lots 5, 6, 7, 8, 9, 10 and the east half of lot 11 in Maisonville Township.

Access is via highway 11 which traverses the west side of the property. Highway 570 extends east from highway 11 through the village of Sese kinika and provides access via bush roads and trails to the south and eastern part of the property. The Ontario Northland Railroad passes through the west part of the property as does the Trans Canada Pipeline (Northern Ontario Natural Gas Pipeline).

Electrical power lines follow highways 11 and 570. Ample water is available in the many lakes and ponds that occur in the area. A trained work force resides in the Kirkland Lake area servicing the operating gold and iron ore mines.

Property

The property comprises 122 unpatented mining claims of which twenty-three are virtually all water claims in Grenfell Township together with ninety-nine mainly land claims in Maisonville Township. The water claims encompass most of Lake Sese kinika.

The 122 contiguous claim group lies within Grenfell and Maisonville Townships, Larder Lake Mining Division, Ontario.

All the claims are registered in the name of Glen Auden Resources Limited, Suite 2400 - 130 Adelaide Street West, Toronto, Ontario M5H 3C2.

Some statistical data on the claims is as listed below.

<u>Claim Number</u>	<u>Expiry Date</u>	<u>Man Days Assessment Work Filed</u>
L 778 368	July 16, 1986	20 days
L 778 369	July 16, 1986	20 days
L 778 370	July 16, 1986	20 days
L 778 371	July 16, 1986	20 days
L 778 372	July 16, 1986	20 days
L 778 373	July 16, 1986	20 days
L 778 376	August 24, 1987	100 days
L 778 377	August 24, 1986	20 days
L 778 378	August 24, 1986	20 days
L 778 379	September 4, 1986	20 days
L 798 863	August 16, 1986	20 days
L 798 864	August 16, 1986	20 days
L 798 865	August 16, 1986	20 days
L 798 866	August 16, 1986	20 days
L 798 867	August 16, 1986	20 days
L 798 868	August 16, 1986	20 days
L 798 869	August 16, 1986	20 days
L 798 870	August 16, 1986	20 days
L 798 871	August 16, 1986	20 days
L 798 872	August 16, 1986	20 days
L 798 873	September 4, 1986	20 days
L 798 874	September 4, 1986	20 days
L 798 875	September 4, 1986	20 days
L 798 876	September 4, 1986	20 days
L 798 877	September 4, 1986	20 days
L 798 878	September 4, 1986	20 days

<u>Claim Number</u>	<u>Expiry Date</u>	<u>Man Days Assessment Work Filed</u>
L 799 289	June 21, 1986	20 days
L 799 290	June 21, 1986	20 days
L 799 291	June 21, 1986	80 days
L 799 678	May 31, 1986	20 days
L 800 344	May 31, 1986	20 days
L 800 345	June 25, 1986	20 days
L 800 346	June 25, 1986	20 days
L 800 347	June 25, 1986	20 days
L 800 348	June 25, 1986	20 days
L 800 349	June 25, 1986	20 days
L 800 827	August 24, 1986	80 days
L 800 828	August 24, 1986	80 days
L 800 829	August 24, 1986	80 days
L 800 830	August 24, 1986	80 days
L 800 831	August 24, 1986	80 days
L 800 832	August 24, 1986	80 days
L 801 217	June 21, 1986	20 days
L 801 218	June 27, 1986	80 days
L 801 219	June 27, 1986	20 days
L 801 220	June 27, 1986	20 days
L 801 221	June 27, 1986	20 days
L 801 222	June 27, 1986	20 days
L 801 876	June 25, 1986	20 days
L 801 877	June 21, 1986	20 days
L 801 878	June 21, 1986	20 days
L 802 331	June 27, 1986	20 days
L 802 332	July 16, 1986	20 days
L 802 333	July 16, 1986	20 days
L 802 334	July 16, 1986	20 days
L 802 335	July 16, 1986	20 days
L 802 336	July 16, 1986	20 days
L 802 337	July 16, 1986	20 days
L 802 338	July 16, 1986	20 days
L 802 339	July 16, 1986	20 days
L 802 340	July 16, 1986	20 days
L 802 341	July 16, 1986	20 days
L 802 342	July 16, 1986	20 days
L 802 343	July 16, 1986	20 days

<u>Claim Number</u>	<u>Expiry Date</u>	<u>Man Days Assessment Work Filed</u>
L 802 346	September 4, 1986	20 days
L 802 347	September 4, 1986	20 days
L 802 348	September 4, 1986	20 days
L 802 349	September 4, 1986	20 days
L 802 353	July 9, 1986	20 days
L 802 354	July 9, 1986	20 days
L 802 355	July 9, 1986	20 days
L 802 356	July 9, 1986	20 days
L 802 357	July 9, 1986	20 days
L 802 358	July 9, 1986	20 days
L 802 359	July 9, 1986	20 days
L 802 360	July 9, 1986	20 days
L 802 365	August 13, 1986	20 days
L 802 366	August 14, 1987	100 days
L 802 367	August 14, 1987	100 days
L 802 368	August 14, 1987	100 days
L 802 369	August 14, 1987	100 days
L 802 744	August 3, 1986	20 days
L 802 745	August 3, 1986	20 days
L 802 746	August 3, 1986	20 days
L 802 747	August 3, 1986	20 days
L 802 748	August 3, 1986	20 days
L 802 749	August 3, 1986	20 days
L 802 750	September 10, 1986	20 days
L 803 374	August 24, 1986	80 days
L 803 375	August 24, 1986	80 days
L 803 376	August 24, 1986	80 days
L 803 377	August 24, 1986	80 days
L 803 378	August 24, 1986	80 days
L 803 379	August 24, 1986	80 days
L 803 380	August 24, 1986	80 days
L 803 381	August 24, 1986	80 days
L 803 382	August 24, 1986	80 days
L 803 383	August 24, 1986	80 days
L 803 384	August 24, 1986	80 days

LEE TWP

753162
753161
753160
753165
753164
753163
753167
753168

767510
767509
767511
767512

HIGHWAY 11

GRENFELL TWP

753175
753174
753173
753172
753171
753170
753177
753178
753179
783223
783222
783221
783225
783224
783226
783227
783228
783229
783230
783231
783232
783233
783234
783235
783236
783237
783238
783239
783240
783241

780781
780780
780779
780778
780777
780776
780775
780774
780773
780772
780771
780770



803374
803376
803377
803378
803379
803380
803381
803382
803383
803384
803385
803386
803387
803388
803389
803390
803391
803392
803393
803394
803395
803396
803397
803398
803399
803400
803401
803402
803403
803404
803405
803406
803407
803408
803409
803410
803411
803412
803413
803414
803415
803416
803417
803418
803419
803420
803421
803422
803423
803424
803425
803426
803427
803428
803429
803430
803431
803432
803433
803434
803435
803436
803437
803438
803439
803440
803441
803442
803443
803444
803445
803446
803447
803448
803449
803450
803451
803452
803453
803454
803455
803456
803457
803458
803459
803460
803461
803462
803463
803464
803465
803466
803467
803468
803469
803470
803471
803472
803473
803474
803475
803476
803477
803478
803479
803480
803481
803482
803483
803484
803485
803486
803487
803488
803489
803490
803491
803492
803493
803494
803495
803496
803497
803498
803499
803500
803501
803502
803503
803504
803505
803506
803507
803508
803509
803510
803511
803512
803513
803514
803515
803516
803517
803518
803519
803520
803521
803522
803523
803524
803525
803526
803527
803528
803529
803530
803531
803532
803533
803534
803535
803536
803537
803538
803539
803540
803541
803542
803543
803544
803545
803546
803547
803548
803549
803550
803551
803552
803553
803554
803555
803556
803557
803558
803559
803560
803561
803562
803563
803564
803565
803566
803567
803568
803569
803570
803571
803572
803573
803574
803575
803576
803577
803578
803579
803580
803581
803582
803583
803584
803585
803586
803587
803588
803589
803590
803591
803592
803593
803594
803595
803596
803597
803598
803599
803600
803601
803602
803603
803604
803605
803606
803607
803608
803609
803610
803611
803612
803613
803614
803615
803616
803617
803618
803619
803620
803621
803622
803623
803624
803625
803626
803627
803628
803629
803630
803631
803632
803633
803634
803635
803636
803637
803638
803639
803640
803641
803642
803643
803644
803645
803646
803647
803648
803649
803650
803651
803652
803653
803654
803655
803656
803657
803658
803659
803660
803661
803662
803663
803664
803665
803666
803667
803668
803669
803670
803671
803672
803673
803674
803675
803676
803677
803678
803679
803680
803681
803682
803683
803684
803685
803686
803687
803688
803689
803690
803691
803692
803693
803694
803695
803696
803697
803698
803699
803700
803701
803702
803703
803704
803705
803706
803707
803708
803709
803710
803711
803712
803713
803714
803715
803716
803717
803718
803719
803720
803721
803722
803723
803724
803725
803726
803727
803728
803729
803730
803731
803732
803733
803734
803735
803736
803737
803738
803739
803740
803741
803742
803743
803744
803745
803746
803747
803748
803749
803750
803751
803752
803753
803754
803755
803756
803757
803758
803759
803760
803761
803762
803763
803764
803765
803766
803767
803768
803769
803770
803771
803772
803773
803774
803775
803776
803777
803778
803779
803780
803781
803782
803783
803784
803785
803786
803787
803788
803789
803790
803791
803792
803793
803794
803795
803796
803797
803798
803799
803800
803801
803802
803803
803804
803805
803806
803807
803808
803809
803810
803811
803812
803813
803814
803815
803816
803817
803818
803819
803820
803821
803822
803823
803824
803825
803826
803827
803828
803829
803830
803831
803832
803833
803834
803835
803836
803837
803838
803839
803840
803841
803842
803843
803844
803845
803846
803847
803848
803849
803850
803851
803852
803853
803854
803855
803856
803857
803858
803859
803860
803861
803862
803863
803864
803865
803866
803867
803868
803869
803870
803871
803872
803873
803874
803875
803876
803877
803878
803879
803880
803881
803882
803883
803884
803885
803886
803887
803888
803889
803890
803891
803892
803893
803894
803895
803896
803897
803898
803899
803900
803901
803902
803903
803904
803905
803906
803907
803908
803909
803910
803911
803912
803913
803914
803915
803916
803917
803918
803919
803920
803921
803922
803923
803924
803925
803926
803927
803928
803929
803930
803931
803932
803933
803934
803935
803936
803937
803938
803939
803940
803941
803942
803943
803944
803945
803946
803947
803948
803949
803950
803951
803952
803953
803954
803955
803956
803957
803958
803959
803960
803961
803962
803963
803964
803965
803966
803967
803968
803969
803970
803971
803972
803973
803974
803975
803976
803977
803978
803979
803980
803981
803982
803983
803984
803985
803986
803987
803988
803989
803990
803991
803992
803993
803994
803995
803996
803997
803998
803999
804000

LAKE

HIGHWAY 11

MAISONVILLE TWP.

BOMPAS TWP



REVISIONS	GEOLOGICAL ENGINEERING SERVICES Frank P. Tagliamonte P. Eng.	
	for GLEN AUDEN RESOURCES LIMITED	
	Title GRENFELL TOWNSHIP PROPERTY	
	CLAIM INDEX MAP	
	Date: July 1985	Scale: 1"=1/2 mi
	Drawn:	Approved:
		File:

Fig. 25

<u>Claim Number</u>	<u>Expiry Date</u>	<u>Man Days Assessment Work Filed</u>
L 803 385	August 24, 1986	80 days
L 803 386	August 24, 1986	80 days
L 803 387	August 24, 1986	80 days
L 803 388	August 24, 1986	80 days
L 803 389	August 24, 1986	80 days
L 803 390	August 24, 1986	80 days
L 803 391	August 24, 1986	80 days
L 803 392	August 24, 1986	80 days
L 803 393	August 24, 1986	80 days
L 803 394	August 24, 1986	80 days
L 803 395	August 24, 1986	80 days
L 803 396	August 24, 1986	80 days
L 803 397	August 24, 1986	80 days
L 803 398	August 24, 1986	80 days
L 803 399	August 24, 1986	80 days
L 803 400	August 24, 1986	80 days
L 803 401	August 24, 1986	80 days
L 803 402	August 24, 1986	80 days
L 803 403	August 24, 1986	80 days
L 803 557	September 4, 1986	20 days
L 803 558	September 4, 1986	20 days
L 803 559	September 4, 1986	20 days
L 803 560	September 4, 1986	20 days

Total of 122 unpatented mining claims.

Topography and Vegetation

Topographical relief in most parts of the property is gentle to moderate. Throughout a good portion of the property, rapid changes in elevation of 50 to 100 feet occur. The iron-rich and iron-poor tholeiites and the diabase dikes form north-south trending ridges.

The highest area of relief is in the northwestern portion of the map-area where a ridge of typically resistant Cobalt Series

sedimentary rocks rise 300 feet above the surrounding tholeiitic volcanic rocks.

Vegetation on the property consists of spruce, balsam and pine with minor stands of poplar and birch. Most of the southern part of the property is covered by Sesekinika Lake. Many summer cottages are on the islands and mainland shores of the lake.

HISTORY OF EXPLORATION

The area was first prospected in the early days of the Kirkland Lake gold rush since some aspects of the geology were similar to that of Kirkland Lake area (syenite porphyry intrusives with associated gold-molybdenite mineralization). Gold was found in 1914 in Con. II, Lot 9 of Maisonville Township and considerable activity took place in the 1930's. Most of this work however, is unrecorded. Numerous pits and trenches occur on the Glen Auden Resources Limited property which contain abundant quartz and/or pyrite mineralization but no record of this work can be located in the assessment files.

The Golden Summit mine which is located approximately 200 metres south of the southern border of the Glen Auden Resources property in Maisonville Twp. in Lot 6 Concession I is reported to have produced gold. An unspecified amount of gold production is unofficially reported by previous workers in the area from the Bennett Mine in Lot 9, Con. II which is located on the Glen Auden

Resources Limited property.

Descriptions of early exploration work and the properties on which work was done is given in the section on Economic Geology. Recent exploration work is described here.

The most recent work in the area was a magnetometer survey carried out on the "Bennett Claim" (Lot 9, Con. II, N 1/2, NW 1/4) by C. Forbes in 1984. This survey located a number of high gradient anomalies associated with magnetic phases of the tholeiitic volcanics. These rocks are intruded by syenite porphyry and host north-south trending shallow dipping and vertical quartz zones. This area corresponds to AREA 12 on the property (see Economic Geology).

Prospecting by C. Forbes on various parts of the property has been successful in discovering a new gold bearing structure. Significant assays of up to .726 oz/ton gold were obtained from grab samples of pyrite bearing quartz vein material from Con. II, Lot 7, S 1/2 (claim 800 346). This unrecorded showing contains a 2' wide glassy brecciated quartz and pyrite vein possibly trending 110 - 130° with numerous parallel and branching veins and veinlets (see Photo 5). Old pits 100 feet north of this old trench were observed to contain pyrite in both the quartz veins and in the iron-rich tholeiitic wallrocks. Considerable chloritization, silicification and carbonate alteration occurs in the vicinity of the quartz veins. The numerous poor exposures

observed imply that a 50 foot wide quartz stockwork may be present. This area corresponds to AREA 5 (see Economic geology).

A north-south trending chert horizon in Lot 6, Con.III, S 1/2 has also been prospected by C. Forbes (1984). Assays as high as .143 oz Au/ton were obtained in grab samples of banded chert with pyrite. This area corresponds to AREA 1 (see Economic geology). In 1983, trenching was performed by Harold Barry in Lot 9, Con.I, N 1/2 and filed as mechanical work, however, sampling was not reported.

In 1979 an airborne electromagnetic and magnetic survey was released by the Ontario Geological Survey (1979) which covered both Maisonville and Grenfell Twps. This data can be used to trace major structures on the property including major shear zones and graphitic chert horizons.

In 1975 Ecstall Mining (Texas Gulf) did ground electromagnetic and magnetic surveys on small grids in various parts of Maisonville Township. Part of this work actually took place on the margin of the property in Lot 6-7, Con.III and V., Slankis, J.L. (1974). Earlier work by Imperial Oil in the same area tested two north trending EM conductors which revealed the presence of graphite - pyrite zones containing sphalerite mineralization. Exploration results have been compiled by Ploeger, F. et al (1979). Descriptions of early exploration work and the properties on which the work was done is described here.

An index of assessment work is shown in Fig. 3.

Bennett Mining Company Limited (33) (AREA 12 see Economic Geology)

In Maisonville Township, concession II, Lots 9 and 10, on former claim L3688, a two-compartment shaft was sunk to a depth of 530 feet and levels were cut at 125 foot intervals. Crosscutting and drifting were reported early in 1927, but in June the plant was dismantled (Kindle 1936, p.11).

About 800 feet (two holes, 133.5 and 174 feet) and 1,200 feet (one hole, 130 feet) southwest of the main shaft, diamond drilling was done in 1958 by Mr. S.A. Pain. The holes intersected diorite and volcanic rocks cut by syenite dikes and quartz veins (ODMNA files, Kirkland Lake).

C.Cole (34) (300 metres northeast of TL4800N/3500E)

In Maisonville Township, near the boundary between concession III, Lot 7, north half, northeast quarter and concession IV, Lot 7, south half, southeast quarter, four pits were sunk by Crown Reserve Mining Company Limited in 1914. The pits are in fractured silicified mafic volcanic rocks cut by a feldspar porphyry dike. The volcanic rocks contain disseminated pyrite and blebs and irregular stringers of pyrite and pyrrhotite with minor amounts of chalcopyrite. Two samples analyzed by The International Nickel Company of Canada Limited contained 0.03 and 0.04 percent nickel, according to notes by W.S. Savage (in ODMNA files, Kirkland Lake); also see Shklanka, R. (1969).

Elzina Mines Limited (35) (AREA 10 see Economic Geology)

In Maisonville Township, concession IV, Lot 10, south half, narrow quartz-carbonate veins cut hornblende syenite and gabbro. The No. 1 vein, striking N60E and dipping 50 to 70N, ranges from 3 to 14 inches wide along its stripped length of 200 feet. Grab samples taken at intervals along the vein returned assays of: .02, .03, .02, .01, and .25 oz gold per ton (D.K. Burke 1938). Three holes have been drilled on the property, but no information on them is available.

The Golden Summit Mining Company Limited (36)
(approximately 200 metres south of TL1576N 4480E)

In Maisonville Township, concession I, Lot 6, north half (formerly the Jensen farm) a gold-bearing quartz vein was discovered south of Kapakita Creek, in 1913. The vein was 6 inches wide and was traced for 150 feet. In 1919, the Golden Summit Mining Company Limited was incorporated to take over the property, and in 1921, a pit on the discovery vein was deepened to a depth of 27 feet. In 1924 the company was reorganized to Golden Summit Mines Limited. During 1929 and ensuing years a shaft collared 450 feet west of the old pit was deepened to 405 feet and lateral work was done on the 125-, 250-, and 375-foot levels. A small mill installed in 1935 produced during 1936 and 1937. By 1937, when mining ended, some stoping had been done on the 125-foot level. From 737 tons of ore, total production worth \$3,738 was recorded with the Ontario Department of Mines for 1936, 1937, and 1945, when mill clean-up was carried out (ODM 1953, p.16).

Most of the gold is in pyrite-containing quartz-carbonate veins cutting basalt, diorite, syenite, and lamprophyre. Assays of .16 oz gold and less were obtained in the shaft station at the 125-foot level, and .08 oz gold or less along the crosscut (Odell 1930). The shaft was sunk in the vicinity of two narrow east-trending quartz veins about 40 feet apart, the south vein being the discovery vein. Immediately east of the shaft is a strong shear zone trending north, parallel to the strike of the flow. The east wall of the shear is a lamprophyre dike about 50 feet wide. The most important vein encountered underground was exposed in the northwest crosscut on the 125-foot level. It was a high-grade quartz vein striking east and ranging in width from 2 inches to 10 inches. It had been intersected and displaced by a flat fault dipping west. The wall rocks adjacent to the vein were highly fractured and veined with red feldspathic material; they contained gold.

Hastings-Moffat (37) (AREA 7, AREA 9 see Economic Geology)

In Maisonville Township, concession II, Lot 8, south half, southwest quarter, five holes were drilled in 1934 by Erie Canadian Mines Limited, the exploration subsidiary of Sylvanite Gold Mines Limited. This drilling totalled 464 feet consisted of quartz veins striking and dipping in several directions, in an area having a diameter of about 400 feet. The host rock is syenite and altered mafic rocks. A few short intersections (average 2 feet) contained gold yielding about .02 oz gold per ton. The best assay of several samples from pits in Lot 8 and 9 yielded .17 oz gold per ton (Erie Canadian 1934 - ODMNA files, Kirkland Lake).

Labine-Smith (41) (Surrounded by the Glen Auden Resources property on Patented Ground-south and east of Area 12.)

In Maisonville Township, concession II, Lot 9, north half, southwest quarter and northwest quarter, and Lot 10, north half, northeast quarter, five pits and trenches were carefully sampled (ODMNA files, Kirkland Lake, report signed G.W. Dixon, August 10, 1914). Most assays for gold were less than .04 oz gold per ton and for silver were less than 1 ounce per ton.

A shaft with an average dip of 35 degrees, had been sunk to a depth of 80 feet on a narrow quartz vein carrying free gold and tellurides.

Sidney, A. Pain (45) (200 metres south of TL3192N/1000E 200 metres from property boundary)

In Maisonville Township, concession II, Lot 10, north half, northwest quarter, native gold was discovered in quartz veins and oxidized wall-rock on the former "Malouf" claim (Burrows and Hopkins 1914, p.35). The decomposed material from the quartz veins, which are narrow and dip gently west, was treated in an arrastra at the base of the hill near the railway track.

Warrer. Peterson (46) (In the northwest corner of CL 803558 AREA 16)

In Maisonville Township, concession IV, Lot 9, south half, northwest quarter, in the northwest quarter of the claim (formerly numbered L46158), is the main (copper) showing. The showing is on the east side of a north-trending ridge of mafic volcanic rocks and consists of syenite and mafic volcanic rocks cut by quartz stringers containing cubic pyrite and fine-grained chalcopyrite. A dike of "pebble" lamprophyre cut by a feldspar porphyry stringer occurs in the northeast corner of one pit. In 1952, three holes (totalling 1,000.5 feet) drilled near the showings intersected a number of weakly mineralized zones in sheared and massive mafic volcanic rocks cut by syenite dikes (ODMNA files, Kirkland Lake).

A. Pudden (47) LOT 8 - AREA 11, AREA 2 LOT 10 - in the vicinity of claim 778377

In Maisonville Township, concession II, Lot 8, north half, northwest quarter and southwest quarter, and in Lot 10, south half, southeast quarter, nine holes, totalling 416 feet, were drilled in 1955. The holes were spaced along a north-striking line about 1/2 mile long. Judging from the azimuth of eight of the nine holes, the veins dip east. The holes intersected quartz veins cutting mafic intrusive rocks containing sparsely disseminated pyrite and chalcopyrite (ODMNA files, Kirkland Lake).

J.G. Russell (48) (In a patented claim 100 metres south of CL 798865 southwest of Olson Lake)

In Maisonville Township, concession III, Lot 9, south half, southeast quarter, near the No. 1 post of claim L15833, a rusty silificied shear zone has been exposed by pits on the west-facing slope of a hill adjacent to a swamp. The strike of the shear zone, which is in mafic volcanic rocks, is N10W and the dip is about 65W. Irregular quartz stringers and feldspathic material cut the sheared rocks, which contain finely disseminated pyrite and minor molybdenite.

About 500 feet southwest, on the western side of a low outcrop, a quartz vein 4 to 6 inches wide strikes

About 500 feet southwest, on the western side of a low outcrop, a quartz vein 4 to 6 inches wide strikes north and dips 20 to 40 west through pillowed volcanic rocks. A narrow streak of darker pyrite-bearing quartz on the footwall is said to have contained gold and tellurides, according to notes by W.S. Savage (in ODMNA files, Kirkland Lake). In the south trench the vein forms a zone having a maximum width of 4 feet.


About 150 feet east of the trench, on the eastern side of a low north-trending ridge of volcanic rocks near the centre of the claim, brecciated andesite is cemented with a fine network of quartz-carbonate-feldspar stringers reported to have contained molybdenite.

On knoll about 250 feet south of the above showing, quartz-carbonate-feldspar stringers, an average of 1 inch wide, cut brecciated andesite containing pyrite and epidote. The eastern side of the knoll is cut by a syenite dike, and a sample from a trench cut through the knoll from east to west is said to have yielded .14 oz gold per ton across 22 feet, according to Savage.

Sesekinika Townsite (49) (AREA 7 see Economic Geology)

According to notes by Sylvanite Gold Mines Limited (in ODMNA files, Kirkland Lake), during 1914 some free gold was found by James L. Hughes on claim L4034 in Maisonville Township. The gold was in a narrow quartz vein dipping 15NW. About 1916, Hughes sold the property to Walter Young (of Toronto) and New York interests.

The part of the "Hastings-Moffat" claim L4035 (which is described separately but is redescribed here by Lovell since it is also part of this property) that is in Maisonville Township, concession II, Lot 8, south half, southwest quarter, was drilled during 1934 by Erie Canadian Mines Limited, the exploration subsidiary of Sylvanite. Five holes, totalling 464 feet, were drilled to intersect a quartz vein 2 feet wide that strikes east and dips 60N. A few short intersections (average 2 feet) contained gold yielding about .02 oz per ton. The best assay from several samples from pits in Lots 8 and 9 yielded .17 oz gold per ton (Erie



MAISONVILLE TWP.

GRENFELL TWP.

NOTE: - * covered with

- No previous work in Grenfell Township Property

REVISIONS	GEOLOGICAL ENGINEERING SERVICES Frank P. Tagliamonte P. Eng.	
	for	GLEN AUDEN RESOURCES LIMITED
	Title	MAISONVILLE TOWNSHIP PROPERTY
		PREVIOUS WORK FIG. 3
	Date July 1985	Scale: 1" = 1/2 mile N.T.S.
	Drawn:	Approved: File:

Canadian Mines Limited 1934).

REGIONAL GEOLOGY

The Maisonville township property is located in the south part of the Kirkland Lake area and is underlain by Archean volcanic, sedimentary, and intrusive rocks which are part of the Abitibi Greenstone Belt. In the south part of the Kirkland Lake area, the Archean rocks are unconformably overlain by Proterozoic sedimentary rocks of the Cobalt series. Diabase dikes cut all of the rock types in the area the Cobalt sediments, which occur in the northwest corner of the property of interest.

In the Kirkland Lake area, the volcanic rocks of the Abitibi Greenstone belt are preserved in a synclinorium located between the Lake Abitibi Batholith and the Round Lake Batholith. The north and south limbs of the synclinorium are cut by two large east-striking fault zones: the Destor-Porcupine Fault Zone, and the Kirkland Lake-Larder Lake Fault Zone, respectively.

Rocks in the area have been affected by subgreenschist regional metamorphism.

The volcanic rocks in the Kirkland Lake Region were formed during cycles of volcanism associated with sedimentation and plutonism. Each cycle consisted of komatiitic volcanism followed by the tholeiitic, calc-alkalic, and finally, alkalic volcanism (Jensen O.G.S., 1979).


The Lower Supergroup on the south limb of the synclinorium,

southeast of Kirkland Lake, contains a cycle of komatiitic, tholeiitic, and calc-alkalic volcanics represented by the Wabewawa, Catherine, and Skead Groups, respectively (Jensen 1979).

The volcanic cycle represented by the Upper Supergroup consists of komatiitic lavas of the Larder Lake Group succeeded by tholeiitic rocks of the Kinojevis Group and calc-alkalic rocks of the Blake River Group. The Maisonville-Grenfell Township property of Glen Auden Resources Limited is underlain by the tholeiitic rocks of the Kinojevis Group and partly by interflow tuffs of calc-alkalic composition of the Blake River Group. This rock series is similar to the Tisdale Group in the Timmins area which contains numerous well known gold deposits. Alkalic volcanic rocks of the Timiskaming Group unconformably overlies the Blake River and kinojevis Groups.

On the north limb of the synclinorium, the oldest rocks are calc-alkalic volcanic rocks of the Hunter Mine Group. Unconformably overlying the Hunter Mine Group is a komatiitic succession called the Stoughton-Roquemaure Group (Jensen, 1976a, 1978b, O.G.S.)

The sinking of volcanic rocks during their accumulation would explain the origin of the synclinorium. Various intrusive differentiates including monzonite, porphyritic syenite, mafic intrusive syenite, and augite syenite intruded the above



MAISONVILLE TWP.
GRENFELL TWP.

NOTE : See next page for legend

REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for	GLEN AUDEN RESOURCES LIMITED	
	Title	General Geology Map	
		Fig. 5	
	Date Oct. 1985	Scale: 1" = 1/2 mile	N.T.S.
	Drawn: C.G.	Approved:	File: M-78

LEGEND

CENOZOIC^a

RECENT

Swamp and stream deposits.

PLEISTOCENE

Sand, gravel, clay.

UNCONFORMITY

PRECAMBRIAN^b

PROTEROZOIC

HURONIAN

COBALT GROUP

GOWGANDA FORMATION

7

- 7a Conglomerate.*
- 7b Arkose.*
- 7c Quartzite and greywacke.*
- 7d Argillite.*

UNCONFORMITY

ARCHEAN

MAFIC INTRUSIVE ROCKS (MATACHEWAN)

6

- 6 Diabase.*

INTRUSIVE CONTACT

FELSIC INTRUSIVE ROCKS (ALGOMAN)

5

- 5a Granite, hornblende-biotite granite, granodiorite.*
- 5b Granite gneiss.*
- 5c Syenite, syenite porphyry, quartz syenite.*
- 5d Mafic syenite, lamprophyre.*

INTRUSIVE CONTACT

MAFIC AND ULTRAMAFIC INTRUSIVE ROCKS (HAILEYBURIAN, EARLY ALGOMAN, AND SOME KEEWATIN)

4

- 4a Serpentinite, peridotite.†*
- 4b Gabbro, diorite, may include some 1a.*

INTRUSIVE CONTACT

METASEDIMENTS, TIGHTLY FOLDED (TIMISKAMING OR KEEWATIN)

3

- 3 Slate, chert, tuff.*

UNCONFORMITY

FELSIC METAVOLCANICS (KEEWATIN)

2

- 2a Dacite, rhyodacite, minor amounts of rhyolite.*
- 2b Silicic and intermediate tuff and agglomerate.*
- 2c Dacite porphyry, amygdaloidal dacite, spherulitic dacite.*

MAFIC TO INTERMEDIATE METAVOLCANICS (KEEWATIN)

1

- 1a Basalt and andesite (pillowed and massive) may include some 4b.*
- 1b Mafic agglomerate and tuff.*
- 1c Altered (bleached) mafic metavolcanics.*
- 1d Andesite porphyry, amygdaloidal andesite, variolitic andesite.*

Ag	Silver.
asb	Asbestos
Au	Gold.
carb	Carbonate.
Cu	Copper.
ep	Epidote.
gf	Graphite.
gt	Garnet.
hem	Hematite.
mag	Magnetite.
Mo	Molybdenum.
Ni	Nickel.
Pb	Lead.
q	Quartz.
qc	Quartz-carbonate.
S	Sulphide mineralization
shee	Scheelite.†
Te	Tellurium.
Zn	Zinc.

mentioned rocks as plugs, dikes and/or sills. Gold mineralization frequently occurs around or within the syenite bodies as well as along north-south trending fault structures. Major north-south faults have been interpreted (Middleton 1976) based on the regional gravity data. These features are also discernible from the aeromagnetic data, O.G.S.(1979).

PROPERTY GEOLOGY

The oldest rocks on the property are the tholeiitic basalts of the Kirrojevus Group and are divided into iron-rich and iron-poor, tholeiitic basalts. The coarser-grained variety (2-5mm) of the basalts were previously mapped as coarse-grained gabbro intrusives, however, more recent mapping in the area by L. Jensen O.G.S., has discovered that these "gabbro intrusions" are in fact coarse-grained iron-rich tholeiitic basalt flows. Similar coarse-grained flows occur on the Maisonville-Grenfell township property, and it is probable that no gabbro intrusives exist on the property to the authors knowledge.

These tholeiitic basalts have been intruded by a series of felsic syenitic dikes and mafic diabasic dikes. Rocks of the felsic variety vary from earlier trondhjemitic phases to porphyritic syenite, syenodiorite to a mafic syenite. Alteration around these dikes includes silicification, epidotization and the introduction of hydrogarnet and carbonate along fractures.

The predominant rock units that occur on the Maisonville-Grenfell Township property are massive (2-5 m) coarse-grained, fine-grained, and pillowed iron-rich and iron-poor, tholeiitic basalts. These mafic volcanics are typically dark green to black in the more iron-rich varieties and a lighter green colour in the iron-poor basalts. The eastern portion of the property is underlain by predominantly iron-poor, tholeiitic basalts with a minor narrow iron-rich tholeiitic basalt sequence containing sheared chert, cherty tuffs and crystal tuff interflow units. Throughout the remainder of the property the coarse-grained iron-rich tholeiites form units 50 to 500 m thick that alternate with finer grained units 2-5 m thick. A few of the fine-grained map units represent a gradational and compositional change in the rocks from iron-rich tholeiite to an iron-poor tholeiite generally confirmed by the magnetism of the rock. The iron-rich tholeiites are strongly magnetic while the iron-poor tholeiites are weakly magnetic. Some of the coarse-grained iron-poor, magnesium tholeiites appear to grade laterally into fine-grained pillowed lavas of the same composition. Minor amounts of pillowed iron-rich tholeiitic basalt also occur.

These iron-rich tholeiitic basalt flows are thought to be the extrusive equivalents of the gabbros occurring in the townships north of the property. The claims covering most of Sesekinika

Lake are underlain by predominantly iron-poor tholeiitic basalts that vary from massive coarse-grained (2-5 mm) flows, to fine-grained massive flows, to pillowed flows with minor variolitic and hyaloclastite phases. These mafic volcanics are medium green and weakly magnetic. Minor syenitic dikes and diabasic dikes with varying amounts of magnetite cut the mafic volcanics. These may correspond to several magnetic anomalies that occur on the regional aeromagnetic map.

Several interflow units of chert, cherty tuff, cherty oxide iron formations, crystal tuff, and tuff breccia of calc-alkalic dacite composition occur on the property overlying the iron-rich tholeiitic basalts. These may represent the waning phases of the volcanism. They may mark periods of quiescence the change from iron-rich tholeiitic basalt to iron-poor, tholeiitic basalts.

The shear zones consist of highly fractured, epidotized, material containing 5-15% pyrite as fine stratiform seams, stringer pyrite and as disseminations.

The lava flows are, in places pillowed, variolitic, amygdaloidal, and hyaloclastic. Little evidence of stretching or flattening is visible in the area. The only trace of movement is the presence of chlorite-carbonate rich slickensides at the flow contacts, and in some pillow selvages of the lava flows.

The presence of interflow tuff units of calc-alkalic composition may indicate an overlap of the later calc-alkalic

Blake River Group into the Kinojevis Group. The crystal tuff unit seems, from field relationships, to occur along the contact between iron-rich and the iron-poor, tholeiite flow contacts, following bedding contacts as determined from pillow tops and from other sedimentary horizon contacts.

Major diabase dikes were found, predominantly in the eastern portion of the property around Kapakita Lake. At this location the dikes cut predominantly iron-poor, tholeiitic basalts. However, throughout the remainder of the property, the similarity between the diabase dikes and the iron-rich coarse-grained (2-5 mm) tholeiitic basalts posed a constant problem throughout the mapping.

The major north trending diabase dikes are thought to follow fault zones and major joint directions.

TABLE OF LITHOLOGIES

PRECAMBRIAN

Huronian (Cobalt Group)

- 7a Conglomerate
- 7b Arkose
- 7c Quartzite, greywacke

unconformity

ARCHEAN

Mafic Intrusive Rocks

- 6 Diabase

Felsic Intrusive Rocks

- 5b Porphyritic syenite
- 5c Syenodiorite
- 5d Mafic Syenite

Metasediments

- | | |
|-----------------------------------|--------------------------------|
| 3a cherty sulphide facies Iron fm | 3d cherty oxide iron formation |
| 3b slate, argillite | 3e graphite |
| 3c laminated, siliceous | 3f cherty tuff |

UNCONFORMITY

MAFIC METAVOLCANICS (Kinojevis Group)

2. Iron-rich tholeiitic basalt

- a. massive, finegrained
- b. pillowed
- c. pillow breccia
- d. massive (2-5 mm) coarsegrained
- f. amygdaloidal
- g. interflow intermediate crystal tuff --
overlap of Blake River Volcanism

1. Magnesium, Iron-poor tholeiitic basalts

- a. massive, finegrained
- b. pillowed
- c. pillow breccia
- d. massive (2-5 mm) coarsegrained
- e. hyaloclastite, fragmental
- f. amygdaloidal
- h. variolitic
- i. chlorite schist

Huronian (Cobalt Group) Metasediments

Fairly flat-lying sedimentary rocks form high ridges in the northwestern parts of the Maisonville property, north of the Whiteclay River. Cliffs about 200 feet high contain intermixed sedimentary rocks including conglomerate, arkose and a quartzitic graywacke. These rocks unconformably overly all other types of rock in the map area.

Unit 7a Conglomerate

The conglomerate forms the majority of the sedimentary rocks on the property. Subangular to subrounded pebbles (and boulders) from a variety of rock types comprise 5 to 40% of the conglomerate, and range from 1 cm. to two meters in diameter. Granite pebbles are predominant, but syenite, granodiorite gneiss, rhyolite, andesite, jasper iron formation, basalt, diabase, and a few pebbles of quartz vein material also were present. The matrix of most of the conglomerate that contains a high proportion of pebbles, is fine to medium grained greenish-grey or pink quartzite or pink or dark-red arkose. The matrix of most of the conglomerate that contains a low proportion of pebbles is a massive fine-grained argillaceous quartzite.

The immaturity of the sediment and subangularity of many pebbles indicates that the source rock was not subjected to chemical weathering, and the pebbles have not been transported far from their source. Many people believe that the conglomerate

is a tillite.

Unit 7b Arkose

This rock is seen in areas where a high proportion of pebbles occur in the conglomerate. The matrix of the conglomerate is a pink or dark-red arkose and often forms thin pink beds of arkose within the conglomeritic unit.

Unit 7c Quartzite, greywacke

This unit forms the matrix of the conglomerate where a low proportion of pebbles occur. This rock has a greenish gray matrix, is medium grained and forms thin beds. Some outcrops in the quartzite-arkose sequences contain some evidence of water action such as graded bedding.

Mafic Intrusive Rocks

Unit 6 Diabase

Diabase dikes are seen cutting all of the rocks in the map area except for the Huronian metasediments. Weathered surfaces are dark grey or rusty brown and fresh surfaces are grey to greyish black. The coarser grained variety is sometimes mistaken for Unit 2d, massive (2-5 mm.) coarse-grained iron-rich tholeiitic basalt, and the darker fine-grained chill margins of the dikes are often mistaken for Unit 2a, the massive fine grained iron-rich tholeiitic basalt. The diabase dikes alter the mafic volcanics only slightly (chlorite and feldspar in the wall-rocks with lesser amounts of magnetite, pyrite, sericite,

and calcite). Most of the diabase dikes fill faults and joints striking within 15 degrees of north.

Felsic Intrusive Rocks

In outcrop, these rocks show up as red to light pink zones cutting the dark green to black metavolcanic country rocks. The dikes show few or no inclusions whereas the larger stocks tend to contain numerous dark green, altered inclusions 1 cm. to 3 cm. in size. In some places, the dikes cut one another indicating that the syenitic rocks were produced during two or more pulses of igneous activity. Crystallization occurred with differentiation which generated other forms of syenite. The syenites are fractured into three strong joint directions and several less well developed irregular fracture directions. These joints are often injected by mineralized silica-rich fluids, that locally are auriferous.

Unit 5b Porphyritic Syenite

This rock occurs as red dikes crosscutting the green metavolcanics on the property. Fresh surfaces are pink to red in colour comprised of fine feldspar and sericite (70-75%) speckled with 10% fine hornblende needles (0.1 to 1 mm. long) or patches of chlorite 1 to 2 mm. in diameter, and 1 to 3% pyrite and magnetite or up to 5% adjacent to shear or fault zones, generally along contacts. Phenocrysts (5-10%) of partly altered euhedral plagioclase, 1 mm. to 4 mm. (average 2 mm.) in diameter also

occur. These dikes have metamorphic haloes including a potassium metasomatism in the surrounding volcanic rocks. Varying amounts of pyrite and chalcopyrite mineralization also occurs with associated spiderweb quartz stringer systems adjacent to contacts with the metavolcanic rocks.

Unit 5c Syenodiorite

This equigranular rock occurs as dikes crosscutting the mafic metavolcanic rocks. Unlike the other syenite dikes, these contain quartz that occurs interstitially to the feldspars. Weathering surfaces are pink to red in colour and are comprised of 85% feldspar and sericite, 5% hornblende needles or chlorite patches, 1 to 3% pyrite, and 5% clear quartz grains. The metamorphic aureole produced by these dikes is minimal with only minor amounts of a red hematitic alteration.

Unit 5d Mafic Syenite

This unit weathers a pink to white colour, while fresh surfaces are light grey. The matrix is aphanitic and contains 30-40% black acicular pyroxene (augite?) phenocrysts from 1-2 mm. in length. The rock is moderately magnetic to nonmagnetic and occasionally contains 1-2% disseminated pyrite. The contacts have metamorphic aureoles including silicification, epidotization and hematization of the country rock. It is difficult to tell true composition from solely hand specimen examination due to the aphanitic matrix but it is thought that

the rock is comprised of mostly feldspar and sericite.

Metasediments

The bulk of the sedimentary rocks in the area consists of interbedded cherty sulphide formations. Other rocks include argillite, graphite, a laminated siliceous sediment and a cherty oxide iron formation. In some localities massive chert beds occur over a considerable distance (400 meters); in others a cherty oxide iron formation with narrow graphite beds is predominant; and in many places a finely laminated siliceous metasediment occurs that often contains considerable sulfides; finally in a few localities massive chert and a cherty tuff occur.

Cherty Sulphide Facies Iron Formation

This unit occurs as 2 m. to 10 m. wide, massive, granular chert beds lying stratigraphically above the iron-rich tholeiitic basalts. In one location these chert horizons can be traced for distances of over 400 metres. Cherty sulphide facies iron formations are predominantly composed of white to light grey chert weathering a buff white colour. The chert occurs in outcrop as follows:

- (a) AREA 1: 50 m. west of No. 2 of 798875
- (b) AREA 3: 100 m. north of No. 2 of 802747 1-2% py
- (c) AREA 4: L1600E/3240N and 20 m. east
- (d) AREA 8: 160 m. W. of No. 1 of 802346 at TL 3192N at 3140E

- (e) AREA 11: TL 3192N at 2480E
- (f) CL80233: TL4800N 2830 E, 180 m. East of No. 1 of 802337,
200 m. east of CL802336 No. 2 of 802337
- (g) CL802332: TL4800N/2720 m. E. and 50 m. south

Mineralization within the chert characteristically occurs as both statiform-type, 1-3 mm. thick layers. and as fine disseminations. Within shear zones, the cherts are highly fractured and contain stringer pyrite and associated epidote and carbonate. In general pyrite constitutes less than 8% of the chert. The cherts reflect the effect of folding within the tholeiites from varying attitudes and many of the chert layers seem disrupted and broken, the dilatent zones being filled by magnetite or pyrite.

Most of the sulphide facies iron formations do not appear to form continuous layers, it appears that they form a series of disconnected, isolated lenses, perhaps in part forming an en-echelon type arrangement. All of the sulphide facies iron formations occur in the top or near to the top of the iron-tholeiites. The sulphide facies iron formation occurs in greater abundance than the oxide facies on the Maisonville Township Glen Auden Resources Ltd. property.

Unit 3b Slate, Argillite

Minor grey slaty sedimentary rocks form units up to .5 m. thick and occur in AREA 1 in trench - 1 (see detailed geology) in

the eastern portion of the property interbedded with chert and cherty tuff units. The slate is highly friable, grey in colour, and occurs along the hanging wall of the cherty tuff, Unit 3f. The unit is no more than 12 cm. in width and contains only trace amounts of pyrite.

Unit 3c Laminated, Siliceous

This unit occurs in: Area 1, Area 2 and in Area 4 (see detailed geology section) as .5 m., and 5 metre wide beds, respectively. The rock is very finely laminated on a scale of 1-2 mm. and is comprised of finegrained white (quartz), light green (epidote), red, maroon (hematitic feldspar?) and dark green (chlorite) laminae. The matrix is aphanitic, strongly siliceous, and where sheared, contains up to 4% stringer pyrite. It is still unsure as to the origin of this unit as it usually occurs wholly within the iron-rich tholeiitic basalts in the middle of the iron-rich sequence, unlike the other metasediments and tuffs that occur stratigraphically along the contact with the iron-rich and the iron-poor tholeiitic basalts.

Unit 3d Cherty Oxide Iron formation

This unit occurs in Area 8 and in Area 16 as narrow 10-100 cm. wide beds within the iron-rich tholeiitic basalts near to the contact with iron-poor, magnesium tholeiitic basalts. The oxide facies iron formation typically consists of alternating layers of magnetite and chert. The magnetite layers are 1 mm. to 20 mm.

thick, and average approximately 5 mm. in thickness, and consist of fine-grained granular magnetite and chert. Chert layers are dark grey to black on fresh and white on weathering surfaces, and vary from few millimetres to 30 cm. or more in thickness yet commonly average approximately 1 to 3 cm., and are almost invariably present in greater amounts than their magnetite-rich layers. The chert looks sugary, recrystallized, granular and contains minor grains of pyrite, pyrrhotite and magnetite.

Unit 3e Graphite

This unit occurs as discontinuous layers from 1 cm. to 3 cm. thick within or interbedded with the sulphide facies iron formations in Area 1, Area 8 and Area 16. The layers are dark grey to black, soft and occur as thin layers within the cherty oxide iron formations.

Unit 3f Cherty Ash Tuff

This unit occurs in Area 1, foot wall to the sulphide facies iron formation. The cherty tuff typically consists of an aphanitic, siliceous, cherty, black matrix with 1-2% clear quartz eyes, 10-15% black biotite phenocrysts and 5% white plagioclase crystals. The cherty tuff contains up to 5% stringer pyrite with associated epidote and carbonate along fractures within shear zones.

MAFIC METAVOLCANICS (Kinojevis Group)

The oldest and most extensive rock units on the property are the 50 m. to 500 m. wide iron-rich and iron-poor magnesium tholeiitic basalts of the kinojevis Group, primarily the massive (2-5 mm.) coarse grained and pillowed flows. On the aeromagnetic map the kinojevis group has a distinctly striped pattern caused by groups of iron-rich tholeiitic basalts alternating with low-iron, magnesium tholeiitic basalts of lower magnetic content.

A typical cycle of the kinojevis Group occurring on the Maisonville-Grenfell Township property consists of a greenish grey magnesium iron-poor basalt at the base, a black to dark green iron-rich, low-magnesium basalt as the main unit, and a thin cherty unit at the top. An overall iron enrichment trend occurs cyclically within the main trend in the volcanic sequence. According to L. Jensen, O.G.S., 1983 the lower 1000 m. of the kinojevis group also contains interflow-units of tuff breccia, crystal tuff and cherty tuff of calc-alkalic dacite and rhyolite composition, as well as chert, argillite, graphite and cherty oxide iron formation. It would appear that the eastern half of the Glen Auden Resources property occupies the lower portion of the kinojevis group. The reason why the beds of interflow tuff and interlayered beds of carbonaceous and argillaceous sediment of the lower 1000 metres of the kinojevis group have the composition of calc-alkalic rocks is not known. Their presence

suggests a pause in the mafic volcanism and they may in fact represent an overlap of the Blake River volcanic event comprised predominantly of calc-alkalic rocks.

The upper part of the pillowed flows of the low-iron basalts is hyaloclastite material occurring as either matrix material within the pillow breccia or as thin 10 cm. - 100 cm. beds topping the pillow breccia unit. The pillow lavas in the iron-rich and iron poor tholeiites are variolitic in many places.

Iron-rich Tholeiitic Basalt

Unit 2a Massive, fine-grained

This rock is characterized by a rusty weathering surface a black to dark green coloured fresh and weathered surface and a high magnetite content, and occur as units 10 cm. to 30 cm. thick that often lie stratigraphically along the contacts of the more massive diabasic to gabbroic textured lava flows. Slightly coarser grained portions (1-2 mm.) also occur within this unit as 10m.-100m. thick beds. This fine-grained basalt grades rapidly upward into the coarse-grained diabasic and gabbroic textured basalt.

Occasional light green clusters of saussuritized plagioclase also occurs. Where traversed by quartz veining the rock is silicified and may contain disseminated pyrite concentrated along the vein selvages.

Unit 2b Pillowed

Some of the massive flows (Unit 2d) are separated from one another by pillowed lava flows from 15m to 200m thick. "The pillows are closely packed and have good pillow ellipsoidal shape. On weathered surfaces, the pillows are 0.3 to 0.9m. in diameter, and are characterized by their strongly weathered, rusty selvages. The selvages are 1 to 2cm. thick, and have pillow fragments and altered glass globules and shards in their thicker portions. "The pillow tops are smooth, convex, curved surfaces and at the pillow base, tails project downward between underlying pillows. Where the pillow lavas directly overlie a more massive lava flow they are slightly flattened parallel to the strike of the lava flow and have no tails."¹ In texture and mineralogy, the fine-grained pillowed basalt differs from the coarser grained massive basalt. Plagioclase laths 0.1 to 1mm. in length comprise 40 to 45 percent of the rock. Subhedral to anhedral platy augite grains 0.2 to 1mm. in size optically enclose the plagioclase and comprise 40 to 45 percent of the rock. Approximately 10 percent magnetite, quartz, and sulfides form the remaining portion of the rock.

Unit 2c Pillow Breccia

The pillow breccia within the iron-rich tholeiitic basalts is rare occurring in a few localities in the east central part of

¹ Jensen, L.S. and Langford, F.F., 1983, Geology and Petrogenesis of the Archean Abitibi Belt in the Kirkland Lake Area, Ontario, Ontario Geological Survey, Open File Report 5455, pp. 520.

the property. The hyaloclastite material in the iron-rich tholeiitic basalt, is similar to that found in the low-iron or iron-poor tholeiite, except for its dark green to black colour and its higher iron content. The hyaloclastite is an aphanitic, tuffaceous rock with a rough brown weathering surface. It is composed of shards and granules of chloritized tachylite in a chloritized matrix.

Unit 2d Massive (2-5 mm.) coarse grained flow

The massive iron-rich basalt has rounded, smooth, dark-green partly rusted weathering surfaces. On fresh surfaces, the basalt is a black to dark green rock mainly because it contains numerous hornblende grains 2-4mm., long and because of its high magnetite content. According to Jensen 1983, O.G.S. the rock is composed of 40 to 50 percent late magmatic hornblende, pseudomorph after subhedral clinopyroxene, and ophitic to euhedral plagioclase which forms 25 to 30 percent of the rock. The remainder of the rock is composed of 5 to 10 percent titaniferous magnetite, 5 to 10 percent chlorite, and 2 to 8 percent interstitial quartz. In most places, the plagioclase is partly altered to epidote, chlorite and albite. According to Jensen 1983, the alteration occurred during the late cooling stage of the basalt. Near to the syenitic dikes the texture of the basalt is partly destroyed by the formation of fine actinolite, epidote and quartz.

Unit 2f Amygdaloidal

This unit occurs towards the upper portion of the flow, and consists of homogeneous medium-grained basalt which contains numerous irregular cooling fractures. The amygdules are from 0.5mm. to 2mm. in size and are filled with quartz and occasionally calcite and epidote. The amygdaloidal basalt often forms a knobby weathering surface due to the more resistant quartz amygdules. This unit is strongly magnetic.

Unit 2g Interflow Intermediate Crystal Tuff

These rocks occur as interflow units 10 cm. to 10 m. thick with a matrix ranging from green to light grey, and crystals ranging from 1 mm. to 4 mm. in size. Many of these units form the upper portion of the iron-rich tholeiitic basalt sequences and are useful as marker horizons for the tops of the iron-rich tholeiitic cycle. On the outcrop, it weathers a buff, light grey to green colour with white, partly kaolinized broken plagioclase crystals, angular to subrounded in shape and minor rounded quartz phenocrysts (1 mm. to 4 mm.). The crystals form from 45 to 70 percent of the rock set in an amorphous matrix. The origin of this unit is thought by the author to be formed from an overlap of the later Blake River calc-alkalic volcanic event with the later Kinojevis Group tholeiitic event.

Unit 2h Variolitic

The pillow lavas in the iron-rich tholeiitic flows are variolitic in many places. The variolites range in size from 1 to 5mm., and occur as layers around the chilled rims of the pillows. On the outcrop, the variolites show up as knobs more resistant to weathering than the enclosing dark green or rusty brown weathered surface of the fine-grained basalt. The variolites are comprised of plagioclase crystals.

Unit 2i Chlorite Schist

The sheared iron-rich tholeiite is strongly foliated, dark green and is composed primarily of chlorite and carbonate. This unit occurs in only a few localities on the property and usually marks lithologic contacts. Trace amounts of disseminated pyrite occur within these zones.

1. Iron-poor tholeiitic basalt

Unit 1a massive, fine-grained

This rock is characterized by a light to medium green weathering and fresh surface, an aphanitic matrix, and a weak magnetic susceptibility and occurs as flows 1.5 m. to 100 m. thick. In general, the massive flows occur toward the base of a formation of iron-poor tholeiitic basalt and are in places separated from one another by pillow-lavas and pillow-breccias ranging from 15 to 100 m. thick. This fine-grained basalt grades rapidly upward into diabasic and gabbroic textured basalt. Where

traversed by quartz spider-web veining the rock is silicified and may contain pyrite along the vein selvages.

Unit 1b Pillowed

In the iron-poor tholeiitic basalts, some of the more massive flows (Unit 1d) are separated by pillowed lava flows that are in greater abundance than in the iron-rich tholeiites. The pillows range from 0.3 to 1 m. in diameter and many have a preferred elongation. Where tightly packed the pillows are ellipsoidal and molded against those below. Some pillowed units are formed of loosely packed, irregular, bulb-shaped pillows. The interstices between the pillows are filled with a light green hyaloclastic material.

The pillows are easily visible because of the chilled rims, which are 2 to 3 cm. thick, that consist of dark green chloritized glass that is deeply weathered than the light coloured pillow interiors.

Unit 1c Pillow Breccia

Pillow breccia within the iron-poor, magnesium tholeiites occurs at the top of the main pillowed lava flow. Many of the pillow-breccia units grade into a solely hyaloclastite bed. The hyaloclastite material surrounding the pillow fragments is tuffaceous, light green in colour, aphanitic, siliceous and contains fragments and shards of chloritized tachylite. As soon as the hyaloclastite unit contains pillow fragments the unit

constitutes a pillow-breccia, that is 10cm. to 50cm. thick and is generally associated with the pillow-lava flow. Where a pillow lava grades into a pillow breccia the pillows become smaller and vary widely in shape and form and are highly fragmented. The proportion of pillow fragments to hyaloclastite matrix ranges from 20 to 80 percent.

Unit 1d massive (2-5 mm.) coarse-grained

The massive iron-poor basalt has a smooth light-green partly rusted weathered and fresh surface. Compositionally the iron-poor basalts are similar to the iron-rich basalts except for the magnetite content. Near to the syenite dike contacts the rocks are chloritized, epidotized and partly silicified where cut by quartz veining.

Unit 1e hyaloclastite, fragmental

Thin hyaloclastite, fragmental units are characteristically interlayered with the lava flows. Many of the pillow lavas grade upward into hyaloclastite composed solely of angular and irregular chloritized glass shards and globules, 0.1 cm. to 2cm. in size. The hyaloclastite is aphanitic, light green and siliceous with a rough, rusty brown weathered surface. As soon as the hyaloclastite unit contains pillow fragments, the unit constitutes a pillow-breccia.

Unit 1f Amygdaloidal

This unit occurs towards the top of the flows and consists of a homogeneous fine-grained basalt. The amygdules are from 0.5 mm. to 2 mm. in size and are filled with quartz and occasionally calcite. This unit is weakly magnetic.

Unit 1h Variolitic

Pillow lavas in the iron-poor tholeiitic flows are variolitic in places. The variolites range in size from 1 to 6 mm., and typically occur as two or more concentric layers just beneath the chilled rim of the pillow. On the outcrop, the variolites show up as grey knobs more resistant to weathering than the enclosing matrix of fine-grained basalt. The variolites are composed primarily of plagioclase crystals.

Unit 1i Chlorite-schist

The sheared iron-poor tholeiites are strongly foliated, medium green in colour and are comprised primarily of chlorite and carbonate. This unit occurs along the iron-poor tholeiite contacts or where the iron-poor tholeiitic basalt is in contact with a metasediment. Trace amounts of disseminated pyrite occurs within these zones.

STRUCTURAL GEOLOGY

The sedimentary rocks and the pillowed flows show strike directions from 050° in the southwestern part of the property to 195° throughout the remainder of the property with dips generally nearing vertical. Evidence for major folding occurs in the southwestern portion of the property just north of Sesekinika Lake from pillow top determinations and from the aeromagnetic map of the area, where the iron-rich tholeiitic basalts are traceable around the nose of a fold in this locality.

Several areas of highly sheared, fractured, carbonatized, silicified, epidotized and quartz-veined zones occur on the Maisonville-Grenfell township property. These shear zones are listed below:

<u>Location</u>	<u>Attitude</u>	<u>Occurrence of shearing and faulting</u>
1) Area 1	185°/80E	sediment-mafic horizon, stringer pyrite
2) Area 2	020°/vert	sediment-mafic horizon, stringer pyrite, quartz veining
3) Area 4	175°/80E	sediment-mafic horizon, stringer pyrite, quartz veining 175/80°E
4) Area 5	175°/vert	fault zone along iron-rich and iron-poor tholeiite contact-quartz veining 110°/vert
5) Area 8	185°/55°W?	
6) Area 9	180°/vert	syenite-mafic contact-quartz veining
7) Area 10	170°/20°W	magnesium tholeiite (assoc syenite dikes)-

quartz veining 170°/20°W

- 8) Area 15 175°/70°E
- 9) Area 16 165°/
- 10) 120m. W of No. 180°/vert iron-rich, iron-poor tholeiite contact
 1 of 778378
- 11) 200m. W of No. 180°/vert
- 1 of 79887
- 12) 220m. W of No. 175°/vert
- 1 of 802347

The location of many faults can be detected from lineaments that show up on the air photographs of the area. These lineaments are long linear valleys and depressions occupied by streams. The lineaments probably represent narrow shear zones or faults which were preferentially eroded by glaciation action. Where exposed in bedrock, the faults and shear zones are less than 15 m. wide with minor chlorite, carbonate, epidote, hematite and sericite with associated sulphide mineralization.

Both the tholeiitic basalt and the syenitic intrusions are fractured into three regular, strong joint directions and several well-developed irregular fracture directions that branch from the strong joint sets. The joint sets in both lithologies are filled with quartz veins. After the syenitic intrusion event the rocks cooled to fracture in a brittle manner.

Field observations indicate that the strongest joint directions from most to least prominent are as follows:

165-175° / 60-70°W
180-185° / vertical
100-105° / 65-75°NE
170-175° / 20-35°E
062-085° / 70°SE to vertical
125-135° / vertical

The north-south shallow and steeply dipping and the southeast steeply dipping veins are filled with dilatent quartz-carbonate with associated gold, molybdenite, pyrite and chalcopyrite.

The fold structure of the map-area is largely determined from attitudes of pillows in lava flows, and from the strikes and dips of contacts between massive flows. Additional information is derived from individual flows and fragmental and sedimentary units which can be used as marker horizons in the volcanic sequence. Pillows, amygdules, and hyaloclastic textures are undeformed throughout the sequence, indicating that no deformation from folding or faulting has affected rocks in the map-area. Features such as variations in thickness and continuity of flows are largely depositional features rather than structural features.

Many of the map-units and individual flows within the map-units of the tholeiitic sequences are continuous for several kilometres and have relatively constant thicknesses. The sedimentary units are discontinuous and may in fact be pockets or lenses rather than continuous beds.

The tholeiitic volcanic rocks on the Glen Auden Resources Limited property occupy the southern part of the southern limb of a synclorium which occurs north of Kirkland Lake and extends eastward to Quebec.

ROCK ALTERATION AND METAMORPHISM

Rocks on the Maisonville-Grenfell Township, Glen Auden Resources Limited property have been affected by subgreenschist regional metamorphism. In many places the regional metamorphism is partly or wholly obscured by a deuteric alteration among the rocks that varies from flow to flow. The amount of alteration also depends on the type and composition of a particular flow.

In the least affected, iron-rich tholeiitic basalt flows, late magmatic hornblende and primary pyroxene are preserved with primary plagioclase in the massive and pillowed lavas. In other flows, the plagioclase is completely replaced by clinozoisite, epidote and calcite. In the pillowed flows, alteration of pyroxene to chlorite or uralite may or may not accompany the plagioclase alteration, and in the massive flows primary hornblende remains unaltered. Secondary quartz, calcite and chlorite occur in the amygdules in the amygdaloidal basalt flows.

In the iron-poor basalts, the plagioclase is replaced by fine-grained clinozoisite, epidote, calcite and albite, especially in the pillowed lavas. Pyroxene may or may not be replaced by chlorite and albite. The replacement of plagioclase

by light-coloured epidote and clinozoisite is responsible for the hardness and the lighter colour of the iron-poor tholeiitic basalts.

Locally, the volcanic rocks are affected by higher hornfelsic grades of metamorphism, particularly near the felsic syenitic dikes. Aureoles from 2 cm. to 300 m. occur around the dikes and more prominently around the longer plug-like bodies. The volcanic rocks in the aureoles are hornblende-hornfelsed and are composed of porphyroblastic diopside and biotite with plagioclase, quartz, magnetite and hornblende. The plagioclase is altered to epidote, kaolinite and albite.

Outward from the syenite dike contacts, the rocks of the hornblende-hornfels facies grade into rocks of epidote (albite) hornfels facies which are homogeneous rocks composed of actinolite, calcite, albite, quartz and epidote according to recent thin section examination (Jensen O.G.S., 1983). Numerous irregular spider-web veining, 1 mm. to 1cm., consisting of epidote, quartz, calcite, and pyrite cuts the metavolcanics. Garnet as well as specularite, quartz, massive epidote, with minor pyrite and chalcopyrite occur at the contacts.

Towards the Winnie Lake stock just east of Kapikita Lake in the eastern part of Maisonville Township, the metamorphic aureole of epidote-albite-hornblende occurs within the iron-poor, magnesium tholeiites.

Alkali metasomatism is common in shear and fault zones especially along lithologic contacts where and easy locus for shearing occurs.

ECONOMIC GEOLOGY

A number of gold occurrences are already known on the Glen Auden Resources Ltd., Maisonville-Grenfell Township property which can be categorized in a variety of settings. The area was first prospected in the early days of the Kirkland Lake gold rush since some aspects of the geology were similar to Kirkland Lake geology. Quartz veins hosted in coarse-grained iron-rich tholeiites with associated syenite porphyry with associated gold-molybdenite mineralization occurs frequently on the property. This type of zone can create an extensive stockwork setting which would be a target for large tonnage-low grade gold deposits. Pyrite mineralization often occurs along the contact of the quartz-veins or veinlets as well as in the alteration zones away from the veins. Highgrading of specific veins within this type of setting appears to be what happened throughout the property in the old days.

A second type of gold occurrence is quartz stockwork zones within syenite intrusives. A pervasive hematization and silicification is often associated with these zones together within pyritic haloes around the veins, that are often auriferous. These are often accompanied by molybdenite and

pyrite mineralization.

A third type of gold occurrence is quartz-stock work zones associated with major North-south trending fault zones. Tensional vein zones trending roughly east-west occur near these north-south trending structures. Pyrite mineralization often occurs throughout the brecciated, rolled veins as matrix pyrite comprising up to 5% of the rock in places.

Banded chert horizons with stratiform-type, disseminated and stringer pyrite represent interflow exhalative chemical sediments which also are targets for large tonnage gold deposits. Several such horizons together with cherty tuff, cherty oxide facies and crystal tuff interflow units occur on the property.

Material mined from the past producing Gold Mines, mainly the Bennett Mine, at the Maisonville-Grenfell township property consisted mainly of quartz veins in silicified tholeiitic basalt, in syenites and in shear zones with tensional quartz veins in syenite and chert. Mineralization consisted of disseminated and stringer pyrite, chalcopyrite and gold. Minor molybdenite, stibnite, silver and gold tellurides and galena were also reported.

The ore zones around the past producing gold mines also show varying degrees of hematization, carbonatization, silicification, epidotization, and chloritization.

The better grade material appears to have been associated

with well silicified sections along the contacts of the syenite-tholeiite or the tholeiite-sediment contacts which contain more pyrite, and intense quartz-carbonate veining. The location of ore-grade material in these past producing mines and on the remainder of the Glen Auden Resources Limited property appear to show a strong preference for certain structural and lithological zones such as sheared sediment iron-rich tholeiitic basalt; sheared cherty oxide iron formations containing quartz veins; major quartz-molybdenum veins steeply dipping striking northeast that have narrow alteration haloes that are pyritiferous and auriferous; in bleached syenites (porphyritic syenite) that are cut by quartz veins with red alteration haloes that are pyritiferous and auriferous; along the crystalluff-tholeiitic basalt contact where the crystal tuff is fractured, cut by quartz veins with red alteration haloes that are pyritic; in major north-south vein systems that have been brecciated or rolled from faults with associated intense epidotization of wallrock with stringer veins subparallel to the strike of the main veins; in roughly east west trending veins dipping steeply south containing molybdenite and pale pyrite; in steeply dipping shear zones striking 160° and dipping 70° to the southwest.

TABLE 2 Description, Location, Results of Lithogeochemical Samples

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
67851	60m W of No. 1 of 802749 near old	2-3% coarsely disseminated pyrite in coarsegrained hi- iron tholeiitic basalt (Unit 2d)	<10/<.2
67852	near old trenches 30m W of No. 1 of 802749	2-3% coarsely diss. pyrite in (Unit 2d)-associated 1cm rusty quartz stringers	20/1
67853	50m W of No. 1 of 802749 old trenches	same as above	50/<.2
67854	10m N of TL 4800N at 7 + 60 E	weakly silicified, carbon- atized, weakly hematized fine grained high iron tholeiitic basalt, (Unit 2a) with 2% finely diss. pyrite -assoc. syenite intrusion	50/.02
67855	CL 801878 old trench 40m W of No. 1 of 801878	Shear zone with associated quartz stockworking within high iron, tholeiitic basalt, trace finely diss. pyrite; assoc. syenite intrusion	140/1.4
<hr/>			
<u>AREA 10</u>			
67856	CL 801878 240m S of No. 1 post of 801877	6cm wide grey quartz vein within syenite intrusion, trace finely diss. pyrite	20/<.2
67857	(old pit) 120m W of No. 1 of 801877	Shear zone with up 3% fine- ly diss. pyrite assoc. with vein selvage (Unit 2a)	60/2.6
67858	CL 801877 (old pit) 145m W of 801877 20m S of 801877	Shear zone with up to 5% diss. pyrite and pyrite along fractures within silicified (Unit 1a)	10/<.2
67859	CL 801877 (old pit) same as 67853	40cm chip sample across shear zone within iron tholeiites and assoc. 1-4cm quartz veins	50/<.2

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
67860	CL 801877 (trench) 180m W of No. 1 801877 20m S	flyrock of milky white quartz vein (6cm) with trace to 1/2% diss. pyrite; assoc. syenite dike trending E-W	60/<.2
67861	CL 801877	6cm wide quartz vein within high iron tholeiitic basalt; assoc. with nearby syenite intrusion	50/<.2
<hr/>			
<u>AREA 16</u>			
67862	220m E of No. 1 post of 801878	oxidized shear zone (12cm-24cm) sericite clay schist; assoc. 4cm-8cm vuggy quartz veining; trace finely diss. pyrite (veining 020°/80NW)	11,000/22.0
<hr/>			
<u>AREA 17</u>			
67863	(old pit) 43m E of No. 1 post of 803559	flyrock of vuggy vein (12cm) with 10% wallrock of silicified, pyritized basalt	50/<.2
67864	trenches located 30m E of No. 1 of 803559	flyrock of coarse crystalline quartz vein with 10% highly silicified, pyritic wallrock near syenite dike	140/19.4
67865	trenches at 10m north of No. 1 of 67864	Grab of alteration envelope (90%) around vuggy quartz vein (10%-of sample); 10-15% coarse cubic pyrite in a chloritic mafic host; near syenite dike	360/11.4
67866	old shaft at 272m E of No. 1 of 803559	70:40% Quartz: (carbonate vein; coarse grained (crystalline); trace pyrite, chalcopyrite, specularite (galena?) Vein is hosted in high iron tholeiitic basalt	<10/<.2
<hr/>			

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
67867	On L 1600 E at 4300 N	Coarse calcite veins (4-6cm) within high iron tholeiitic basalt; assoc. syenite stringers; contain up to 3% coarsely diss. chalcopyrite, pyrite.	<10/<.2
67868	160m W of No. 1 of 798868	fracture zone within iron- poor tholeiitic basalts; 2-3% stringer pyrite along fractures; silicified wall- rock	<10/<.2
67869	Old pit in SW corner of CL 798866	flyrock of quartz (carbonate) vein with trace pyrite; within unaltered Iron-poor pillowed tholeiitic basalt	<10/<.2
67870	Old trench in centre of claim 798866	flyrock of quartz: epidote (50:50%) vein material; de- void of sulphides; altered silicified wallrock-coarse needles of epidote	40/<.2
<u>AREA 4</u>			
67871	20m E of line 1600 E 3240 N	contact of high iron tholeiitic basalt with sheared sediments	20/<.2
67872	15m E of L 1600 E/32 + 50N	silicified, foliated sheared metasediments with 3% finely disseminated pyrite	<10/<.2
67873	20m E of L 1600E/ 32 + 60N (old pit)	Sheared wallrock (high iron tholeiites?) intensely silicified, hematized, strongly limonitic; 3-5% diss. pyrite and fracture con- trolled pyrite	10/<.2
67874	old pit 20m E of L 1600E/32 + 58N	flyrock of white, 6cm wide quartz vein; trace pyrite	120/<.2
67875	L 31 + 92N/17 + 50E	24cm wide quartz vein with diss. molybdenite and strong- ly pyrite mafic xenoliths	700/1.8

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
<u>AREA 12</u>			
67876	shaft at Old Bennett Mine	Quartz carbonate vein flyrock strongly hematitic along fractures strong jarosite; with trace pyrite, trace chalcopyrite	<10/<.2
67877	Old Bennett Mine	Grey to dark grey (variably magnetic chert with calcite along fractures; strong hematite stain, assoc. epidote clots; 3-5% finely diss. pyrite and trace chalcopyrite	40/<.2
<hr/>			
<u>AREA 4</u>			
67878	Old pit on line 31 + 92N/1400E	fracture zone within high iron tholeiitic basalt; carbonate along fractures; assoc. epidote; 1% diss. and fracture coating pyrite	50/<.2
<u>AREA 12</u>			
67879	trench at 42m E of No. 1 of 802745	flyrock of rusty quartz vein with 1cm selvage of strongly pyritized wallrock; 4% finely diss. pale pyrite; host is coarse (3-5mm) high iron tholeiitic basalt	1500/3.4
67880	trench-same as above	flyrock of intensely pyritized silicified wallrock adjacent to vein contact; 5-8% finely diss. pale pyrite	600/2.2
67881	trench-same as above	unaltered outcrop in trench of coarse-grained high iron basalt with 2% diss. pyrite along hanging wall of vein	<10/<.2
67882	old pit at 100m E and 100m S of No. 4 of 802744	Pyritic cherty and argillaceous sediments	<10/<.2

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
67883	20m W of L 1600 E/2730N	2 metre wide quartz vein with xenoliths of iron tholeiitic basalt; assoc. hematite stain- ing	60/<.2
67884	old pit at 100m W of O.N.R.	flyrock of massive, coarsely crystalline white quartz with trace galena, pyrite, chalco- pyrite	<10/4.4
67885	L 1600E/2840N	old pit-quartz flyrock; quartz is coarsely crystalline; vuggy; trace finely diss. pyrite; vein is 24cm wide.	50/<.2
67886	L 1600/2800N	old pit- 3-7' wide vein network; quartz is white, trace diss. pyrite; wall- rock alteration is limited to 3cm of silicified, pyritized; assoc. syenite dike.	<10/<.2
67887	trench at 35m W of No. 1 of 802745	flyrock 50:50% wallrock: quartz vein material wallrock is intensely carbonatized and pyritized and quartz vein is vuggy with trace pyrite	1750/5.2
67888	same as 67887	Sample of 90:10% quartz: wall- rock. The quartz is darker grey with finely diss. pyrite	30/<.2
67889	same as 67887	quartz vein, medium grained	1500/5.2
67890	East central claim 778379	flyrock of 15cm wide quartz vein; coarse grained, vuggy, trace pyrite with vein sel- vages altered to epidote and chlorite	1830/2.4
67891	94m W of L 1600E/2760N	Subcrop of high iron basalt with 3% coarsely cubic pyrite, 5% epidotized feldspars	10/<.2

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
<u>AREA 15</u>			
67892	70m east of No. 3 of 802334	Iron-rich tholeiitic basalt with 1% diss. pyrite	<10/<.2
67893	190m E of No. 3 post 802334	10-15cm wide quartz vein; grey to white, strong iron stain along fractures, trace pyrite with 5% heavily pyritized wallrock xenoliths	<10/<.2
67894	same as 67893	Wallrock of 10-15cm wide quartz vein, alteration in- cludes an intense epidotization carbonatization with up to 8% fine cubic pyrite. Host is Unit 2d	20/<.2
<hr/>			
<u>AREA 9</u>			
67895	L 2450/15 + 65N	Fracture zone within Iron-poor tholeiitic basalts with 1/2%- 1% fracture controlled pyrite; assoc. syenite intrusion	<10/<.2
67896	old trench at L 24 + 50E/ 15 + 60N	Iron-poor basalt injected by red feldspathic stringers, calcite stringers, up to 5% pyrite along fractures, trace as fine disseminations	240/<1.0
67897	60m E of L 2450E/ 15 + 60N	flyrock from old trench of quartz-carb vein brecciated; with 1/2% fine matrix pyrite around qc fragments in breccia	650/<.2
<hr/>			
67898	L 4800N/2630E	Silicified iron-rich tholeiite with up to 5% diss. pyrite	40/<.2
67899	25cm S of L 4800N/2440E	Iron-rich tholeiites with hematite staining and 1-2% diss. pyrite	<10/<.2
67900	25m S of L 4800N/2440E	Massive white quartz vein with epidote and trace pyrite and specularite	<10/<.2
18280	80m S of L 4800N/2700E	Cherty argillaceous sediments with stratiform pyrite (1-2%)	10/<.2

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
18281	L 4800N/2840E	old pit-of cherty argillaeous sediments with 3% pyrite as blebs and streaks parallel to bedding	<10/<.2
18282	200m E of No. 1 of 802337	as 18281 - 1% pyrite, increase in epidote stringers, cherts are slightly magnetic	<10/.2
18283	170m E of No. 1 of 802746	24cm wide quartz vein with 1% diss. Wallrock is silicified for 24cm.	<10/<.2
<hr/>			
<u>AREA 5</u>			
18284	NE corner of CL 800346	Shear zone 50% diss. pale pyrite within brecciated quartz carbonate vein (up to 30% matrix pyrite); trace diss. molybdenite	.96/oz/ton
18285	NE corner of CL 800346	Wallrock of vein; high iron tholeiite; coarsegrained; epidote along fracture surfaces	650/<.2
<hr/>			
<u>AREA 15</u>			
18286	65m W of No. 2 of 802335	Shear zone 074/80°E within high iron tholeiites; 3% finely disseminated pyrite	20/<.2
18287	L 3220E/ 400N Near old pit	Rusty 10cm wide quartz vein with 3% pale coarse cubic pyrite and diss. pyrite	70/5.2
18288	Near pit 165°/55E	50:50 wallrock: vein material; wallrock is carbonatized, chlorite, epidote with 2% finely diss. pyrite Vein is grey, coarsely crystalline with trace fine pyrite, chalcopyrite, molybdenite	40/1.6

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
18289	140m S of No. 1 of 802340 and 65m W	Quartz-carbonate vein (30cm wide); light grey coarsely crystalline with epidote, chlorite inclusions; trace diss. pyrite, chalcopyrite, and stibnite	10/2.4
18290	same as 18289 old trench	Intensely epidotized wall-rock of vein in 18289 medium grained; granular; 80% epidote; 2% diss. pyrite	30/2.0
18291	230m W of No. 2 of 802340	Tholeiitic dacite ? with diss. pyrite; carbonate and epidote along fractures	<10/<.2
18292	10m W of No. 1 post of 802339	50:50 vein: wallrock sample; vein is grey, coarsely crystalline with trace diss. pyrite: wallrock is chloritized, with 1% diss. pyrite	<10/<.2
18293	Centre of CL 802333	Silicified, pillowed high-iron tholeiite; assoc. ladder veining; trace diss. pyrite; associated syenite dike	20/<.2
<hr/>			
<u>AREA 14</u>			
18294	190m E of No. 3 post of 802336	Cherty sediments with 3% stratiform pyrite	<10/<.2
<hr/>			
<u>AREA 15</u>			
18295	73m W of No. 2 of 802336	Quartz vein (18cm wide) 142/85°NE; chlorite inclusions; trace diss. pyrite	10/<.2
18296	65m W of L 3220E/ 4200N	Quartz vein; (10cm wide); dark grey; coarsely crystalline; trace diss. pyrite; 20% inclusions of silicified, green, chloritized wallrock with trace pyrite	<10/<.2

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
<u>AREA 8</u>			
67951	100m N of No. 1 of 802346	Sediments; graphite-magnetite horizon; mixed sediment-volc. sequence; shear zone with 2-3cm grey massive magnetite beds with trace to 1/2% pyrite	<10
67952	150m W of No. 1 of 802346	Chert black; 2cm wide; 3-4% fracture controlled pyrite; 1% diss. pyrite	<10
67953	Same as 67952	Chert; lighter grey colour; 3% diss. pyrite	<10
<hr/>			
<u>AREA 11</u>			
67954	40m S of L 3190N/ 2460E	Cherty and/or silicified rock with quartz-carbonate veining; trace pyrite and chalcopyrite	265 ppb
67955	40m E of L 2450E/ 3400N	Old pit; magnesium rich tholeiite with epidote calcite along fractures; trace pyrite	<10
67956	5m S of L 3192E/ 2500E	Old pit; Cherty argillite; < 15% pyrite as 'blebs' and grains; flyrock	30 ppb
67957	4m N of L 3192N/ 2520E	Massive grey chert with 1% diss. pyrite	90 ppb
<hr/>			
<u>AREA 2</u>			
67958	L 2450E/2480E	Old pit; flyrock from shear zone with high-iron tholeiites; coarse calcite and feldspathic stringers, tr-1/2% pyrite along fractures	5 ppb
67959	20m E of L 2450E/ 24 + 80N	trench; shear zone; chloritic high-iron tholeiite with magnetite seams; 2-3% stringer pyrite	65 ppb
67960	20m E of L 2450E/ 2500N	Silicified, banded sediment; calc-silicate? 4-5% stratiform pyrite within shear zone	80 ppb

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
<u>AREA 7</u>			
18297	100m N of No. 1 of 802368	Quartz vein; white to light grey; coarsely crystalline with trace disseminated pyrite	210/<.2
18298	old pit 180m N of No. 1 of 802368	Quartz vein (45cm wide); trace to 1% diss. pyrite; trace diss. molybdenite	270/<.2
<hr/>			
<u>AREA 3</u>			
18299	150m S of No. 3 of 778378	Chert, banded with pyrite and epidote bands; 2% pyrite as fine cubes and stratiform bands	90/<.2
18300	30m E of No. 1 of 182991 location	Chert with 1% pyrite; massive	40/<.2
<hr/>			
67644	76m E of L 1600E/ 2120N	Intensely silicified low iron tholeiite; qtz stringer zone; pervasive epidotization (25cm) 2-3% pyrite; assoc. syenite dike (5cm)	15 ppb
67645	190m E of L 1600E/ 2120N	Old pit, brecciated, rehealed fracture zone; veins have trace pyrite, molybdenite, malachite. Wallrock is carbona- tized and silicified.	<10
67646	as 67645	90% carbonatized, silicified high iron tholeiite with trace pyrite; assoc 10% quartz stringers	<10
67647	10m N of S CL line of 802746 (near power line)	Old pit; high iron tholeiite with 1% sulphides; hematite, carbonate, quartz	<10 ppb
67648	30m N of S CL of 802746	Old pit; iron tholeiite with trace pyrite; epidote; carbonate, quartz	35 ppb
67649	225m E of No. 3 of 802337	Old pit; iron tholeiite with 1-2% pyrite; hematite; epidote	<10

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
67961	20m E of L 2450E/ 2450N	fracture zone within high iron tholeiite with coarse calcite, epidote, chlorite and 1% diss. chalcopyrite	15 ppb
67962	as 67961	Breccia zone with intense quartz-calcite stringers wallrock was silicified, epidotized, carbonatized and hematized with coarse pale pyrite	30 ppb
67963	140m N of No. 1 of 778378	Fractured iron-poor tholeiite with 2-3% diss. pyrite and fracture controlled pyrite	<10
<hr/>			
<u>AREA 6</u>			
67964	Old pit on power-line 10m E of No. 2 of 778378	Quartz-stringer zone with sample of silicified, carbonatized host rock and 5-10% coarse diss. pyrite within crystal tuff	<10
67965	L 1600E/1170N	Silicified crystal tuff with 4% diss. pyrite cut by 1cm wide quartz stringers	<10
67966	L 1600E/1170N	5cm wide white, vuggy quartz vein, 2% pale pyrite	10
67967	200m W of No. 1 of 798877 L 5192E/ 31 + 90N	Unit 1a, iron-poor tholeiites injected by fine ep-quartz stringers with diss. pyrite	
67968	Same as above	75cm wide prevasively epidotized rock along fault	
<hr/>			
<u>AREA 1</u>			
67969	50m W of No. 1 of 798878	Sheared, brecciated chert, shaley argillite horizon with up to 5% stringer pyrite	
67970	240m W of No. 1 of 798878	Fractured iron-poor tholeiite with 2% stringer pyrite	

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
67971	E of Pit 2-1	Iron-poor tholeiitic basalt with 4% stringer pyrite	
67972	Trench 4-1	Siliceous sediment horizon with massive pyrite seams, diss. pyrite, and stringer pyrite	
67973	Pit 1-1	Sheared, Brecciated, chert with chlorite-carbonate slips and 5-10% stringer pyrite	
67974	Pit 1-1	Black chert, 8% stringer pyrite	
67975	TR 1-1	Grey fractured chert, 5% stringer pyrite	475 ppb
<hr/>			
67976	210m N of No. 3 of 798875	Iron-rich tholeiitic basalt, silicified with 3% stringer pyrite	
67977	L 1576N/2480E	Iron-poor tholeiite next to porphyritic syenite dike, country rock silicified, epidotized with 2% pyrite and trace chalcopyrite	110 ppb
<hr/>			
<u>AREA 9</u>			
67978	10m S of L 1576N/ 2510E	Intensely epidotized zone within iron-poor tholeiite, tr pyrite	300 ppb
67979	L 1576N/2515E	Massive pyrite lense with 40% pyrite, 1m wide	
<hr/>			
67980	L 1560N/3100E	2m wide shear zone with 3-4% stratiform pyrite bands	
67981	100m W of No. 1 of 801219	Silicified, hematized, epidotized iron-rich tholeiite with trace pyrite and chalcopyrite	
67982	L 1576N/3660E	Strongly foliated, iron tholeiite with chlorite-carbonate slips, quartz-carbonate stringers, tr pyrite, chalcopyrite	50 ppb
<hr/>			

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
<u>AREA 5</u>			
67983	40m N of L 1576N/ 3660E	Intensely silicified tholeiitic basalt with silica-chlorite stringers with trace pyrite	
67984	90m N of L 1576N/ 3660E	Silicified tholeiitic basalt, chlorite-carbonate slips with coarse calcite, hematite stringers, trace pyrite	
67985	90m N of L 1576N/ 3660E	Sericite schist-sheared hyaloclastite material, trace pyrite	
67986	140m N of 1576N/ 3660E	Quartz-fracture zone with quartz stringers with pyritic host rock fragments	
67987	Same as 67986	Wallrock material of vein, iron-poor tholeiite is silicified, epidotized, car-ized with 4% finely diss. pyrite	
<hr/>			
<u>AREA 9</u>			
67988	65m E of L 2450E/ 1300N	Pyritic material in iron tholeiitic basalt, 5% diss. pyrite	115 ppb
67989	10m E of L 2450E/ 1170N	Quartz vein with 50% WR: 50% vein, 4% coarsely diss. pyrite	11,500 ppb
67990	L 2450E/1170N	Iron-rich tholeiite with 3-4% stringer pyrite	
67991	70m E of 1600E/ 1380N	30-45cm wide white bull quartz; trace pyrite	
67992	70m E and 40m S of L 1600E/ 1380N	Iron-poor pillowed basalt, strongly epidotized, silicified with trace pyrite	
67993	70m E of L 1600E 1380N	Quartz vein with 10% wallrock inclusions with 3% pyrite	

SAMPLE NO.	LOCATION	DESCRIPTION	Au Ag ppb/ppm
<u>AREA 1</u>			
67994	AREA 1	Intensely silicified, carbonatized sediment with 5% stringer pyrite	
67995	AREA 1	Silicified sediments; 3% finely diss. pyrite	
<hr/>			
<u>AREA 2</u>			
67996	L 2450E/2480N	Silicified sediment, sheared, epidote-carbonate along slips tr. pyrite	
67997	L 2450E/2470N	Strongly foliated iron-rich tholeiitic basalt with 10% stringer pyrite	
<hr/>			
67998	AREA 6	Flyrock from pit, strongly pyritic iron-rich tholeiite with 5-7% diss. pyrite	
67994	CL 803558	Weakly sheared iron-rich tholeiite with 5-7% diss. pyrite	
68000	CL 803558	12-20cmwide shear zone within iron-rich tholeiitic basalts, assoc. drusy 2cm qtz-ep veining	
<hr/>			
<u>AREA 16</u>			
67906	AREA 16	Grey cryptocrystalline vein with assoc. sheared pyritic iron rich tholeiite	
67907	AREA 16	Cherty-oxide iron formation sub-parallel to shear zone, assoc. epidote-garnet-calcite stringer	

AREA 1 (see Figure 9)

In the eastern part of the Maisonville-Grenfell township property north-northwest of Kapakita Lake a series of trenches and pits occur. The main pit is in a shear zone along the chert-tholeiitic basalt contact. The banded chert zone represents an exhalative chemical sediment horizon that has undergone an intensive period of fracturing with associated alteration including sericitization and epidotization. Previous assays from prospector C. Forbes (1984) gave values up to 143 oz. Au/ton from a grab sample of banded chert with stratiform-type pyrite. During the geological mapping a sample of grey, fractured chert with 5% stringer pyrite returned 475 ppb Au.


A pace and compass grid was put in for better control in the mapping and sampling of these pits and trenches. The No. 2. post of claim 798875 is the 0+00N/0+00W point for the grid.

The main pit (PIT 1-1) is in a strongly foliated tholeiitic basalt (Unit 2d) with chlorite-carbonate slip surfaces and up to 4% stringer pyrite. The basalt is composed of actinolite, chlorite, biotite, feldspar, pyroxene and magnetite. Varying quantities of epidote, quartz, and calcite are dispersed throughout the rock. Footwall to the basalt is a sheared chert-cherty tuff horizon with up to 10% stringer and stratiform pyrite. The chert horizon here is striking roughly 185° and dips

	Au (ppb)	Mo (ppm)	As (ppm)	Zn (ppm)
67969		50	21.5	1100
67972		45	10.5	
67973		100	35.1	900
67974			65.2	900
67975	475	160	21.5	270
67994			38.3	1500



798875



REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for	GLEN AUDEN RESOURCES LIMITED	
	Title	Detailed Geology AREA 1	
		Fig. 9	
	Date: Oct 1985	Scale: 1:100	N.T.S.:
	Drawn: NC	Approved:	File: M-78

steeply to the east.

In the trenches north of PIT-1, a grab sample returned 475 ppb or .146 oz./ton Au. from a grey chert with 5% stringer pyrite. Slate, cherty tuff and a pyritic laminated siliceous sediment?, also occur here. The chert horizons in this area are mixed chemical and volcanoclastic sediments. The cherts are often banded, the banding consisting of varying proportions of chert, sulphide, carbonate and epidotized tuffaceous laminae, with thicknesses ranging from .1mm to 4mm.

In the trenches to the south of PIT-1, the chert horizon is at least 1 metre wide and contains up to 8% stringer pyrite. A 5 metre wide crystal tuff horizon also occurs here, hanging wall to the chert. The tuff is strongly hematized and contains up to 2% disseminated pyrite with up to 30% subhedral broken feldspar crystals ranging from 1mm. to 4mm. in diameter.

In the trench south of here, (TR5-1), the sheared pyritic chert horizon is slightly narrower and contains 8-10% stringer pyrite. Finally in the last trench, TR6-1, the chert appears to have pinched out leaving the crystal tuff hanging wall to the strongly foliated sheared iron-rich tholeiite.

Both west and east of the sheared sediment horizon the iron-rich tholeiites are intensely fractured and contain fine quartz-carbonate stringers with associated garnet-epidote-carbonate along the fractures with up to 3%

stringer pyrite. The mineralogy implies that upper green-schist facies metamorphism occurs here.

Disseminated sulphides coupled with stratiform-type and stringer sulphides are associated with this stratabound exhalative chert horizon which can be traced using induced-polarization methods.

Weakly to moderately anomalous copper values from 240 to 560 ppm. occur within the sedimentary chert-argillite horizon with associated stringer pyrite. The brecciated iron-tholeiite also returns values up to 560 ppm. copper. Similarly, moderately to strongly anomalous zinc values up to 1500 ppm. occur within both the fractured chert and basalt sequences containing 5-8% stringer pyrite. Weakly anomalous lead values up to 20 ppm. also occur here.

Four out of five samples of the brecciated chert horizon along its 100 metre length return strongly anomalous in molybdenum, up to 160 ppm. There also is an important enrichment in arsenic reaching values of up to 65.2 ppm. within the cherts. These two elements are important as pathfinder elements in the Kirkland Lake, Matachewan Matheson and Timmins gold areas.

AREA 2 (see Figure 10)


On line 2450E between 2480N and 2520N, a series of trenches occur in a sheared fractured, siliceous sediment horizon in

contact with an iron-rich tholeiitic basalt. Weakly anomalous gold values up to 80 ppb. occur here within the laminated, siliceous sediment containing 4-5% stratiform pyrite; a highly fractured iron tholeiite with 40% stratiform pyrite; in a highly fractured iron tholeiite with 4% stringer pyrite. A small pace and compass grid covers the area for better control in mapping and sampling.

The main shear zone is striking 025° and is dipping vertically. A grey, sugary textured quartz vein parallels the sediment horizon and varies from 40-50cm. in width. The vein contains calcite-chlorite-epidote-hematite coated fracture surfaces. Mineralization consists of up to 2% finely disseminated pyrite.

Approximately 100 metres to the south of these trenches an east-west trench occurs that appears to have been searching for the southern extension of the shear zone. The trench is in silicified, epidotized, hematized iron-rich tholeiite. It appears that the shear zone is either to the east or west of this trench hidden by overburden cover.

All of the samples taken here return moderately to strongly anomalous in molybdenum, up to 116 ppm. Weakly to moderately anomalous copper and zinc values up to 740 ppm. and 240 ppm., respectively occur within both the fractured basalt with up to 3% stringer pyrite and in the fractured laminated sediment with up



	Au (ppb)	Mo (ppm)	Cu (ppm)
67958			
67959	65	116	550
67960	80	44	740

REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for	GLEN AUDEN RESOURCES LIMITED	
	Title	Detailed Geology AREA 2	
		Fig. 10	
	Date: Oct 1985	Scale: 1:100	N.T.S.:
	Drawn: NC	Approved:	File: M-78

to 5% stringer and stratiform pyrite. Weakly anomalous gold values up to 80 ppb. occur within the laminated siliceous sedimentary horizon.

AREA 3

An east north-east trending cherty sulphide facies iron formation occurs 200 metres north of the No. 2 post of 802747. The chert horizon is a 1-3 metre wide bed that occurs stratigraphically along the variolitic iron-poor magnesium tholeiite and the iron-rich tholeiite contact. The chert can be traced eastward from the claim line for 60 metres. The strike varies from 065° to 080° and dips vertically. Two rock grab samples were taken here and both returned weakly anomalous gold values of up to 90ppb.

The northern contact is the variolitic basalt (unit 1h) with an intense silica-epidote alteration along hairline fractures and an associated pink carbonate. Minor amounts of disseminated pyrite occur throughout the matrix. The chert (Unit 3a) is grey, aphanitic with 1-3% disseminated, stratiform, and stringer pyrite. The chert is moderately fractured and is semi-banded to massive with epidote-pyrite bands and epidotized, carbonatized fracture surfaces. The southern contact of the chert is the coarse grained iron-rich tholeiitic basalt (Unit 2d). The rock is strongly magnetic and contains spider-web silica-epidote

stringers striking 080°/80S and 025°/80W. It appears that the chert may pinch out to the east and is lost in the overburden to the west.

Weakly anomalous copper up to 310 ppm. and moderately anomalous zinc values up to 590 ppm. occur within the chert horizon. The banded variety of chert, the banding consists of epidote and carbonate probably tuffaceous in origin, carries the anomalous zinc values together with slightly higher than background copper values. The weakly anomalous gold value (85 ppb) also occurs in the banded variety as compared to 40 ppb. Au in the massive variety. A significant enrichment in tungsten with moderately anomalous values up to 14 ppm. occurs in both the banded and the massive chert varieties. Weak to moderately anomalous molybdenum values from 16-30 ppm. occur in the banded and the massive varieties, respectively. Finally, weakly anomalous arsenic values up to 5 ppm. occur in both varieties of chert.

The anomalous tungsten and molybdenum values are important as indicator elements to known gold deposits in the area.

AREA 4 (see figure 11)

In the southwest corner of claim 798864 on line 1600E from 3220N to 3260N a north-south trending sheared laminated, siliceous sediment (Unit 3c) occurs within an iron-rich

tholeiitic basalt flows (Unit 2h, Unit 2d).

Weakly anomalous gold values up to 115 ppb. occur here associated with quartz veining within the iron rich tholeiitic basalt. The shear zone appears to be striking 175° and dips near vertical. The veining is subparalleling the main shear direction at $175^{\circ}/75-80^{\circ}\text{E}$ with minor veining at $080^{\circ}/85\text{S}$.

The shear zone is confined to the contact between the massive coarse-grained iron-rich tholeiitic basalt (Unit 2d) and the laminated, siliceous metasediment (Unit 3c). Shearing is indicated by strongly foliated dark green basalts with chlorite-carbonate slips, stringer pyrite, epidotization and hematization. Up to 3% disseminated pyrite occurs within the coarse grained tholeiite that is cut by an erratic quartz stringer zone associated with the shear zone.

The siliceous sediments occur as an 8 metre wide bed with an aphanitic, siliceous matrix with pale green, maroon, beige and purple 1-2 mm. laminae. Up to 4% pyrite mineralization occurs as stratiform and stringer types with associated carbonate, hematite and limonite coatings along fractures.



Minor quartz stringers from 3-5 mm. in width strike 158° and dip 070° to the west and occur within the sheared siliceous sediment horizon. The sheared sediment horizon can be traced for 40 metres until it is lost in the overburden. Approximately 200 metres east of the shear zone in Area 4, on the southern claim



	Au (ppb)	Mo (ppm)	W (ppm)
67871	15	64	
67872	5	13	12
67873	10	43	
67874	115	8	
67878	700	47	

798863

798864




REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for GLEN AUDEN RESOURCES LIMITED		
	Title Detailed Geology AREA 4		
	Fig. #		
	Date: Oct. 1985	Scale: 1:1000	N.T.S.:
	Drawn: NC	Approved:	File: M-78

line of claim 798864 a 24 cm. wide roughly northeast striking vein occurs dipping 046° NW. The vein contains 25% strongly pyritic host rock inclusions and disseminated molybdenite. Strongly anomalous gold values occur within this vein up to 700 ppb. This vein has previously been pitted to uncover it along strike but the pits are filled with overburden cover.

Disseminated sulphides coupled with stringer sulphides within both the sedimentary horizon and the fractured iron-rich tholeiite can be traced using induced polarization method.

Two weakly to strongly anomalous gold values occur within quartz veins with disseminated molybdenite and trace amounts of pyrite. The contact of the iron-rich tholeiites and the sediments returns strongly anomalous molybdenum values up to 64 ppm. and weakly anomalous copper up to 260 ppm. The sheared sedimentary horizon contains weakly anomalous lead, up to 102 ppm., weakly anomalous molybdenum, up to 13 ppm. and moderately anomalous tungsten, up to 12 ppm. The sheared portion of the iron-rich tholeiite with an intense silicification and hematization with 5% pyrite returns strongly anomalous molybdenum up to 43 ppm. All five rocks grab samples return weakly to strongly anomalous in molybdenum ranging from 8 ppm. to 64 ppm. This is an important indicator element to various gold mines in the area.





	Au (ppb)	Mo (ppm)	W (ppm)
67871	15	64	
67872	5	13	12
67873	10	43	
67874	115	8	
67878	700	47	

798863

798864

LIGOE

3300N
3280N
3260N
3240N
3220N
3200N



REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for	GLEN AUDEN RESOURCES LIMITED	
	Title	Detailed Geology AREA 4	
		Fig. II	
	Date: Oct. 1985	Scale: 1:4000	N.T.S.
	Drawn: NC	Approved:	File: M-78


AREA 5 (see fig. 12)

In the south-eastern corner of the Maisonville-Grenfell township property in the northeastern corner of claim 800346 a series of trenches occur. The trenches are following east-west tensional veins along a major north-south trending fault zone along the contact between the iron-rich and the iron-poor tholeiites. This fault zone shows up on the aerial photographs as a strong lineament along a linear depression occupied by a stream.

In the outcrop the shearing is indicated by a strong foliation 165° - 175° , dipping vertically. A major fracture system striking approximately 110 - 120° dipping 50° to vertically is prominent that is injected by auriferous quartz veins. In the vicinity of the shear zone the iron-poor magnesium pillowed tholeiites are strongly silicified with chlorite-carbonate-epidote-hematite along fractures.


On line 1680N at 360E an old trench occurs in a fracture zone with quartz-carbonate stringers following two main directions $050^{\circ}/67\text{NW}$ and $012^{\circ}/52^{\circ}\text{W}$.


Strongly anomalous gold values up to .96 oz/ton gold occur north of here, west of the beaver pond (see fig. 12) in a previously trenched brecciated, rolled 100 cm. wide quartz-pyrite-molybdenite vein. The vein contains up to 60%



800346

BEAVER DAM





	Au(ppb)	Mo(ppm)
18284	.96oz/ton	44
18285	650	23
67986	365	80
67987	10	150

REVISIONS

ROBERT S. MIDDLETON
EXPLORATION SERVICES INC.

for GLEN AUDEN RESOURCES LIMITED

Title
Detailed Geology
AREA 5

Fig. 12

Date: Oct 1985	Scale: 1:100	N.T.S.:
Drawn: NC	Approved:	File: M-78

matrix pyrite and fine grey disseminated molybdenite. The wallrock of this vein returned 650 ppb. Au.

A pace and compass grid was put in over the shear zone with flags every 10 metres. The shear zone is striking 170° and dips near vertical while the auriferous veins are striking from $110-120^{\circ}$ with numerous parallel and branching veins and veinlets. Considerable chloritization, silicification and carbonate alteration occurs in the vicinity of the quartz veins and up to 200 metres in length and width of quartz stockwork may occur here. Another gold bearing vein approximately, 200 meters south-southeast of the main vein west of the beaver pond, occurs within iron tholeiite. The vein strikes 112° and dips 52° northeast. The vein here is 30 cm. wide and contains up to 4% pyrite. Moderately anomalous gold values up to 365 ppb. occur here. This area, therefore may be a 200 metre x 150 metre wide quartz stockwork zone and further work is warranted.


Strongly anomalous molybdenum values up to 150 ppm. occur within the wallrocks of the veins while the veins themselves carry value up to 80 ppm. Mo. The molybdenum-gold association here is obvious whereby anomalous gold values are coincident with anomalous molybdenum values.

AREA 6 (see fig. 13)

In the south western part of the property, on line 1600E at

1160N several pits and trenches occur in fractured iron-rich tholeiitic basalt and silicified crystal tuff cut by quartz stringers. The crystal tuff horizon is strongly fractured and is cut by an erratic quartz stockwork system that has intensely silicified the hostrock. A major shear zone occurs 40 m. west of the crystal tuff horizon along the iron-poor and iron-rich tholeiitic basalt contact. The shear is striking 175° and dips steeply to the east. The shearing event probably formed the east-west trending tensional fractures within the more competent felsic crystal tuff that were later injected by quartz-pyrite veins.

In detail, the crystal tuff horizon is stratigraphically topping the iron-rich tholeiitic basalts and forms a bed from 10 m. to 20 m. in width. The rock consists of up to 30%, 1 mm. to 4mm. subangular broken feldspar crystals set in an aphanitic, siliceous, grey to dark grey groundmass. A pervasive hematization exists along the contact producing a pink-red rock that looks similar to the porphyritic syenite, unit 5b. Mineralization includes 5-10% disseminated pyrite along the silicified vein selvages and up to 3% pale pyrite within the quartz-carbonate stringers. The veins also contain inclusions of highly pyritic wallrock. The vein widths vary from 2 cm. to 10 cm. and strike 110°/75E and 060°/vertical, respectively. The quartz-stockwork system can be traced for 60 metres to the north



REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for	GLEN AUDEN RESOURCES LIMITED	
	Title	Detailed Geology	
		AREA 6	
		Fig. 13	
	Date: Oct 1985	Scale: 1:100	N.T.S.:
	Drawn: NC	Approved:	File: M-78

and then is lost in the overburden. The system is also lost in overburden to the south of the pit on the hydroline. Southwest of pit 1 (see fig. 13) a trench exists striking 080° within the sheared iron tholeiites.

This fracture zone occurs on a roughly E-W trending fold axis occurring in the Southwestern portion of the property. A weakly anomalous molybdenum value of 18ppm. occurs within this quartz fracture zone.

AREA 7

In the southwestern corner of the property along the eastern claim line of Claim 802748 a trench and pit occur on what used to be called the Sesekinika Townsite property (see previous history). The pit is in a fractured, silicified iron-rich tholeiite cut by porphyritic syenite (Unit 5b) with associated quartz veins striking 170°/20°W. The main vein in the pit is 50 cm. in width and contains 1% pale pyrite and molybdenite mineralization. To the south the trench is in a fractured, silicified iron-rich basalt with a 35 cm. wide quartz vein that contains molybdenite and pale pyrite. To the west the vein pinches down to 6 cm. and then is lost in the overburden. A quartz stockwork system covers an area of 200 metres x 100 metres in this location, and possibly more. Weakly anomalous gold values up to 270 ppb. occur in this area. Moderately to strongly

anomalous molybdenum up to 120 ppm. also occur here.

Work carried out in this area in the early 1900's included pitting, trenching and drilling on the Sesekinika Townsite property No. 49 (see O.D.M.N.A. Files, Kirkland Lake) by Sylvanite Gold Mines Limited.

During 1914 some free gold was found here in a narrow quartz vein dipping 15°NW.

Drilling by Erie Canadian Mines Limited, totalling 464 feet, five holes intersected a 2 foot wide quartz vein that strikes east and dips 60N.

AREA 8

In the south-eastern corner of claim 802342, on line 3220E between 3100 to 3300N, several trenches occur in sheared iron-rich tholeiitic basalts cut by porphyritic syenite dikes, with associated cherty oxide facies iron formation (Unit 3d). An intense quartz stockwork system occurs over this area, with associated disseminated pyrite. The more major vein directions are 175°/80E, 080°/vertical and 010°/55°E. The main north-south trending trench may have been following a vein, however the trench contents are now masked by overburden. To the southwest approximately 100 metres, a 5-10 metre wide cherty sulphide iron formation (Unit 3a) occurs. The horizon is striking 110° and dips steeply to the south. The chert is a dark grey to light

grey in colour with up to 4% stringer pyrite and 1% stratiform pyrite.

Moderately to strongly anomalous zinc values up to 2100 ppm. and weakly anomalous copper values up to 270 ppm. occur within the grey cherty-sulfide horizon to the south. Weakly anomalous molybdenum values up to 23 ppm. occur within both cherty oxide and the cherty sulfide facies iron formations.

AREA 9 (Sesekinika Townsite Property)

On line 2450E from 1100N to 1700N several trenches and pits occur. At 1640N the trenches are in iron poor tholeiitic basalt with associated north-south porphyritic syenite dikes and associated quartz veining. Moderately to strongly anomalous gold values from 650 to 11,500 ppb. occur within, iron-poor basalt injected by red feldspathic and calcite stringers with 5% stringer pyrite, and a brecciated quartz-calcite vein (6 cm.?) with 1/2% matrix pyrite around the quartz fragments. The vein system may be going roughly east-west here, unfortunately none were found in outcrop. All grab samples were taken from flyrock in the trenches. Going south for approximately 40 metres on TL1576N at 2550E, a 1 m. wide massive pyrite bed occurs comprised of 50% coarsely crystalline pyrite. Deep overburden cover to the north and south hindered possible strike extension of this horizon. A weakly anomalous gold value of 300 ppb. occurs within

an intensely fractured epidotized iron-poor tholeiitic basalt with trace pyrite near to this horizon. Following the iron-rich, iron-poor tholeiitic basalt horizon southward from TL1576N for 150 metres another trench occurs in the iron-rich tholeiitic basalt, with 5% disseminated pyrite. A grab sample from here yielded 115 ppb. gold.

At 2450E/1170N several more trenches occur. The trenches are in the iron-rich tholeiitic basalt unit along the contact with the iron-poor tholeiites. One sample from flyrock of a quartz vein with 50% wallrock material and 4% coarsely disseminated pyrite yielded 11,500 ppb. gold.

Strongly anomalous molybdenum values up to 120 ppm. occur within fractured iron tholeiite with 3% disseminated pyrite. A moderately anomalous copper value up to 1620 ppm. occurs with a strong arsenic enrichment up to 126 ppm (As) within the massive pyrite lense.

The strong molybdenum, and arsenic enrichment are important here as they are pathfinder elements to gold mines in the area.

AREA 10

In the northwest corner of the property south of the White-Clay River, in the northeast corner of claim 801877, several trenches, pits and one shaft occur within narrow quartz-carbonate veins that cut syenite (Unit 5c) and iron-poor

(Unit 1a). This area was worked on by Elzina Mines Limited in the early 1900's (see Previous History). The trenches and pits are in: syenite intrusions cut by narrow 6-10 cm. wide grey quartz veins with finely disseminated pyrite; roughly north south trending shear zones ($175^{\circ}/20^{\circ}W$) injected by narrow quartz veins within the iron-poor tholeiites; quartz veins in iron-rich tholeiitic basalts. The syenite intrusions seem to be trending roughly north-south along the contact between the iron-rich (Unit 2a) and the iron-poor (Unit 1a) tholeiitic basalt flows. Rock samples return weakly anomalous gold values up to 60 ppb. in the narrow quartz veins while a 50 cm. wide chip sample across the shear zone returns 50 ppb. The main vein direction is east-west dipping steeply to the south with a minor direction of 175° /steeply to the west. Mineralization includes trace to 1% finely disseminated pyrite in the veins up to 3% disseminated pyrite along vein selvages. The shear zone consists of 5% disseminated and stringer pyrite within silicified iron-poor tholeiitic basalt.

Weakly anomalous copper values and arsenic values up to 440 ppm. and 7.0 ppm. occur with weakly anomalous gold values up to 45 ppb., and with moderately anomalous tungsten values up to 10 ppm. within the narrow quartz veins.

Area 11

In the northwest corner of claim 802349 on TL 3192N at 2500E, a series of pits occur within a fractured chert-argillite horizon. The pit is in a grey chert horizon cut by quartz-carbonate stringers, within the iron-rich tholeiitic basalt sequence. Grab samples from the main pit return from 90 ppb. to 265 ppb. in gold. The fractured chert horizon is lying stratigraphically towards the top of the iron-rich tholeiitic basalt sequence near to the iron-poor tholeiitic basalt contact.

The main chert horizon represents an exhalative chemical sedimentary horizon mineralized with 3% disseminated pyrite with minor chalcopyrite. The horizon is striking roughly 190° and occurs just east of a north-south trending shear zone.

In detail the chert is grey, sugary textured with erratic fine, 1 cm. wide quartz-carbonate veining. A cherty argillite unit also occurs here with up to 15% disseminated pyrite.

Approximately 160 metres north of the chert horizon another pit occurs in strongly epidotized iron-poor magnesium tholeiite with trace amounts of pyrite.

Moderately anomalous copper values occur in this area up to 760 ppm. coincident with strongly anomalous molybdenum values of up to 230 ppm. and weakly anomalous gold of 265 ppb. within the chert horizon. Moderately anomalous zinc values up to 610 ppm.

with strongly anomalous arsenic values of 32.2 ppm. also occur within the fractured chert. One chert sample returns 480 ppm. molybdenum.

AREA 12

In the southwestern corner of CL778379 just west of the Bennett Mine Shaft a series of pits and trenches occur. The trenches and pits are in a fracture zone within the iron-rich tholeiitic basalts. The tholeiites are intruded by syenite porphyry dikes and host north-south and east-west trending shallow dipping and vertical quartz zones.

Several moderately to strongly anomalous gold values up to 1830 ppb. occur here within vuggy quartz veins with strongly pyritic vein selvages.

The main roughly north-south trending trench extends for 100 metres and is following a vein striking 170° - 180° and dipping 036° W. The vein is approximately 50 cm. in width here and has an intensely pyritized selvage. The pyritized vein selvage consists of 5-8% disseminated pyrite and returns gold values of up to 1750 ppb. The vein itself returns 30 ppb. gold. Several pits occur north of this main trench in quartz veins within iron-rich tholeiites and associated syenite porphyries. Moderately anomalous gold values up to 600 ppb. occur here. Approximately 250 metres northeast from here, a 15 cm. wide vuggy quartz vein

with trace pyrite and a strongly epidotized, chloritic vein selvage returns 1830 ppb. gold.

Approximately 40-80 metres west of the main trench a 2-4 metre wide vein occurs striking 185° and dips vertically. Several small trenches occur within the iron-rich tholeiite cut by quartz stringers. Weakly anomalous gold values occur within this area.

The Bennett Mine shaft occurs approximately 200 metres east of the main trench zone. A two-compartment shaft was sunk to a depth of 530 feet and levels were cut at 125 foot intervals. Cross cutting and drifting were reported early in 1927. Approximately 2000 feet of diamond drilling was done in 1958 just west of the shaft. The holes intersected diorite and volcanic rocks cut by syenite dikes and quartz veins (O.D.M.N.A. files, Kirkland Lake).

About 150 metres and 250 metres north of the main shaft two quartz veins striking possibly northeast occur within the iron-rich tholeiite. The veins returned 700 ppb. and 1830 ppb. gold respectively. The veins vary from 15 cm. to 30 cm. in width and contain pyrite and molybdenite mineralization.

It appears that this area may be a 400 metre x 400 metre wide quartz stock work zone probably related to the east west and east-northeast striking porphyritic syenite dikes cutting the volcanic rocks.

Several moderately to strongly anomalous molybdenum values up to 140 ppm. occur within both the veins and the iron tholeiitic wallrock material. The values are generally higher in the vein selvage material than they are in the veins. One sample of the fractured iron tholeiite returned an arsenic enrichment of up to 11.9 ppm. with coincident anomalous molybdenum values of up to 95 ppm.

One vein sample returned 1200 ppm copper. Two of the moderately anomalous gold samples have coincident tungsten enrichments up to 16 ppm.

AREA 13

In the eastern portion of the property along the north-south access road heading off of the Highway 570, a series of A.E.M. anomalies occur in claims 778372 and 800349. To the north of here, approximately two hundred metres, several two and three channel EM anomalies occur that were previously drilled by Noranda, 1984. They encountered cherty sulphide facies iron formations with graphitic seams. From the regional mapping in Maisonville Township by H. Lovel O.G.S., a major north-south trending fault extends south from Wolf Lake along these EM anomalies.

AREA 14

In the northeastern portion of the property near TL4800N at 2880E, extending southward through claims 802333 and 802336, a 5 metre wide cherty sulphide facies iron formation occurs. The chert is striking 170° in the north and changes to 185° in the south. Unfortunately no grab samples returned anomalous gold values.

In detail, the chert grades into a cherty argillite in places and contains from 1% to 3% disseminated and stratiform-type pyrite with minor amounts of magnetite and epidote. The maximum strike length of this exhalative horizon is 800 metres.

Moderately anomalous copper and zinc values of up to 500 ppm. and 400 ppm. respectively occur within the southern extension of the chert-sulphide horizon. Strongly anomalous molybdenum values of up to 78 ppm. also occur within this horizon.

AREA 15

In the north-central section of the property, just east of the exhalative chert horizon discussed in AREA 14, a roughly north-south trending shear zone occurs within iron-rich tholeiitic basalts cut by syenite dikes and quartz stockwork

veins. Several gold occurrences in the area are associated with quartz veins hosted in iron-rich tholeiites with associated syenite porphyries. The Bennett Mine, for example, occurs within this setting.

Weakly anomalous gold values from 40 to 70 ppb. occur here from narrow quartz veins with up to 3% pyrite. The main vein system is striking 170°-175° and 90°-110° and dips 50°-70° east and 45°-55° north, respectively.

Several pits and trenches occur on narrow quartz veins within strongly pyritized vein selvages within strongly pyritized vein selvages within the iron-rich tholeiitic basalts. An intense epidotization and carbonatization with up to 8% cubic pyrite accompanies the vein system. The veins are generally grey, coarsely crystalline with epidote-chlorite inclusions and trace-1/2% disseminated pyrite, chalcopyrite and molybdenite.

Several roughly southeast trending porphyritic syenite dikes cut the iron-rich tholeiites. The syenites generally are cut by intense ladder veining and have a red hematitic alteration associated with them.

Several samples of quartz vein and vein selvage material returns moderately anomalous in copper, up to 1400 ppm. Disseminated chalcopyrite was observed in these samples. Weakly anomalous zinc values (up to 400 ppm.) are associated with these copper values.

Several moderately anomalous molybdenum values up to 95 ppm. occur within strongly epidotized, chloritized wallrock of the southeast trending quartz veins. One weakly anomalous gold value up to 65 ppb. occurs with an anomalous molybdenum sample of up to 44 ppm. and a moderately anomalous copper sample of 1400 ppm. The sample is of a 10 cm. wide rusty quartz vein with 3% pale cubic and disseminated pyrite. The vein is striking 165° and dipping 70°E. One sample of intensely epidotized wallrock of a southeast striking vein returns 59 ppm. Mo.

AREA 16

A north south trending shear zone occurs approximately 200 metres west of the No. 1 of 803559 along the iron-rich tholeiitic basalts and the cherty oxide facies iron formation contact. Strongly anomalous gold values occur within this shear zone up to .96 oz/ton. Several quartz stringers occur subparallel to the main shear zone with strongly pyritic vein selvages.

Approximately 200 metres to the north of this shear a trench occurs in weakly sheared iron-tholeiite cut by syenite dikes.

AREA 17

Several pits and trenches occur in iron-rich tholeiites in the top northeastern corner of the property along the southern claim line of 803557. The iron-tholeiites are cut by an

extensive quartz stockwork system. Several syenite dikes, trending roughly northeast cut the volcanics here. The trenches in the area are trending in several directions and most are filled with overburden cover.

Several samples of flyrock return gold values up to 360 ppb. from a vein selvage, of strongly chloritic basalt with 10-15% cubic pyrite. The quartz veins with 10% pyritic wallrock inclusions return weakly anomalous gold values of up to 140 ppb. and strongly anomalous molybdenum of up to 73 ppm. The intensely chloritized wallrock with 15% pyrite also returns 14 ppm. tungsten as well as 360 ppb. gold.

In detail, the shear zone strikes 175° and dips steeply to the east. The shearing has produced a strongly foliated iron-rich basalt with chlorite-carbonate slips that is injected by narrow 1-4 cm. wide grey quartz-epidote veins and a fractured, cherty oxide facies iron formation that is cut by an erratic display of quartz veining. According to the preliminary map of Maisonville township several holes were drilled approximately 150 metres north of this shear zone on a sulphide prospect in 1956, however due to overburden cover the old hole locations are hidden. Several trenches, however, were found within weakly sheared iron-rich tholeiites with trace pyrite and are cut by syenite dikes with associated quartz stringers.

CONCLUSIONS

1. Several disseminated sulphide zones both with quartz stockwork settings and with stratabound exhalative chert zones occur on the property in Maisonville and Grenfell townships.
2. Similar host rocks, including iron tholeiites and associated porphyritic syenite dikes occur on the Glen Auden Resources Limited property that exhibit similar geological settings to some gold deposits in the Kirkland Lake, Matachewan, Matheson and Timmins area.
3. Specifically, the Glen Auden Resources Limited property has syenite porphyry intrusives with associated gold-molybdenite mineralization similar to the Kirkland Lake area.
4. Tensional vein zones trending east-west occur near major north-south trending structures.
5. Both the north-south trending shallow dipping and vertical quartz zones and the east-west steeply dipping quartz zones carry anomalous gold values.
6. The contacts of the porphyritic syenite dikes and intrusions are important in localizing shearing and in forming tensional vein structures and stockwork systems.
7. Several exhalative chert horizons occur on the property including both sulphide facies and oxide facies iron formation. The sulphide facies occur in greater abundance.
8. The Maisonville-Grenfell Township property is totally underlain by Archean mafic volcanics (tholeiitic basalts) that are part of the Kinojevis Group (Jensen, L.S. (1978)).
9. The Kinojevis Group of rocks is similar to the Tisdale Group in Timmins which contains numerous well known gold deposits.
10. Gold mineralization frequently occurs around or within the syenite bodies as well as along north-south fault structures.
11. Several north-south trending fault structures occur on the Maisonville-Grenfell Township property, many of which can

Maisonville-Grenfell Township property, many of which can be identified on aerial photographs, regional gravity data Middleton, R. as well as on the aeromagnetic data, O.G.S. (1979).

12. Molybdenite mineralization seems to be an important component to gold mineralization.
13. Several interflow intermediate to felsic crystal tuff horizons occur on the property which may represent a slight overlap of the Blake River Volcanism, calc-alkalic event.
14. Arsenic and Tungsten values seem to be important indicators of gold enrichment in veins and selvages.

RECOMMENDATIONS

The Maisonville and Grenfell Townships property contains a number of gold showings which have been only partly exposed by old hand trenches, most of which are hidden by overburden, as well as a small zone (Bennett) which has been in part, mined. A number of the showings have geological settings which suggest potential for large tonnage gold deposits. These targets could be trenched and perhaps drilled at an early stage; however, detailed mapping and induced polarization surveys should be carried out to trace the extent and direction of these zones.

Disseminated sulphides are associated both with quartz-stockwork settings and stratabound exhalative chert zones which can be traced using induced polarization methods. Tensional vein zones trending roughly east-west occur near major north-south trending structures and therefore two line directions may be required since the majority of the rocks trend

northwesterly to north-south. Trenching of a number of showings on the property using a bulldozer and backhoe should be done as well to follow up and expose any shallow I.P. anomalies. Diamond drilling of deeper I.P. anomalies and known gold showings should be done as follow-up to the trenching.

The I.P. surveys should be conducted with an "a" spacing of 12.5 metres with n=1,2,3 and 4, on 50 metre grid line intervals on north-south, east-west or both line directions.

Area 1 An I.P. survey, on 50 metre east-west lines over claims 798874, 798875 and 798878, with trenching of near surface I.P. anomalies and in the vicinity of the old pits and trenches.

Area 2 Trenching in the vicinity of the old trenches and pits.

Area 4 and Area 12 An I.P. survey, on 50 metre east-west lines is recommended over claims 798864, 801217, 778379 and the eastern half of claim 798863 with trenching of any near surface anomalies.

Area 5 An I.P. survey for tracing the extent of the zone on 50-metre east-west and north-south lines over claims 800346 and 800348 to find the extent of the north-south fault zone and the east-west auriferous tensional veins. Trenching is also recommended in the vicinity of the trenches.

Area 7 An I.P. survey covering the eastern half of claim 802748 and all of claim 801876 extending north into claims 802747 and 801220 with north-south and east-west lines spaced 50 metres apart with trenching of any near surface I.P. anomalies.

Area 8 Trenching with a bulldozer and backhoe is recommended in the vicinity of the trenches to better expose the overburden covered outcrops in the southeastern corner of claim 802342.

Area 9 Bulldozer work is recommended in the vicinity of the

old treches and pits, in order to establish the true vein directions.

- Area 11 Trenching in the vicinity of the old pits and trenches followed by I.P. survey over claim 802343 and claim 802349 on 50 metre east-west lines.
- Area 13 An I.P. survey is recommended over claims 800349, 800347, 7783772, 778371, 800345 and 778368 to find the extension of the pyritic graphite horizon that shows up on A.E.M. anomalies.
- Area 16 An I.P. survey is recommended over claim 803558 and 803559 on east-west lines spaced 50 metres apart to find the north and south strike extension of the shear zone that returned .96 oz./ton Au.
- Area 17 Bulldozer and backhoe work is recommended in the vicinity of the pits and trenches in order to establish true vein directions followed by an I.P. survey over claims 803557 and 803560 on a specific grid system at 50 metre line spacings.

BUDGET

Phase I

Line-cutting at \$300/claim on 23 claims \$ 6,900

Sub Total \$ 6,900

Phase II

I.P. Surveys on specific areas 55 miles at \$1,300/mi. \$ 71,500
Trenching 40 days at \$500/day 20,000
Sampling and assaying 7,000

Sub Total \$ 98,500

Phase III

Diamond drilling of 2 known showings with a series of short holes = 2800 ft. at \$25/ft. 70,000
Assaying and Supervison 10,000

Sub Total \$ 80,000

Total of Phase I, II, and III \$185,400
Contingency at 10% 18,540

TOTAL \$203,940

Further drilling and trenching to be contingent on results of Phases I, II and III.

Respectfully Submitted,

Nadia Cairra

Nadia Cairra, B.Sc.
October 2, 1985

RECEIVED

NOV 19 1985

MINING LANDS SECTION

REFERENCES

- Fyon, Andy J. and
Crocket, J.H. Gold Exploration in the Timmins Area Using
Field and Lithochemical Characteristics
of Carbonate Alteration Zones O.G.S. Study
26.
- Grunsky, E.C. No. 58 Abitibi Alteration Study
- Jensen, L.S. Geology of Thackeray, Elliot, Tannahill,
and Dokis Townships, O.G.S. Report 165
- Jensen, L.S. and Langford
F.F. 1983 Geology and Petrogenesis of the Archean
Abitibi Belt in the Kirkland Lake Area,
Ontario; O.G.S. Open file Report 5455, p520
- Jensen, L.S. No. 10 Regional Stratigraphy and Structure
of the Timmins-Kirkland Lake Area, District
of Cochrane and Timiskaming and the Kirkland
Lake - Larder Lake Area
- Kerrich, R. Geochemistry of Gold Deposits in the Abitibi
Greenstone Belt, Special Volume 27
The Canadian Institute of Mining and
Metallurgy
- Lovell, H. L., Geology of the Bourkes Area, O.G.S., Report
92 1971
- Lovell, H. L., Geology of the Matachewan Area 1967
- Pyke, D. R. Geology of the Timmins Area, O.G.S. Report
p. 219
- Tagliamonte, Frank P. Report on the Property of Glen Auden
Resources Limited and Adola Mining Corp.
Maisonville and Grenfell Townships.

CERTIFICATION

I, Nadia M. Cairra, B.Sc., of Timmins, Ontario, certify that:

1. I am a graduate of the University of British Columbia, Vancouver, B.C., with a B.Sc. degree in Geology obtained in 1981.
2. I have been practising my profession in Canada since 1981.
3. I have no direct or indirect interest in the properties, leases or securities of Glen Auden Resources Ltd., Maisonville-Grenfell Township property, nor do I expect to receive any.

Dated this October 2, 1985 Timmins, Ontario.

Nadia M. Cairra, B.Sc.

A P P E N D I X

R. S. MIDDLETON EXPLORATION SERVICES

(N. CAIRA)

PROJ: M-78

WO NO: 85-0520

PAGE: 1

SAMPLE ID	CU PPM	ZN PPM	PB PPM	AG PPM	AU PPR	AU-FA OZ/TON	MO PPM	AS PPM	SR PPM	W PPM
067851	89	39	3	<.2	<5	---	7	.7	<.2	<4
067852	2100	41	9	.3	20	---	63	<.2	<.2	5
067853	650	51	5	<.2	45	---	22	.4	<.2	<4
067854	490	20	4	<.2	45	---	31	1.7	<.2	28
067855	79	18	6	.4	135	---	83	1.0	.4	12
067856	32	9	13	<.2	15	---	25	.5	<.2	4
067857	88	61	5	.7	35	---	6	2.8	<.2	8
067858	440	33	<1	<.2	10	---	<1	7.0	<.2	6
067859	240	53	<1	<.2	45	---	<1	.5	<.2	6
067860	21	7	<1	<.2	60	---	4	.9	<.2	<4
067861	19	18	12	<.2	45	---	140	.7	<.2	10
067862	180	13	21	5.5	11000	.320	6	5.3	<.2	<4
067863	44	28	6	<.2	45	---	6	1.0	<.2	<4
067864	21	20	300	4.9	140	---	73	.9	<.2	4
067865	42	55	230	2.9	360	---	23	.3	<.2	14
067866	200	19	260	<.2	<5	---	7	.7	<.2	<4
067867	2200	72	5	<.2	5	---	7	.5	<.2	<4
067868	450	25	<1	<.2	5	---	1	.8	<.2	<4
067869	20	13	<1	<.2	<5	---	4	1.2	<.2	<4
067870	15	15	<1	<.2	35	---	4	.9	.2	<4
067871	260	36	<1	<.2	15	---	64	1.2	<.2	<4
067872	62	110	102	<.2	5	---	13	2.1	.2	12
067873	46	24	<1	<.2	10	---	43	2.5	.2	<4
067874	12	7	<1	<.2	115	---	8	.4	.2	<4
067875	19	9	<1	.4	700	---	47	.8	.2	<4
067876	130	33	<1	<.2	5	---	10	1.9	<.2	<4
067877	64	54	<1	<.2	35	---	40	1.0	<.2	<4
067878	110	85	23	<.2	50	---	95	11.9	.4	<4
067879	19	14	3	.8	1500	.030	27	.7	<.2	<4
067880	73	62	<1	.6	600	---	8	1.1	<.2	20
067881	69	49	<1	<.2	5	---	9	.9	<.2	<4
067882	1000	540	8	<.2	5	---	13	19.0	.2	1
067883	1300	42	.1	<.2	62	---	10	.5	<.2	4
067884	82	43	1300	1.1	<5	---	3	4.0	<.2	<4
067885	19	10	8	<.2	50	---	9	.7	.2	<4
067886	11	6	3	<.2	<5	---	4	.4	<.2	<4
067887	21	20	7	1.3	1750	.030	140	1.3	.2	16
067888	21	43	<1	<.2	35	---	15	.7	.2	4
067889	38	25	3	1.3	1500	.040	28	1.2	<.2	6
067890	29	4	4	.6	1820	.050	15	.5	.2	<4
067891	13	10	<1	<.2	10	---	19	<.2	.2	40
067892	32	50	<1	<.2	<5	---	2	<.2	<.2	<4
067893	38	5	8	<.2	<5	---	21	.7	<.2	<4
067894	110	9	27	<.2	20	---	59	.9	<.2	<4
067895	69	48	<1	<.2	<5	---	4	.3	<.2	<4

304 CARLINGVIEW DRIVE
REXDALE, ONTARIO
M9W 5G2

(416) 675-3870

3750 - 19TH STREET
SUITE 105
CALGARY, ALBERTA
T2E 6V2
(403) 276-8701

FILE: 15_0520
DATE: 10/09/85
MATRIX: AQ HFG

BARRINGER MAGENTA

WO NO: 85-0520

PAGE: 2

K. S. MIDDLETON EXPLORATION SERVICES (N. CATRA) PROJ: N-78

SAMPLE ID	CU PPM	ZN PPM	PB PPM	AG PPM	AU PPM	AU-FA OZ/TON	MO PPM	AS PPM	SB PPM	W PPM
067096	150	54	2	.3	235	---	3	1.8	<.2	3
067097	150	17	17	<.2	210	---	48	.6	<.2	<4
067098	120	55	<1	<.2	35	---	3	.5	<.2	16
067099	71	56	<1	<.2	45	---	3	<.2	<.2	14
067100	34	10	2	<.2	5	---	5	.5	<.2	15
18280	290	300	35	<.2	10	---	7	5.6	<.2	<4
18281	200	58	5	<.2	10	---	9	2.1	<.2	<4
18282	84	49	<1	<.2	25	---	12	.6	<.2	14
18283	85	5	<1	<.2	25	---	14	.4	.2	30
18284	21	7	11	1.4	4750	.900	49	2.5	.2	12
18285	72	34	<1	<.2	350	---	23	<.2	<.2	<4
18286	100	73	<1	<.2	15	---	1	<.2	<.2	<4
18287	1400	11	170	1.3	65	---	44	.6	<.2	<4
18288	570	39	160	.4	40	---	17	.5	<.2	<4
18289	46	5	85	.6	10	---	13	.2	<.2	<4
18290	85	6	44	.5	30	---	56	.4	<.2	<4
18291	110	47	6	<.2	<5	---	5	1.3	<.2	<4
18292	110	19	51	<.2	15	---	95	.4	<.2	<4
18293	6	42	<1	<.2	25	---	2	<.2	<.2	<4
18294	560	400	<1	<.2	45	---	78	2.6	.2	<4
18295	32	8	22	<.2	10	---	6	.4	<.2	<4
18296	40	8	47	<.2	5	---	100	.4	<.2	<4
18297	15	8	36	<.2	270	---	23	.4	.2	<4
18298	9	5	5	<.2	5	---	120	.4	<.2	<4
18299	310	590	3	<.2	85	---	16	4.5	.2	14
18300	160	110	<1	<.2	40	---	30	4.4	<.2	12

R. S. MIDDLETON EXPLORATION SERVICES

(H. CATRA)

PROJ: M-78

WQ NO: 85-0583

PAGE: 1

SAMPLE ID	CU PPM	ZN PPM	FE PPM	AG PPM	AU PPM	MO PPM	AS PPM	SB PPM
67967	280	11	2	<.2		2	<.2	<.2
67968	12	9	<1	<.2	<.2		<.2	<.2
67969	240	1100	11	<.2	<.5	50	21.5	
67970	560	50	<1	<.2	<.5	<1	.2	<.2
67971	59	52	3	<.2	<.5	7	10.5	.2
67972	210	700	19	.4	10	45	35.1	.9
67973	280	900	15	.3	<.5	100	65.2	.7
67974	170	270	10	<.2	<.5	13	21.5	1.3
67975	540	1500	18	.5	47.5		38.3	.9
67976	98	75	<1	<.2	10		1.4	<.2
67977	55	90	4	<.2	110	7	2.4	
67978	18	37	7	<.2	300	2	<.2	<.2
67979	140	42	12	.6	<.5	10	126	.6
67980	340	50		<.2	<.5	4	.9	<.2
67981	800	190	1000	1.4	20	75	.7	<.2
67982	74	51	12	<.2	10	9	.9	<.2
67983	78	30	<1	<.2	20	7	.9	<.2
67984	39	47	<1	<.2	<.5	1	<.2	<.2
67985	29	15	<1	<.2	<.5	2	<.2	<.2
67986		6	2	<.2	365	80	.5	<.2
67987	10		1	<.2	10	150	.3	<.2
67988	120	20		<.2	115	120	1.0	<.2
67989	42	34	<1		100	8	2.4	<.2
67990	260	38	<1	.3		17	2.0	<.2

CERTIFICATION

I, Nadia M. Caira, B.Sc., of Timmins, Ontario, certify that:


1. I am a graduate of the University of British Columbia, Vancouver, B.C., with a B.Sc. degree in Geology obtained in 1981.
2. I have been practising my profession in Canada since 1981.
3. I have no direct or indirect interest in the properties, leases or securities of Glen Auden Resources Ltd., Maisonville-Grenfell Township property, nor do I expect to receive any.

Dated this October 2, 1985 Timmins, Ontario.

Nadia Caira
Nadia M. Caira, B.Sc.


RECEIVED
NOV 19 1985
MINING LANDS SECTION

10



52 / 467

TL1576N



	Au(ppb)	Mo(ppm)
18284	.96oz/ton	44
18285	650	23
67986	365	80
67987	10	150

REVISIONS

ROBERT S. MIDDLETON
EXPLORATION SERVICES INC.

for GLEN AUDEN RESOURCES LIMITED

Title

Detailed Geology
AREA 5

Fig. 12

Date: Oct 1985

Scale: 1:100

N.T.S.:

Drawn: NC

Approved:

File: M-78

Claim List for 26 Claims

Claim Number

L 798 873 -
L 798 874
L 798 875
L 798 876
L 798 877
L 798 878

L 800 345 ✓
L 800 346 ✓

L 801 219 ✓
L 801 220 ✓
L 801 221 ✓
L 801 222 ✓

L 802 331 ✓

L 802 341 ✓
L 802 342 ✓
L 802 343 ✓

L 802 346 ✓
L 802 347 ✓
L 802 348 ✓
L 802 349 ✓

L 802 366 ✓
L 802 367 ✓
L 802 368 ✓
L 802 369 ✓

L 802 748 ✓
L 802 750 ✓



The Mining Act

In the matter of mining claims:

L 798873, et al,
in Maisonville Township
as listed on Report
of Work #315.

On consideration of an application from the recorded holder, Glen Auden Resources Limited
under Section 77 Subsection 22 of The Mining Act, I hereby order that the time for filing reports and plans in support of
Geological assessment work recorded on September 3, 19 85
be extended until and including November 15, 19 85.

1985.11.19

Date

[Signature]

Signature of Director, Land Management Branch

Copies: AR
Glen Auden Resources Limited
136 Cedar Street South
P.O. Box 1637
Timmins, Ontario
P4N 7W8

R.S. Middleton Exploration Services Inc
P.O. Box 1637
Timmins, Ontario
P4N 7W8
Attention: Nadia Cairn

Mining Recorder
Timmins, Ontario

7

705-264-4246.

Marylin.

called. ✓
Oct. 25

Rec. on file
18 Oct.
in Kirkland
Lake
2072/85
M.P. 10/25/85

REGISTERED

October 23, 1985

Report Of Work #315

Glen Auden Resources Limited
136 Cedar Street South
P.O. Box 1637
— Timmins, Ontario
P4N 7W8

Dear Sirs:

RE: Mining Claims L 798873, et al,
in Maisonville Township

I have not received the reports and maps (in duplicate)
for the Geological Survey on the above-mentioned claims.

As the assessment "Report of Work" was recorded by the
Mining Recorder on September 3, 1985 the 60 day period
allowed by Section 77 of the Mining Act for the submission
of the technical reports and maps to this office will
expire on November 2, 1985.

If the material is not submitted to this office by November 2, →
1985 I will have no alternative but to instruct the Mining
Recorder to delete the work credits from the claim record
sheets.

For further information, please contact Mr. Arthur Barr
at (416)965-4888.

Yours sincerely,

S.E. Yundt
Director
Land Management Branch

Whitney Block, Room 6643
Queen's Park
Toronto, Ontario
M7A 1W3
Phone: (416)965-4888

AB/mc

cc: R.S. Middleton Exploration Services Inc Mining Recorder
P.O. Box 1637 Timmins, Ontario
Timmins, Ontario P4N 7W8
Attention: Nadia Cairra

Enc).

Mining Lands Section

File No 8629

Control Sheet

TYPE OF SURVEY GEOPHYSICAL
 GEOLOGICAL
 GEOCHEMICAL
 EXPENDITURE

MINING LANDS COMMENTS:

Lg.
L.S.

J. Hurst
Signature of Assessor

Nov 20/85.
Date

2.8629

198873 ✓

74 ✓

75 ✓

76 ✓

77 ✓

78 ✓

800345 ✓

46 ✓

801219 ✓

20 ✓

21 ✓

22 ✓

802331 ✓

802341 ✓

42 ✓

43 ✓

346 ✓

47 ✓

48 ✓

49 ✓

366 ✓


67 ✓

68 ✓

69 ✓

802748 ✓

50 ✓



LEGEND

BRANDAMERIAN
 HUMBOLDTIAN (Cobalt Group)
 7a Conglomerate
 7b Arkose
 7c Quartzite, greywacke

ARCHAIC
 MAFIC INTRUSIVE ROCKS
 6 Squigranular Diabase
 intrusive contact

FELSIC INTRUSIVE ROCKS
 5b Porphyritic Syenite
 5c Syenodiorite
 5d Mafic Syenite
 intrusive contact


METASEDIMENTS
 3a Cherty sulphide iron formation
 3b Slate, argillite
 3c Laminated, siliceous
 3d Cherty oxide iron formation
 3e Graphite
 3f Cherty tuff

MAFIC METAVOLCANICS (Kinojevis Group)
 2 Iron-rich tholeiitic basalts
 1 Iron-poor tholeiitic basalts
 a Massive, fine-grained
 b Pillowed
 c Pillow breccia
 d Massive (2-5mm) coarsegrained
 e Hyaloclastic, fragmental
 f Amygdaloidal
 g Crystal tuff (felsic)—overlap of Blake River Group
 h Variolitic
 i Chlorite schist

carb carbonatization
 sil silicification
 ep epidotization
 hem hematization
 gt garnet
 mt magnetite
 py pyrite
 cp chalcocopyrite
 gm galena
 mo molybdenite
 gr graphite
 q.v quartz vein

str strong
 tr trace
 wk weak

vein inclined, vertical
 bedding inclined, vertical
 pillow lava top (arrow)
 foliation
 fault
 shear
 geological contact
 outcrop
 fracture inclined, vertical
 shaft
 pit
 trench
 rock sample location
 quartz stringer zone
 swamp
 claim post located
 claim post unlocated
 AEM anomaly
 Ground magnetometer anomaly
 1500/5.5 Au(ppb) Ar(ppm)
 * map matching points




REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for GLEN AUDEN RESOURCES LIMITED		
	Title PROPERTY GEOLOGY		
	Sheet I		
	Date: OCT. 1985	Scale: 1:4000	N.T.S.:
	Drawn: C.G	Approved:	File: M-78




28629

REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for GLEN AUDEN RESOURCES LIMITED		
	Title PROPERTY GEOLOGY Sheet 2		
	Date: OCT. 1985	Scale: 1:4000	N.T.S.:
	Drawn: C.G.	Approved:	File: M-78

NOTE: Refer to Sheet 1 for legend





CONCESSION III
CONCESSION II

T.L. 3192 N

L 3000 N

L 2500 N

L 2000 N

CONCESSION II
CONCESSION I

SESEKINIKA
LAKE

T.L. 1576 N

L 1500 N

L 1000 N


LOT 10

L 1600 E

LOT 9

L 2450 E


LOT 8




NOTE: Refer to Sheet 1 for legend

REVISIONS	ROBERT S. MIDDLETON EXPLORATION SERVICES INC.		
	for GLEN AUDEN RESOURCES LIMITED		
	Title		
	PROPERTY GEOLOGY Sheet 3		
	Date: OCT 1985	Scale: 1:4000	N.T.S.:
	Drawn: C.G.	Approved:	File: M-78

28629



Scale, 1 inch to 50 miles
N.T.S. reference 42/A1, 42/A8



SYMBOLS

- Glacial striae.
- Small bedrock outcrop.
- Area of bedrock outcrop.
- Bedding, top unknown; (inclined, vertical).
- Bedding, top indicated by arrow; (inclined, vertical, overturned).
- Lava flow; top (arrow) from pillows shape and packing.
- Schistosity; (horizontal, inclined, vertical).
- Geological boundary, observed.
- Geological boundary, position interpreted.
- Fault; (observed, assumed). Arrows indicate horizontal movement.
- Jointing; (horizontal, inclined, vertical).
- Drag folds with plunge.
- Anticline, syncline, with plunge.
- Drill hole; (vertical, inclined).
- Vein. Width in inches.
- Shaft; depth in feet.
- Magnetic attraction.
- Motor road. Provincial highway number encircled where applicable.
- Other road.
- Trail, postage, winter road.
- Building.
- District boundary, approximate position only.
- Township boundary, approximate position only.
- Lot and Concession lines, approximate position only.
- Property boundary, approximate position only.
- Claim line, surveyed, approximate position only.
- Location of mining property, surveyed. (See "List of Properties")
- Location of mining property, unsurveyed. (See "List of Properties")

LEGEND

- CENOZOIC***
- RECENT**
Swamp and stream deposits.
- PLEISTOCENE**
Sand, gravel, clay.
- UNCONFORMITY
- PRECAMBRIAN^b**
- PROTEROZOIC**
- HURONIAN**
- COBALT GROUP**
- GOWGANDA FORMATION**
- 75 Conglomerate.
- 76 Archaic.
- 77 Quartzite and greywacke.
- 78 Argillite.
- UNCONFORMITY
- ARCHEAN**
- MAFIC INTRUSIVE ROCKS (MATACHEWAN)**
- 4 Diabase.
- INTRUSIVE CONTACT
- FELSIC INTRUSIVE ROCKS (ALGOMAN)**
- 5a Granite, hornblende-biotite granite, granodiorite.
- 5b Granite gneiss.
- 5c Syenite, syenite porphyry, quartz syenite.
- 5d Mafic syenite, lamprophyre.
- INTRUSIVE CONTACT
- MAFIC AND ULTRAMAFIC INTRUSIVE ROCKS (HAILEYBURIAN, EARLY ALGOMAN, AND SOME KEEWATIN)**
- 4a Serpentine, peridotite.
- 4b Gabbro, diorite, may include some ta.
- INTRUSIVE CONTACT
- METASEDIMENTS, TIGHTLY FOLDED (TIMISKAMING OR KEEWATIN)**
- 3 Slate, chert, tuff.
- UNCONFORMITY
- FELSIC METAVOLCANICS (KEEWATIN)**
- 2a Dacite, rhyolite, minor amounts of rhyolite.
- 2b Siliceous and intermediate tuff and agglomerate.
- 2c Dacite porphyry, amygdaloidal dacite, spherulitic dacite.
- MAFIC TO INTERMEDIATE METAVOLCANICS (KEEWATIN)**
- 1a Basalt and andesite (pillowed and massive) may include some 4b.
- 1b Mafic agglomerate and tuff.
- 1c Altered (bleached) mafic metavolcanics.
- 1d Andesite porphyry, amygdaloidal andesite, variscite andesite.

*Unconsolidated deposits. Cenozoic deposits are represented by the higher coloured parts on the map.
^bBedrock geology. Outcrops and inferred extensions of each rock map unit are shown. Where in places a formation is too narrow to show colour and must be represented in black, a short black bar appears in the appropriate block.
 †Appears on companion sheets Map 2213 or Map 2214.

- Ag Silver.
- asb Asbestos.
- Au Gold.
- carb Carbonate.
- Cu Copper.
- ep Epidote.
- gf Graphite.
- gn Garnet.
- hem Hematite.
- mag Magnetite.
- Mo Molybdenum.
- Ni Nickel.
- Pb Lead.
- q Quartz.
- qt Quartz-carbonate.
- s Sulphide mineralization.
- shw Scheelite.
- Te Tellurium.
- Zn Zinc.

SOURCES OF INFORMATION
 Geology by H. L. Lovell and assistants, 1965 and 1966. Geology is not tied to surveyed lines.
 Geological and geophysical maps and reports of mining companies.
 Preliminary maps P. 329 Benoit Township issued 1966 and P. 429 Maisonville Township issued 1967, scale 1 inch to 1/4 mile.
 Cartography by D. V. Impey and assistants, Ontario Department of Mines, 1970.
 Basemaps derived from maps of the Forest Resources Inventory, Ontario Department of Lands and Forests, with revisions by H. L. Lovell.
 Magnetic declination in the area was approximately 9°30' W, 1966.

- LIST OF PROPERTIES**
- BENOIT TOWNSHIP**
18. Bergstrand [circa 1921]
 19. Bunting (Johnson) [circa 1920]
 20. Carlson, G. [circa 1942]
 21. Chamandy, K. [circa 1948], and Mokta Canada Limited (1968)
 22. Collier [circa 1921]
 23. David Mines Ltd.
 24. Erickson, K. J.
 25. Hagen, Ole
 26. Keefe, Daniel G. (W. half of Skjonsby-Klanderland)
 27. Noranda Exploration Company
 28. Potter, J. S. (Wickstead-Mahor) [circa 1920]
 29. Scovold-Nelson [circa 1921]
 30. Skjonsby, K. E.
 31. Sward, Axel (part of the former Webb claims, circa 1921)
 32. Thompson-McLeod [circa 1921]
- MAISONVILLE TOWNSHIP**
33. Bennett Mining Co. Ltd. [circa 1929]
 34. Cole [circa 1951]
 35. Elcira Mines Ltd. [circa 1938]
 36. Golden Summit Mining Co. Ltd., The
 37. Hastings-Moffat [circa 1924]
 38. International Nickel Co. of Canada Ltd., The
 39. Kerr Addison Mines Ltd. "Gosse Lake group"
 40. Kerr Addison Mines Ltd. "Kasapika group"
 41. Labrie-Smith [circa 1914]
 42. Langs, John G., trustee
 43. Lee, Mrs. O. M., estate
 44. Murray, John E.
 45. Pann, Sidney A.
 46. Peterson, Warren [circa 1953]
 47. Puddon, A. [circa 1953]
 48. Russell, J. G.
 49. Sesekinika Township [circa 1934]
 50. Ventures Claims Ltd.
 51. Wickstead [circa 1948]
- Date in square brackets refers to year of last major work.

Map 2215
BENOIT AND MAISONVILLE TOWNSHIPS
 TIMISKAMING DISTRICT

Scale 1:31,680 or 1 Inch to 1/2 Mile

