

THESE TERMS GOVERN YOUR USE OF THIS DOCUMENT

Your use of this Ontario Geological Survey document (the “Content”) is governed by the terms set out on this page (“Terms of Use”). By downloading this Content, you (the “User”) have accepted, and have agreed to be bound by, the Terms of Use.

Content: This Content is offered by the Province of Ontario’s *Ministry of Northern Development and Mines* (MNDM) as a public service, on an “as-is” basis. Recommendations and statements of opinion expressed in the Content are those of the author or authors and are not to be construed as statement of government policy. You are solely responsible for your use of the Content. You should not rely on the Content for legal advice nor as authoritative in your particular circumstances. Users should verify the accuracy and applicability of any Content before acting on it. MNDM does not guarantee, or make any warranty express or implied, that the Content is current, accurate, complete or reliable. MNDM is not responsible for any damage however caused, which results, directly or indirectly, from your use of the Content. MNDM assumes no legal liability or responsibility for the Content whatsoever.

Links to Other Web Sites: This Content may contain links, to Web sites that are not operated by MNDM. Linked Web sites may not be available in French. MNDM neither endorses nor assumes any responsibility for the safety, accuracy or availability of linked Web sites or the information contained on them. The linked Web sites, their operation and content are the responsibility of the person or entity for which they were created or maintained (the “Owner”). Both your use of a linked Web site, and your right to use or reproduce information or materials from a linked Web site, are subject to the terms of use governing that particular Web site. Any comments or inquiries regarding a linked Web site must be directed to its Owner.

Copyright: Canadian and international intellectual property laws protect the Content. Unless otherwise indicated, copyright is held by the Queen’s Printer for Ontario.

It is recommended that reference to the Content be made in the following form: <Author’s last name>, <Initials> <year of publication>. <Content title>; Ontario Geological Survey, <Content publication series and number>, <total number of pages>p.

Use and Reproduction of Content: The Content may be used and reproduced only in accordance with applicable intellectual property laws. *Non-commercial* use of unsubstantial excerpts of the Content is permitted provided that appropriate credit is given and Crown copyright is acknowledged. Any substantial reproduction of the Content or any *commercial* use of all or part of the Content is prohibited without the prior written permission of MNDM. Substantial reproduction includes the reproduction of any illustration or figure, such as, but not limited to graphs, charts and maps. Commercial use includes commercial distribution of the Content, the reproduction of multiple copies of the Content for any purpose whether or not commercial, use of the Content in commercial publications, and the creation of value-added products using the Content.

Contact:

FOR FURTHER INFORMATION ON	PLEASE CONTACT:	BY TELEPHONE:	BY E-MAIL:
The Reproduction of Content	MNDM Publication Services	Local: (705) 670-5691 Toll Free: 1-888-415-9845, ext. 5691 (inside Canada, United States)	Pubsales@ndm.gov.on.ca
The Purchase of MNDM Publications	MNDM Publication Sales	Local: (705) 670-5691 Toll Free: 1-888-415-9845, ext. 5691 (inside Canada, United States)	Pubsales@ndm.gov.on.ca
Crown Copyright	Queen’s Printer	Local: (416) 326-2678 Toll Free: 1-800-668-9938 (inside Canada, United States)	Copyright@gov.on.ca

LES CONDITIONS CI-DESSOUS RÉGISSENT L'UTILISATION DU PRÉSENT DOCUMENT.

Votre utilisation de ce document de la Commission géologique de l'Ontario (le « contenu ») est régie par les conditions décrites sur cette page (« conditions d'utilisation »). En téléchargeant ce contenu, vous (l'« utilisateur ») signifiez que vous avez accepté d'être lié par les présentes conditions d'utilisation.

Contenu : Ce contenu est offert en l'état comme service public par le *ministère du Développement du Nord et des Mines* (MDNM) de la province de l'Ontario. Les recommandations et les opinions exprimées dans le contenu sont celles de l'auteur ou des auteurs et ne doivent pas être interprétées comme des énoncés officiels de politique gouvernementale. Vous êtes entièrement responsable de l'utilisation que vous en faites. Le contenu ne constitue pas une source fiable de conseils juridiques et ne peut en aucun cas faire autorité dans votre situation particulière. Les utilisateurs sont tenus de vérifier l'exactitude et l'applicabilité de tout contenu avant de l'utiliser. Le MDNM n'offre aucune garantie expresse ou implicite relativement à la mise à jour, à l'exactitude, à l'intégralité ou à la fiabilité du contenu. Le MDNM ne peut être tenu responsable de tout dommage, quelle qu'en soit la cause, résultant directement ou indirectement de l'utilisation du contenu. Le MDNM n'assume aucune responsabilité légale de quelque nature que ce soit en ce qui a trait au contenu.

Liens vers d'autres sites Web : Ce contenu peut comporter des liens vers des sites Web qui ne sont pas exploités par le MDNM. Certains de ces sites pourraient ne pas être offerts en français. Le MDNM se dégage de toute responsabilité quant à la sûreté, à l'exactitude ou à la disponibilité des sites Web ainsi reliés ou à l'information qu'ils contiennent. La responsabilité des sites Web ainsi reliés, de leur exploitation et de leur contenu incombe à la personne ou à l'entité pour lesquelles ils ont été créés ou sont entretenus (le « propriétaire »). Votre utilisation de ces sites Web ainsi que votre droit d'utiliser ou de reproduire leur contenu sont assujettis aux conditions d'utilisation propres à chacun de ces sites. Tout commentaire ou toute question concernant l'un de ces sites doivent être adressés au propriétaire du site.

Droits d'auteur : Le contenu est protégé par les lois canadiennes et internationales sur la propriété intellectuelle. Sauf indication contraire, les droits d'auteurs appartiennent à l'Imprimeur de la Reine pour l'Ontario.

Nous recommandons de faire paraître ainsi toute référence au contenu : nom de famille de l'auteur, initiales, année de publication, titre du document, Commission géologique de l'Ontario, série et numéro de publication, nombre de pages.

Utilisation et reproduction du contenu : Le contenu ne peut être utilisé et reproduit qu'en conformité avec les lois sur la propriété intellectuelle applicables. L'utilisation de courts extraits du contenu à des fins *non commerciales* est autorisée, à condition de faire une mention de source appropriée reconnaissant les droits d'auteurs de la Couronne. Toute reproduction importante du contenu ou toute utilisation, en tout ou en partie, du contenu à des fins *commerciales* est interdite sans l'autorisation écrite préalable du MDNM. Une reproduction jugée importante comprend la reproduction de toute illustration ou figure comme les graphiques, les diagrammes, les cartes, etc. L'utilisation commerciale comprend la distribution du contenu à des fins commerciales, la reproduction de copies multiples du contenu à des fins commerciales ou non, l'utilisation du contenu dans des publications commerciales et la création de produits à valeur ajoutée à l'aide du contenu.

Renseignements :

POUR PLUS DE RENSEIGNEMENTS SUR	VEUILLEZ VOUS ADRESSER À :	PAR TÉLÉPHONE :	PAR COURRIEL :
la reproduction du contenu	Services de publication du MDNM	Local : (705) 670-5691 Numéro sans frais : 1 888 415-9845, poste 5691 (au Canada et aux États-Unis)	Pubsales@ndm.gov.on.ca
l'achat des publications du MDNM	Vente de publications du MDNM	Local : (705) 670-5691 Numéro sans frais : 1 888 415-9845, poste 5691 (au Canada et aux États-Unis)	Pubsales@ndm.gov.on.ca
les droits d'auteurs de la Couronne	Imprimeur de la Reine	Local : 416 326-2678 Numéro sans frais : 1 800 668-9938 (au Canada et aux États-Unis)	Copyright@gov.on.ca

**Uranium and Thorium
Deposits of
Northern Ontario**

by

James A. Robertson and Kerry L. Gould

**This project was partially funded by
the Ministry of Northern Affairs**

1983



Ontario

**Ministry of
Natural
Resources**

**Hon. Alan W. Pope
Minister**

**W. T. Foster
Deputy Minister**

Copyright OMNR 1983
Printed in Canada

ISSN 0706-4551
ISBN 0-7743-8439-5

Publications of the Ontario Ministry of Natural Resources are available from the following sources. Orders for publications should be accompanied by cheque or money order payable to the Treasurer of Ontario.

Reports, maps, and price lists (personal shopping or mail order):
Public Service Centre, Ministry of Natural Resources
Room 1640, Whitney Block, Queen's Park
Toronto, Ontario M7A 1W3

Reports and accompanying maps only (personal shopping):
Ontario Government Bookstore
Main Floor, 880 Bay Street
Toronto, Ontario

Reports and accompanying maps (mail order or telephone orders):
Publications Services Section, Ministry of Government Services
5th Floor, 880 Bay Street
Toronto, Ontario M7A 1N8
Telephone (local calls), 965-6015
Toll-free long distance, 1-800-268-7540
Toll-free from area code 807, 0-ZENITH-67200

Every possible effort is made to ensure the accuracy of the information contained in this report, but the Ministry of Natural Resources does not assume any liability for errors that may occur. Source references are included in the report and users may wish to verify critical information.

Parts of this publication may be quoted if credit is given. It is recommended that reference to this report be made in the following form:

Robertson, J.A., and Gould, K.L.
1983: Uranium and Thorium Deposits of Northern Ontario; Ontario
Geological Survey, Mineral Deposits Circular 25, 152p.

FOREWORD

This circular briefly summarizes data on some four hundred uranium and/or thorium deposits of northern Ontario and supercedes Mineral Resources Circular 9, published in 1968. It is a part of a group of publications providing an up-to-date data base on Ontario's uranium deposits, their classification, distribution, resources and factors affecting their development or further resources.

The most significant of the deposits described are fossil placers lying at or near the base of the Middle Precambrian Huronian Supergroup, including the producing and past-producing mines in the Elliot Lake - Agnew Lake area.

Should a market for thorium develop, the Elliot Lake uranium deposits would constitute a major source.

E.G. Pye
Director
Ontario Geological Survey

CONTENTS

Introduction	1	Kamis Occurrence	24
Acknowledgements	4	Lacnor Mine	25
Explanation of this Circular	4	Milliken Mine (Rio Algom Limited)	25
Uranium and Thorium Deposits of Algoma District	8	Nordic Mine (Rio Algom Limited)	26
Aberdeen Township	8	Silvermaque Occurrence	27
Rio Tinto Prospect	8	Stanleigh Mine (Rio Algom Limited)	27
Albanel Township	8	Jollineau Township	28
Little White River Occurrence	8	Gardiner Occurrence	28
Beange Township	9	Joubin Township	28
Candore Prospect	9	Abeta Occurrence	28
Callinan Flin-Flon Occurrence	9	Milliqua Prospect (Pardee Reef of Pardee Zone)	28
Rio Algom (Quirke Group West) Occurrence	10	Nasco Occurrence	29
Rio Algom (Span-North) Occurrence	10	Pardee Prospect (Pardee Zone)	29
Bolger Township	11	Pecors Lake (West) Prospect (Pardee Zone)	30
Gui-Por Prospect	11	Stancan Occurrence	30
Moon Lake Mines Occurrence (Moon Lake Zone)	11	St. Mary's Occurrence (Pardee Zone)	31
Nordic Group West Prospect (Moon Lake Zone)	12	Vite Occurrence (Pecors Zone)	31
Bouck Township	12	Kamichisitit Township	32
Benner Occurrence	12	Kee No.2 Occurrence	32
Denison Mine	12	Kincaid Township	32
Gemico Prospect	13	Hathaway Occurrence	32
Kerr-McGee 150-151 Occurrence	14	Kirkwood Township	33
Quirke Mine No.1 (Rio Algom Limited)	14	Abconore Occurrence	33
New Quirke Mine No.2 (Rio Algom Limited)	15	Labelle Township	33
Spanish American Mine (Rio Algom Limited)	16	Franz Occurrence	33
Buckles Township	16	Ottawa Associates Occurrence	34
Can-Met Mine (Denison Mines Limited)	16	Labonte Township	34
Conecho Prospect	17	McCombe Syndicate Occurrence	34
Panel Mine (Rio Algom Limited)	17	Napray Occurrence	35
Roche Long Lac Prospect	18	Larson Township	35
Roman Island Occurrence	19	McDonough Occurrence	35
Stanrock Mine (Denison Mines Limited)	19	Long Township	35
Day Township	20	Location X Occurrence	35
Cullis Lake Occurrence	20	Pronto Mine (Rio Algom Limited)	36
Gaiashk Township	20	McAughy Township	36
Corner Lake Occurrence	20	Gimby Occurrence	36
Iso Mines Occurrence (Whiskey Zone)	20	Morin Township	37
Pecors Lake East Prospect (Whiskey Zone)	21	McIntyre Porcupine Occurrence	37
Whiskey Prospect (Whiskey Zone)	21	Nicholas Township	37
Giles Township	22	Crazy Lake Occurrence	37
Soo-Tomic Occurrence	22	Peever Township	38
Greenwood Township	22	Damascus Occurrence	38
Phillips Occurrence	22	Dolan Occurrence	38
Gunterman Township	23	Hennessy Occurrence	38
Buckles Mine (Rio Algom Limited)	23	J.G. McCombe Occurrence	38
Crotch Lake Occurrence	23	Patrick Occurrence	39
Genex Prospect	24	Ranwick Prospect	39
		Van Lake Occurrence	39
		Piche Township	40
		Golden Arrow Occurrence	40
		Inspiration Occurrence	40
		Poulin Township	41
		Cobalt Consolidated Occurrence	41
		Reilly Township	41
		Ranger Occurrence	41

Rix Township	42	Gaudette Township	59
Canagau Dike Occurrence	42	Alur Occurrence	59
Root Township	42	Gunterman Township	59
Aubrey Falls Occurrence	42	International Occurrence	59
Sagard Township	42	Thorncrest Radioactivity Occurrence	60
Caneonti Prospect	42	Jarvis Township	60
Grey Trout Lake Prospect	43	Reserve Lake Occurrence	60
Harvard Prospect	44	Johnson Township	60
Shedden Township	44	Desbarats Radioactivity Occurrence	60
Denvic Lake Occurrence	44	Juillette Township	60
Slater Township	44	Fano Radioactivity Occurrence	60
Batchawana Uranium Mines		Matinenda Radioactivity Occurrence	60
Occurrence	44	Kamichisitit Township	61
Bobcam Occurrence	45	Blind River Occurrence	61
Camray Occurrence	45	Kirkwood Township	61
Theano Point Prospect	46	Kirkwood Township Radioactivity	
Mosher-Byles Occurrence	46	Occurrence	61
Robb-Murmac Prospect	46	Labonte Township	61
Soo-Tomic Occurrence	47	Huclif Porcupine Radioactivity	
Smilsky Township	47	Occurrence	61
Fausten Occurrence	47	Jalore Occurrence	61
Jay-Dee Occurrence	48	Lapaska Occurrence	62
Labine-McCarthy Occurrence	48	McDevitt Occurrence	62
Spragge Township	49	Tooker Radioactivity Occurrence	62
Spragge Creek Occurrence	49	Lecaron Township	63
Timmermans Township	49	Kirkpatrick Lake Radioactivity	
Anabar Occurrence	49	Occurrence	63
Big Game Occurrence	49	Lefroy Township	63
Buffana Prospect	50	Frye Radioactivity Occurrence	63
Coffee Lake Occurrence	51	Webb Occurrence	63
Denison Occurrence	51	Lehman Township	63
Dominion Prospect	51	Bracemar Radioactivity Occurrence	63
Jeanette Prospect	52	Mack Township	64
Martin Occurrence	52	Black Lake Radioactivity	
Moon Lake Occurrence	53	Occurrence	64
Picton Occurrence	53	Maeck Township	64
Pistol Lake Prospect	54	Seabrook Lake Carbonatite Complex	64
Zenmac Occurrence	54	McMurray Township	64
Tupper Township	55	Firesand River Carbonatite Complex	64
Pitchore Prospect	55	Surluga Occurrence	64
Vankoughnet Township	55	Monestime Township	64
Vankoughnet Occurrence	55	Russian Lake Radioactivity	
Viel Township	55	Occurrence	64
Gaitwin Occurrence	55	Montgomery Township	65
Mattaini Occurrence	56	Demorest Lake Occurrence	65
Rawhide Occurrence	56	Morin Township	65
Minor Uranium and Thorium Occurrences of		Sutherland Occurrence	65
Algoma District	58	Raimbault Township	65
Albanel Township	58	Boymar Occurrence	65
Arco Triller No. 2	58	God's Lake Mines Occurrence	65
Bolger Township	58	Zenmac Radioactivity Occurrence	66
Peerless Uranium Mining Corporation		Rix Township	66
Limited (Exploration Area)	58	Falconbridge Occurrence	66
Day Township	58	Roche Occurrence	66
Sowerby Occurrence	58	Rollins Township	66
Duncan Township	58	James Occurrence	66
Duncan Township Occurrence	58	Sagard Township	67
Maud Lake Occurrence	59	Mount Lake Area	67
Gaiashk Township	59	Salter Township	67
Canuc Occurrence	59	Shunck Occurrence	67

Slater Township	67	Bridges Township	81
Manwood Radioactivity Occurrence	67	Coulee Occurrence (Kimber Lake - West Occurrence)	81
Pennwood Occurrence	67	Fairservice Occurrence	81
Smilsky Township	68	Headway Occurrence	82
Ranrouyn Occurrence	68	Docker Township	82
Strain Township	68	Kimber Lake (East) Occurrence (C.S. Stephens)	82
Madawanson Lake Occurrence	68	Drope Township	83
Striker Township	68	Bluett Lake Occurrence	83
Cyr Occurrence	68	Langton Township	83
Tarbutt Township	68	Loudon Prospect	83
Tarbutt Mines Occurrence	68	Preston Occurrence	83
Thessalon Township	68	MacNicol Township	84
West Thessalon Occurrence	68	Foot Lake Occurrence	84
Timmermans Township	69	Hawk Lake Occurrence	
Anuwon Occurrence	69	(Byberg-Campbell-Macfarlane Occurrence)	84
Fano Radioactivity Occurrence	69	Kenoratomic Occurrence	85
Fort Norman Radioactivity Occurrence	69	Richard Lake Prospect (New Campbell Island)	85
Viel Township	70	Temple Township	86
Weston Radioactivity Occurrence	70	Bottle Bay Occurrence	86
Whitman Township	70	Tustin Township	86
Crestland Occurrence	70	Ascot Occurrence	86
Uranium and Thorium Deposits of Cochrane District	71	Petursson Lake Occurrence	87
Pitt Township	71	Wabigoon Township	87
Calmor Occurrence	71	Benchmark Occurrence	87
Minor Uranium and Thorium Occurrences of Cochrane District	72	Minor Uranium and Thorium Occurrences of Kenora District	88
Habel Township	72	Occurrences on Unsurveyed Land	88
Soweska River Occurrence	72	Belair Lake Radioactivity Occurrence	88
Pitt Township	72	Chase Lake Radioactivity Occurrence	88
Coral Rapids Occurrence	72	Hornby Lake Occurrence	88
Shackleton Township	72	Huston Lake Radioactivity Occurrence	88
Provencher Occurrence	72	Johnson Lake Radioactivity Occurrence	88
Uranium and Thorium Deposits of Kenora District	73	Oak Lake Occurrence	89
Deposits on Unsurveyed Land	73	Octopus Lake Radioactivity Occurrence	89
Aerobus Lake Occurrence	73	Paterson Lake Occurrence	89
Bamaji Lake Occurrence	73	Pineneedle Lake Radioactivity Occurrence	89
Bearhead Lake Prospect	74	Pine Road Occurrence	89
Cameron-Alcock Occurrence	75	Reynar Lake Occurrence	90
Can-Fer Occurrence	75	Sydney Lake Radioactivity Occurrence	90
Davidson Lake Occurrence	76	Bridges Township	90
Halliday Occurrence	76	Parth Occurrence	90
Headvue Occurrence	77	Wilson Occurrence	90
Jenson Occurrence	77	McBrien Township	90
Pancer Occurrence	77	Coal Creek Occurrence	90
Perch Lake Occurrence	78	McMaster Township	91
Pirson Occurrence (Medicine Stone Lake Occurrence)	78	Asarco Occurrence	91
Schryburt Lake Occurrence	79	McNicol Township	91
Snook Lake Occurrence	79	Viceroy Occurrence	91
Tourist Lake Prospect	79		
Treelined Lake Occurrence	80		
Vermilion Lake Occurrence	80		
Wolf Island Occurrence	80		

Tustin Township	91	May Township	107
Bee Lake Occurrence	91	Dominion Gulf Occurrence	107
Paddington Lake Occurrence	91	Parkin Township	108
Wabigoon Township	92	Assembly 17A Occurrence	108
Quibell Occurrence		Porter Township	108
(Meehan Occurrence)	92	Brewis-White Occurrence	108
Uranium and Thorium Deposits of Nipissing District	93	Chemical Research Occurrence	109
Pardo Township	93	Hunter Lake Occurrence	110
Pickle Crow Occurrence	93	New Mylamaque (Mymar) Occurrence	110
Vogt Township	93	Roberts Township	111
Proscop Prospect	93	Nordic Resources Prospect	111
Minor Uranium and Thorium Occurrences of Nipissing District	95	Roberts Lake Occurrence	111
Strathcona Township	95	Shakespeare Township	112
Neil Occurrence	95	Alexander Occurrence	112
Uranium and Thorium Occurrences of Rainy River District	96	Broulan Reef Occurrence (Sutherland Creek Occurrence)	112
Mainville Lake Occurrence	96	Dominion Gulf Occurrence	113
Minor Uranium and Thorium Occurrences of Rainy River District	97	McDonald Bay Occurrence	113
Trottier Township	97	Stetham Township	114
Heward Lake Occurrence	97	Jonsmith Occurrence (McKinnon Occurrence)	114
Uranium and Thorium Occurrences of Sudbury District	98	Stobie Township	114
Baldwin Township	98	Stobie Township Occurrence	114
Broulan Reef Central Occurrence	98	Turner Township	115
Broulan Reef Lookout Tower Occurrence	98	Harrison Occurrence	115
Canadian Johns-Manville Occurrence	99	Minor Uranium and Thorium Occurrences of Sudbury District	116
Jellicoe (Espanola Bay) Occurrence	99	Baldwin Township	116
Jellicoe Occurrence	99	Dominion Gulf Occurrence	116
Plum Occurrence	100	United Cobalt Occurrence	116
Creelman Township	100	Bigwood Township	116
Leslie Occurrence	100	French River Occurrence (South Side of Main Channel)	116
Drury Township	101	Cassidy Township	116
Alanen-Maki Occurrence	101	Aubinadong River Occurrence	116
Kerr Addison Prospect	102	Clary Township	116
Ermatinger Township	102	Noranda Occurrence (Marcia Lake Occurrence)	116
Balboa Occurrence	102	Collins and Chewett Townships	117
Grigg Township	103	Nemegosenda Lake Occurrence	117
Canadian Johns-Manville Occurrence	103	Creelman Township	117
Hallam Township	103	North Creelman Occurrence	117
Aggressive Occurrence	103	Curtin Township	117
Hutton Township	104	Bridges Occurrence	117
Assembly Occurrence	104	Demorest Township	117
Assembly 1A Occurrence	104	Noranda Occurrence (Yorston Lake Occurrence)	117
Hyman Township	105	Fraleck Township	118
Noranda Occurrence	105	Ingamar Occurrence	118
Agnew Lake Mine (Agnew Lake Mines Limited)	106	Gallagher Township	118
East Bay Occurrence	106	Borden Lake Occurrence	118
MacLennan Township	107	Harrow Township	118
Pictou Occurrence (Leclerc Occurrence)	107	Stencill Occurrence	118
		Hyman Township	118
		Richore Occurrence	118
		Langlois Township	119
		Kram-Glowaski Occurrence	119
		MacLennan Township	119
		Chevrette Occurrence	119
		Massey Bay Occurrence	119

Marconi Township	119	Bomby Township	130
Canadian Johns-Manville		Hemlo Occurrence	130
Occurrence	119	Conmee Township	130
Can-Fer Occurrence	120	Kakabeka Falls Occurrence	130
McNish Township	120	Dorion Township	130
Saville Occurrence	120	Dorion Amethyst Mine	130
Moncrieff Township	120	Goldie Township	130
Hollinger Radioactivity		Nelson Occurrence	130
Occurrence	120	Hele Township	131
Nairn Township	120	Tessier-Williamson Occurrence	131
Deauville Occurrence	120	Laberge Township	131
Porter Township	121	Molson Lake Occurrence	131
Macfie Occurrence	121	Punkari Occurrence	131
Pennbec Occurrence	121	McCoy Township	131
Wadge Occurrence	121	Port Monroe (Potvine) Occurrence	131
Seagram Township	121	McIntyre Township	131
Denison Occurrence	121	Port Arthur Occurrence	131
Shakespeare Township	122	McTavish Township	132
Blue Occurrence	122	Enterprise Mine	132
Delcan Occurrence	122	Pearl Lake Occurrence	132
Pick Occurrence	122	O'Neill Township	132
Satellite Occurrence	122	Cramette Occurrence	132
Schlesinger Occurrence	123	Pic Township	132
Stetham Township	123	Smith Occurrence	132
Astrabrun Radioactivity		Walsh Township	132
Occurrence	123	Deadhorse Creek (Diatreme)	
Turner Township	123	Occurrence)	132
Denison (Bull Lake) Occurrence	123	McKellar Creek (Diatreme)	
Noranda (Pilgrim Creek)		Occurrence	133
Occurrence	123		
Victoria Township	124	Uranium and Thorium Deposits of	
Sugar Lake Occurrence	124	Timiskaming District	134
Towle-Humpage Occurrence	124	Lebel Township	134
		Kaplan Occurrence	134
Uranium and Thorium Deposits of Thunder			
Bay District	125	Minor Uranium and Thorium Occurrences of	
Occurrences on Unsurveyed Land	125	Timiskaming District	135
Greenwich Lake Occurrence	125	Boston Township	135
New InSCO Prospect (Prairie Lake		Empire Occurrence	135
Complex)	126	Lunge Occurrence	135
New Santiago (Sturgeon Narrows)		Yost Occurrence	135
Occurrence	126	Bucke Township	135
Sandy Stone (Charron Lake)		Harrison-Hibbert Occurrence	135
Occurrence	127	Cane Township	135
Sandy Stone (Kassagimini Lake)		Cane Township Occurrence	135
Occurrence	127	Coleman Township	136
Daley Township	128	Kerr Lake Mine	136
Longlac Occurrence	128	Gillies Limit Township	136
Dorion Township	128	Silvermaque Mine	136
Innes Lake Occurrence	128	Hudson Township	136
		Spencer Occurrence	136
Minor Uranium and Thorium Occurrences of		Lebel Township	136
Thunder Bay District	129	Bidgood Kirkland Mine	136
Occurrences on Unsurveyed Land	129	Hurd Occurrence	136
Celotti Occurrence	129	Lebel Occurrence	136
Pic Bamooos Occurrence (Port		Tudhope Township	137
Coldwell Complex)	129	Sauve-Inch Occurrence	137
Squaw Lake Occurrence	129		
Univex Occurrence	129	References	138
Yzerdraat Occurrence	129		
		Index	147

Figures

- | | |
|--|---|
| 1. Ontario Districts and Counties | 2 |
| 2. Principal Uranium Deposits of Ontario | 3 |

Tables

- | | |
|--|---|
| 1. Classification of Uranium Deposits in Ontario | 5 |
| 2. Production Data for the Blind River - Elliot Lake - Agnew Lake Camp | 6 |

Conversion Factors for Measurements in Ontario Geological Survey Publications

CONVERSION FROM SI TO IMPERIAL			CONVERSION FROM IMPERIAL TO SI		
<i>SI Unit</i>	<i>Multiplied by</i>	<i>Gives</i>	<i>Imperial Unit</i>	<i>Multiplied by</i>	<i>Gives</i>
LENGTH					
1 mm	0.039 37	inches	1 inch	25.4	mm
1 cm	0.393 70	inches	1 inch	2.54	cm
1 m	3.280 84	feet	1 foot	0.304 8	m
1 m	0.049 709 7	chains	1 chain	20.116 8	m
1 km	0.621 371	miles (statute)	1 mile (statute)	1.609 344	km
AREA					
1 cm ²	0.155 0	square inches	1 square inch	6.451 6	cm ²
1 m ²	10.763 9	square feet	1 square foot	0.092 903 04	m ²
1 km ²	0.386 10	square miles	1 square mile	2.589 988	km ²
1 ha	2.471 054	acres	1 acre	0.404 685 6	ha
VOLUME					
1 cm ³	0.061 02	cubic inches	1 cubic inch	16.387 064	cm ³
1 m ³	35.314 7	cubic feet	1 cubic foot	0.028 316 85	m ³
1 m ³	1.308 0	cubic yards	1 cubic yard	0.764 555	m ³
CAPACITY					
1 L	1.759 755	pints	1 pint	0.568 261	L
1 L	0.879 877	quarts	1 quart	1.136 522	L
1 L	0.219 969	gallons	1 gallon	4.546 090	L
MASS					
1 g	0.035 273 96	ounces (avdp)	1 ounce (avdp)	28.349 523	g
1 g	0.032 150 75	ounces (troy)	1 ounce (troy)	31.103 476 8	g
1 kg	2.204 62	pounds (avdp)	1 pound (avdp)	0.453 592 37	kg
1 kg	0.001 102 3	tons (short)	1 ton (short)	907.184 74	kg
1 t	1.102 311	tons (short)	1 ton (short)	0.907 184 74	t
1 kg	0.000 984 21	tons (long)	1 ton (long)	1016.046 908 8	kg
1 t	0.984 206 5	tons (long)	1 ton (long)	1.016 046 908 8	t
CONCENTRATION					
1 g/t	0.029 166 6	ounce (troy)/ ton (short)	1 ounce (troy)/ ton (short)	34 285 714 2	g/t
1 g/t	0.583 333 33	pennyweights/ ton (short)	1 pennyweight/ ton (short)	1.714 285 7	g/t

OTHER USEFUL CONVERSION FACTORS

1 ounce (troy)/ton (short)	20.0	pennyweights/ton (short)
1 pennyweight/ton (short)	0.05	ounce (troy)/ton (short)

NOTE—Conversion factors which are in bold type are exact. The conversion factors have been taken from or have been derived from factors given in the Metric Practice Guide for the Canadian Mining and Metallurgical Industries published by The Mining Association of Canada in cooperation with the Coal Association of Canada.

CONVERSION FACTORS FOR URANIUM

Number of tonnes U x 1.29987 = number of short tons U₃O₈
 Number of short tons U₃O₈ x 0.76931 = number of tonnes U

ABSTRACT

This, the second edition of the uranium-thorium deposit inventory, describes briefly the deposits of uranium and/or thorium in northern Ontario, which for the purposes of this circular is defined as that part of Ontario lying north and west of the Grenville Front. The most significant of the deposits described are fossil placers lying at or near the base of the Middle Precambrian Huronian Supergroup. These include the producing and past-producing mines of the Elliot Lake - Agnew Lake area. Also included are the pitchblende veins spatially associated with Late Precambrian (Keweenawan) diabase dikes of the Theano Point - Montreal River area. Miscellaneous Early Precambrian pegmatite, pitchblende-coffinite-sulphide occurrences near the Middle-Early Precambrian unconformity fringing the Lake Superior basin, and disseminations in diabase, granitic rocks, alkalic complexes and breccias scattered throughout northern Ontario make up the rest of the occurrences.

Exploration for uranium in Ontario was carried out initially in the period 1948 to 1957 with production at Elliot Lake starting in 1955. Peak production was reached in 1959, but cut back due to lack of civilian markets. Exploration resumed in 1965 with optimism for the future of nuclear power development. However, neither markets nor discoveries matched expectations and the exploration pace again slackened. Further exploration took place in the late 1970s as the price of uranium rose sharply on the open market. Long-term contracts were signed between Ontario Hydro and Preston Mines Limited (now Rio Algom Limited) and Denison Mines Limited, ensuring production well into the 21st century, and also ensuring the orderly expansion of the mining and milling facilities, the townsites at Elliot Lake, and the regional infrastructures. Eldorado Nuclear is locating its principal refinery facilities at Blind River.

However, on the international scene, there are still delays in the licensing of nuclear power stations, and predictions of demand in the medium term are still being lowered as these delays, high capital costs, and high interest rates force cancellations or deferral of planned nuclear stations. On the supply side, recent discoveries of high grade deposits particularly in Saskatchewan, Northern Territory of Australia, and Central Africa have led to concentration of the continuing exploration in these areas. Nevertheless, given a modest increase in uranium price relative to costs, Ontario with its resources and infrastructure and operating know-how at Elliot Lake may well be able to compete successfully for future contracts.

Production from the Elliot Lake camp to the end of 1981 totalled 98 300 tonnes U (256 million pounds U_3O_8) for an average recovered grade of 0.099 percent U (2.36 pounds U_3O_8 per ton) and an additional 671 tonnes U (1.75 million pounds U_3O_8) recovered by leaching at Agnew Lake. There has also been minor production of thorium and yttrium. Should a market for thorium develop, the Elliot Lake uranium deposits would constitute a major resource.

URANIUM AND THORIUM DEPOSITS OF NORTHERN ONTARIO

by

James A. Robertson¹ and Kerry L. Gould²

INTRODUCTION

This circular briefly summarizes data on uranium and/or thorium deposits in northern Ontario and supercedes Mineral Resources Circular 9, published in 1968 (Robertson 1968a). It is one of a group of Mineral Resource Group publications providing up-to-date information on Ontario's uranium deposits, their classification, distribution, resources, and factors affecting their development or further resources. The reader is referred to Robertson et al. (1981); Robertson (in press); Robertson and Gould (1981); and Runnalls (1981). Background on the Canadian uranium exploration and extraction industries is provided by the annual reports of the Uranium Resources Appraisal Group (Energy, Mines and Resources, Canada); international information is given by the biannual reports of the Organization for Economic Organization and Development (OECD), the annual deliberations of the Uranium Institute, and the industry workshop sponsored annually by the United States Department of Energy.

A new edition of this circular has been necessitated by the considerable amount of data which has entered the public realm as a result of:

1. declassification of previously confidential material;
2. new information collected by the Ministry's regional and geological staff, some of which has been published in maps and reports of the Ontario Geological Survey, and some

of which is filed in the Geoscience Data Files of the Ontario Geological Survey or in the files of the Resident Geologists;

3. two cycles of exploration for uranium completed since the first edition;
4. new production data; and
5. the naming of townships in northern Ontario previously designated by numbers.

The principal uranium and thorium deposits of Ontario occur in the Blind River - Elliot Lake - Agnew Lake area and the Bancroft area. The deposits in the former area are fossil placers at or near the unconformity separating the Middle Precambrian Huronian Supergroup and the Early Precambrian (Robertson 1981, 1983; Adams and Button 1981; Houston and Karlstrom 1981), and those in the latter area are pegmatitic granite dikes, pegmatites, metasomatic replacement deposits and calcite-fluorite-apatitic veins in rocks of the Grenville Supergroup of the Grenville Province (Robertson et al. 1981; Gordon et al. 1981; Robertson 1981). Only the pyritic quartz-pebble conglomerate of the Elliot Lake - Agnew Lake camp and certain pegmatites of the Bancroft area have supported production.

For the purposes of this inventory, the boundary between northern and southern Ontario has been taken as the Grenville Front, rather than a district or administrative boundary, because of the geological implication.

Figure 1 shows Ontario district and county boundaries. Figure 2 summarizes the geology of Ontario and shows the distribution of the principal deposit types using a classification comparable with that used by the Uranium Resource Appraisal Group (Energy, Mines and Resources, Canada) and international publications such as OECD (Organization for Economic Cooperation and Development), and IUREP (International Uranium Resources Evaluation Project). Table 1 shows the classification of deposit types used in Ontario. For expanded discussion of the Ontario deposits the reader

¹Policy Advisor, Mineral Resources Branch, Ontario Ministry of Natural Resources, Toronto.

²Geological Assistant, Mineral Deposits Section, Ontario Geological Survey, Toronto. Manuscript approved for publication by the Chief Geologist, January 13, 1982. This report is published with the permission of E.G. Pye, Director, Ontario Geological Survey.



Figure 1. Ontario Districts and Counties.




Figure 2. Principal Uranium Deposits of Ontario.

is referred to Robertson (1981) and the maps showing distribution of these deposit types with respect to geology (Robertson et al. 1981; Robertson 1982; Robertson and Gould 1981).

Table 2 lists the production data for the Blind River - Elliot Lake - Agnew Lake camp. Minor amounts of thorium and yttrium have also been produced but data is not in the public realm.

This circular also includes the pitchblende occurrences of the Theano Point - Montreal River area where the pitchblende veinlets are found in or near the contact zone of the Late Precambrian (Keweenawan) diabase dikes (Nuffield 1956; Robertson 1968a, 1978). There has been no production from the Theano Point area but it is of historical interest as the first recorded occurrence of pitchblende in Canada and the rediscovery of the area in 1948 was an important step leading to the discovery of Elliot Lake -- a world class deposit. The circular also lists a number of pitchblende-coffinite-sulphide occurrences, possibly related to the Early Precambrian - Middle Precambrian unconformity adjacent to the Lake Superior Basin.

A number of other deposit types such as Early Precambrian pegmatites, disseminations in diabase, alkalic complexes and breccias and two areas where there are deposits possibly genetically related to Early Precambrian volcanic soils are described.

Uranium and thorium are potential by-products from some carbonatite complexes which contain the material of commercial interest, and the Elliot Lake uranium deposit is a major potential source of thorium.

ACKNOWLEDGMENTS

The writers are greatly indebted to the Resident Geologists and Mineral Resource Coordinators attached to the Ministry's regional staff in northern Ontario for much of the data summarized in the following paper. Many company officials, prospectors, and individuals have given access to their records and their assistance over many years is gratefully acknowledged.

The senior author wishes to acknowledge the continued assistance afforded him by the professional and support staff of the Ontario Geological Survey during the preparation and processing of this report.

EXPLANATION OF THIS CIRCULAR

Uranium occurrences are listed alphabetically by district, township and deposit name. It should be noted that since publication of the first edition all numbered townships in Ontario have been named. Further, as many showings and occurrences change hands frequently, it is generally more convenient to use a geographic or a well known name as the permanent name for an occurrence. This may mean that a deposit is named for the discoverer or a previous owner rather than the present or latest owner. An index to names and location is provided at the back of this circular. The reader may find it convenient to refer to the Mineral Map of Ontario (Map 2310) or the map of uranium and thorium deposits of Ontario, which shows the districts, townships, mining districts, and the location of the deposit).

Low grade uranium deposits or ones on which little or no work has been done are listed separately under "Minor Uranium and Thorium Occurrences" for each district.

In general, only the most up-to-date references are quoted for description and location. However, where there are additional references of particular significance, these may also be included. Publications such as the annual "Canadian Mines Handbook", published by Northern Miner Press, or the annual "Survey of Mines" and "Survey of Energy Resources", published by the Financial Post, contain information on many of the companies listed.

Data has been collected continuously since the preparation of the first edition. Information on property developments does not extend beyond the end of 1980.

Data is given in the measurement format in which it was originally recorded and a list of conversion factors is given on page xi. Note that early measurement of radioactivity and early assay methods, particularly those based on a non-discriminating radiation measuring device, should not be necessarily considered reliable but only indicative of anomalous radioactivity. In recent years field examination of occurrences has generally included the use of discriminating radiation measurement devices. In more detailed studies, the operation of equilibrium is also addressed. In re-examining old showings the "state of the art" at the time of the original work should be borne in mind.

**TABLE 1.
CLASSIFICATION OF URANIUM DEPOSITS
IN ONTARIO**

Magmatic Deposits

1. Carbonatite bodies and alkalic complexes, including fenites
2. Calc-alkaline extrusive rocks

Magmatic - Metamorphic Deposits

3. Pegmatite:
 - a) red zoned
 - b) red unzoned
 - c) white
4. Calc-silicate (skarn) rocks and pyroxenite

Sedimentary (Detrital Deposits)

5. Quartz-pebble conglomerate:
 - a) pyrite-uranium-thorium
 - b) iron oxide-thorium
6. Polymictic conglomerate, semipelitic rocks

Circulating Mineralized Solutions*

7. a) Pitchblende with simple mineralogy (Athabasca type)
- b) Pitchblende with multi-metal mineralogy (Bohemia type)
- c) Mineralization of unspecified nature, generally related to an unconformity and found in argillite, greywacke, sandstone, carbonate rock, and lignite.
- d) Uraninite-calcite-apatite-fluorite veins.

Unclassified Deposits

8. Usually small occurrences about which little is known

* Largely equivalent to the vein deposits of the OECD (1979) and of Robertson (1975), or the "hydrogenic deposits" of McMillan (1977).

Mine	Year	Production Million \$
Rio Algom		
Lacnor Mine ^b	1960	14.02
Milliken Mine ^d	1958	11.89
	1959	21.63
	1960	23.69
Nordic Mine	1957	22.59
	1958	26.88
	1959	25.54
	1960	23.14
	1965	30.10
	1966	27.52
Panel Mine ^b		
Pronto Mine ^e	1955	0.49
	1956	7.28
	1957	11.02
	1958	12.22
	1959	13.64
	1960	3.35
Quirke Mine	1956	1.62
	1957	22.84
	1958	22.90
	1959	23.05
	1960	22.79
All Mines	1961	72.93
	1962	46.10
	1963	48.86
	1964	25.90
Total Rio Algom		588.07
Stanrock	1958	7.00
	1959	17.99
	1960	22.71
	1961	22.62
	1962	20.70
	1963	10.91
	1964	8.44
	1965	0.81
	1966	0.75
Total		111.94
Grand Total*		<u>\$ 1,203.84</u>

Table 2 Notes

Sources: Runnalls (1981) and Company Annual Reports

*to end 1966, representing 128,160,096 pounds U₃O₈ from 56,077,179 tons milled

^aConsolidated Denison Mines Limited 1957-1959.

^bNorthspan Uranium Mines Limited includes dollar value for Lacnor, Panel, and South American Mines 1957-1959.

^cStanleigh Uranium Mining Corporation Limited 1958-1959.

^dMilliken Lake Uranium Mines Limited 1959-1959.

^ePronto Uranium Mines Limited 1955-1959.

URANIUM AND THORIUM DEPOSITS OF ALGOMA DISTRICT

ABERDEEN TOWNSHIP

RIO TINTO PROSPECT

Commodity
Uranium, thorium.

Radioactive Minerals
Unknown.

Location
Latitude 46°29'55"N, Longitude 83°43'50"W.
Aberdeen Township.
Map Reference: OGS Map 2419.

Geology
The prospect is underlain by mafic metavolcanics and pink medium to coarse grained feldspathic quartzite, brown to green arkosic quartzite, and quartz-pebble conglomerate of the Middle Precambrian Huronian Matinenda Formation.

Economic Features
Assays of samples from drill holes by Amax Exploration Incorporated averaged 0.31 pounds U₃O₈ per ton over 3.0 feet.

History of Development
1955: Six drill holes were completed by Sagamore Exploration.
1956: Six drill holes, totalling 2025 feet, were located in N1/2 lot 3, concession IV, S1/2 lot 3, concession V, S1/2 lot 4, concession V, and N1/2 lot 4, concession V. The holes were drilled by Rio Tinto Canadian Exploration Limited.
1968: Airborne magnetic, electromagnetic and radiometric surveys were performed by Amax Exploration Incorporated.
1969: Ten drill holes totalling 6914 feet were drilled by Amax Exploration Incorporated.
1970: A geological survey was completed by Amax Exploration Incorporated.

References
Giblin et al. (1979)
Robertson (1968a, p.4)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.
Technical File No. 63.3026, Amax Exploration Incorporated, 1970.
Technical File No. 63.2443, Amax Exploration Incorporated, 1969.
Aberdeen Township, Drill Log Report No. 10, Amax Exploration Incorporated, 1969.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File No. 000001.

ALBANEL TOWNSHIP

LITTLE WHITE RIVER OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°34'45"N Longitude 82°58'34"W.
Albanel Township.
Map Reference: OGS Map 2419.

Geology
This occurrence lies in the Southern Province and is underlain by Middle Precambrian Huronian metasediments. Surface exposures consist of Mississagi quartzite of the Hough Lake Group.

Economic Features
Selected samples gave assay values of 0.04 percent U₃O₈ (0.8 pounds) (Assessment Files Research Office, Ontario Geological Survey, Toronto, Technical File No. 63A.534).

Radioactivity of 10 to 20 times the background is not uncommon. Locally this may rise to 40 times the background where quartz pebbles are present.

History of Development
1968: Geological survey was completed by G.E. Parsons.
1969: Ground magnetic survey and one hole to 3002 feet were completed by the Hanna Mining Company.
1974: A multi-sensor airborne geophysical survey was done by Fort Norman Explorations Incorporated.

References
Giblin et al. (1979).
Siemiakowska et al. (1975).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.
Albanel Township, Drill Log Report No. 17, The Hanna Mining Company, 1969.

Technical File No. 63A.534, G.E. Parsons, 1968.
 Technical File No. 63.3461, The Hanna Mining Company, 1969.
 Technical File No. 2.1575, Fort Norman Explorations Limited, 1974.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
 File No. 000265.

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.
 Beange Township, Drill Log Report No. 10, Pitchgoma Mines Limited, 1953-1954.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
 File No. 000115.

BEANGE TOWNSHIP

CANDORE PROSPECT

Commodity
 Uranium.

Radioactive Minerals
 Unknown.

Location
 Latitude 46°30'42"N, Longitude 82°43'10"W.
 Beange Township.
 Map Reference: OGS Map 2419.

Geology
 Sedimentary rocks lie unconformably on Early Precambrian granite which was intersected by seven drill holes ranging in depth from 311 feet to 652 feet. Slightly radioactive thin oligomictic pebble-conglomerate bands in Upper Mississagi Formation with pyrite in the matrix were intersected in drilling.

Economic Features
 The bands are normally less than 6 inches thick, and assayed less than 0.02 percent U₃O₈. The best assay was 0.043 percent U₃O₈ over 1 foot.

History of Development
 1953-1954: Geological and radiometric surveys were performed. Seventeen drill holes totalling approximately 7500 feet were drilled for Pitchgoma Mines Limited.
 1955: One hole was drilled to 2244 feet, near Dunlop Lake by Delta Minerals Limited.
 1966: The claim group was optioned to Denison Mines Limited.
 1966-1967: Three drill holes totalling 5028 feet were drilled by Denison Mines Limited.
 1975: The claim group was sold to Gateway Uranium Mines Limited.

References
 Giblin et al. (1979).
 Robertson (1968a, p.46).

CALLINAN FLIN-FLON OCCURRENCE

Commodity
 Uranium.

Radioactive Minerals
 Unknown.

Location
 Latitude 46°32'00"N, Longitude 82°47'26"W.
 Beange Township.
 Map Reference: ODM Map 2015.

Geology
 This occurrence lies in the Southern Province and is underlain by Middle Precambrian Huronian metasediments comprising greenish, coarse-grained arkose and quartzite with interbeds and lenses of slightly radioactive pyritized pebble conglomerate of the Mississagi Formation of the Hough Lake Group.

Economic Features
 A geiger survey showed numerous readings of 0.02, 0.03, and 0.05 mR/h (millirontgens/hour). The area with the highest readings is on the southwest shore of Gibbery Lake. Here, readings of 0.10 and 0.14 mR/h were obtained over a length of 350 feet in quartz-pebble conglomerate closely associated with pea-green quartzite.

History of Development
 1955: Ground radiometric and geological surveys were performed. Three drill holes totalling 871 feet were completed on former claim numbers SSM 25756 and SSM 25745. One hole to 522 feet was drilled in Rainbault Township (formerly Township 157). Work was done by Callinan Flin-Flon Mines Limited.

1957: One hole to 50 feet was drilled by Consolidated Callinan Flin Flon Mines Limited.

1970: Airborne magnetic and electromagnetic surveys were performed for A. St. Denis.

References
 Robertson (1963, p.53,61).

Robertson (1968a, p.47).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Beange Township, Drill Log Report
No. 14, Callinan Flin-Flon Mines
Limited, 1955.

Technical File No. 63A.239, Callinan
Flin-Flon Mines Limited, 1955.

Technical File No. 63.2736, A. St
Denis, 1970.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000116

RIO ALGOM (QUIRKE GROUP WEST) OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°30'31"N, Longitude 82°42'16"W.
Beange Township.
Map Reference: ODM Map 2015.

Geology
This occurrence is underlain by Middle
Precambrian Huronian metasediments
comprising coarse-grained, green arkose with
slightly radioactive oligomictic pebble bands
of the Mississagi Formation of the Hough
Lake Group.

Economic Features
Unknown.

History of Development
1954-1955: Geological and geophysical
surveys and three drill holes were completed
by Algom Uranium Mines Limited.
1954: Algom Uranium Mines Limited merged
with other companies to form Rio Algom
Mines Limited. The name was later changed
to Rio Algom Limited.

References
Robertson (1963, p.55).

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000117.

RIO ALGOM (SPAN-NORTH) OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°31'05"N, Longitude 82°43'56"W.
Beange Township.
Map Reference: ODM Map 2015.

Geology
This occurrence is underlain by Middle
Precambrian Huronian metasediments,
comprising Mississagi quartzite and arkose of
the Hough Lake Group, with interbeds and
lenses of slightly radioactive pebble
conglomerate. Scattered thin pebble bands
with minor pyrite were intersected in drill
holes.

Economic Features
Assays averaged 0.32 percent U₃O₈ over
0.5 feet.

History of Development
1954-1955: Geological and airborne
scintillometer surveys were completed. Ten
drill holes totalling 8558 feet were drilled by
Panel Consolidated Uranium Mines Limited.
1957-1958: Five drill holes totalling 2833 feet
were drilled by Northspan Uranium Mines
Limited.
1956: Panel Consolidated Uranium Mines
Limited merged into Northspan Uranium
Mines Limited and property was transferred
to Span-North Mining Claims Limited.
1960: Span-North Mining Claims Limited
merged with Algom Uranium Mines Limited.

References
Robertson (1963, p.75-77).
Robertson (1968a, p.47-48).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.
Beange Township, Drill Log Report
No. 11, Panel Consolidated Uranium
Mines Limited, and Northspan
Uranium Mines Limited, 1954-1958.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000118

BOLGER TOWNSHIP**GUI-POR PROSPECT****Commodity**

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°23'57"N, Longitude 82°49'20"W.
Bolger Township.

Map Reference: ODM Map 2014.

Geology

The uranium mineralization is located within the Matinenda Formation of the Elliot Lake Group. Three diamond drill holes encountered oligomictic pebble bands, and the assays averaged 0.040 percent U₃O₈ over 5.2 feet.

History of Development

1954-1955: Geological and scintillometer surveys were completed. Fifteen diamond drill holes totalling 6923 feet were drilled by New Jersey Zinc Exploration Company (Canada) Limited.

Further surface exploration work was done by Rio Tinto Canadian Exploration Limited.

1974: Airborne magnetic, electromagnetic and gamma ray spectrometer surveys were completed by Fort Norman Explorations Incorporated.

References

Robertson (1963, p.65).

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Technical File No. 63A.303, New Jersey Zinc Exploration Company (Canada) Limited, 1954-1955.

Technical File No. 2.1575, Fort Norman Exploration Incorporated, 1974.

Bolger Township, Drill Log Report No. 13, Gui-Por Uranium Mines and Metals Limited, 1954-1955.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000111.

MOON LAKE MINES OCCURRENCE (MOON LAKE ZONE)

Commodity

Uranium and thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°24'10"N, Longitude 82°49'30"W.
Bolger Township.

Map Reference: ODM Map 2014.

Geology

Only thin bands and lenses of slightly radioactive conglomerates considered to belong to the Matinenda Formation of the Elliot Lake Group were present.

Economic Features

Average assay results from four drill holes were 0.054 percent U₃O₈ over 3.2 feet.

History of Development

1954-1955: Geological and scintillometer surveys were completed. Four holes were drilled totalling 2188 feet in Bolger Township, and twenty holes were drilled in Timmermans Township. All work was completed by New Jersey Zinc Exploration Company (Canada) Limited.

1957: Surface mapping was completed by Rio Tinto Canadian Exploration Limited.

1967: An airborne magnetometer survey was completed by Denison Mines Limited.

1974: Airborne electromagnetic, magnetometer, radiometric, and resistivity surveys were completed by Fort Norman Explorations Incorporated.

References

Robertson (1963, p.70).

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Technical File No. 63A.303, New Jersey Zinc Exploration Company (Canada) Limited, 1955.

Bolger Township, Drill Log Report No.14, Moon Lake Uranium Mines Limited, 1955.

Technical File No. 63A.342, Moon Lake Uranium Mines Limited, 1958.

Technical File No. 2.1575, Fort Norman Explorations Incorporated, 1974.

Bolger Township, Drill Log Report No. 39, Fort Norman Explorations Incorporated, 1976.

Timmermans Township, Drill Log Report No. 43, Fort Norman Explorations Incorporated, 1976.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000112.

NORDIC GROUP WEST PROSPECT (MOON LAKE ZONE)

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°23'58"N, Longitude 82°45'16"W.
Bolger Township.
Map Reference: ODM Map 2014.

Geology
Conglomerate of the Bruce Formation is exposed along the north boundary of the area. Minor intersections of oligomictic conglomerate bearing traces of U₃O₈ were encountered.

History of Development

1953-1958: Geological and geophysical surveys and extensive drilling totalling over 17,044 feet was completed by Algom Uranium Mines Limited.

1960: Algom Uranium Mines Limited amalgamated with Rio Algom Mines Limited.

1966: An airborne magnetic survey was completed by B.B. Scott and L.T. Chandler.

1970: An airborne spectrometer survey was completed by Kerr-McGee of Canada Limited.

1978: An airborne geophysical survey was completed by Fort Norman Explorations Incorporated.

References

Robertson (1963, p.54).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Bolger Township, Drill Log Report No. 10, 20, 21, 22, and 23, Algom Uranium Mines Limited.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000113.

BOUCK TOWNSHIP

BENNER OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°29'53"N, Longitude 82°38'40"W.
Bouck Township.
Map Reference: ODM Map 2114.

Geology
Diamond drilling on this claim intersected uranium-bearing conglomerate of the Mississagi Formation from depths of 2679 feet to 2707 feet.

Economic Features

Assays of the mineralized zone showed 0.025 percent U₃O₈ over 11 feet, from 2679 to 2690 feet and 0.061 percent U₃O₈ over 13.25 feet from 2693.5 to 2706.75 feet.

History of Development

1954: One diamond drill hole totalling 2785 feet was drilled by Ralph Benner.

1975: This area now belongs to Denison Mines Limited.

References

Robertson (1967).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Bouck Township, Drill Log Report No. 13, R. Benner, 1954.

DENISON MINE

Commodity
Uranium, yttrium, and thorium.

Radioactive Minerals
Brannerite, monazite, uraninite,
uraniothorite, and uranothorianite.

Location
Latitude 46°29'35"N, Longitude 82°35'54"W.
Bouck Township.
Map Reference: OGS Map 2419.

Geology

The main orebody (Denison Reef) occurs from 0 to 180 feet above the Early

Precambrian basement and 210 to 230 feet below an argillite layer. It is a tabular-shaped mass covering a projected surface area of 8000 feet by 14,000 feet. The ore zone averages 16.7 feet thick and strikes S65E with an average dip of 19°S.

The ore zone is composed of two layers known as the "A" and "B" Zones. They average 8 feet in thickness. The average depth of the ore zone is 1000 feet at the north end and 3000 feet at the south end. The "C" Zone above the "A" and "B" averages 6 by 100 feet in size. The "D" and "E" Zones are conglomerate beds 50 and 70 feet below the argillite.

The ore-bearing conglomerate is a uranium-bearing oligomictic quartz-pebble conglomerate.

Economic Features

From 1961 to 1980, production from the Denison Mine was 34,376,100 tons of ore, from which 41,297 tons U₃O₈ were recovered for an average grade of 0.12 percent U₃O₈.

History of Development

1956: The No. 1 shaft (1857 feet deep and six compartments) was completed. The trackless mining method was initiated.

1957: The second shaft (eight compartments to 2750 feet) was completed. Crushing and acid leach plants began production. The mill was rated at 6000 tons per day. Mining by the room and pillar method was used.

1960: Consolidated Denison Mines Limited and Can-Met Explorations Limited amalgamated to form Denison Mines Limited.

1970: A 6200 foot connecting drive was completed between the two mines.

1973: Denison Mines Limited amalgamated with Stanrock Uranium Mines Limited. The mill rate was up to 7000 tons per day, producing 12,000 to 13,000 pounds of U₃O₈ and 400 pounds of yttrium.

1978: Production was 2,404,000 tons of ore with an average millhead grade of 2.15 pounds per ton; 4,989,000 pounds of U₃O₈ were recovered.

Dewatering, expansion, and technological upgrading took place at the Stanrock and Can-Met Mines.

The capacity of the central processing plant increased to 15,000 tons per day. Mining capacity increased to 21,000 tons per day.

1980: Underground expansion was completed including the dewatering of the Stanrock Mine. The mill is to be completed in 1981. Production was 2,510,000 tons of ore with an average millhead grade of 1.87 pounds per ton; 4,451,823 pounds of U₃O₈ were recovered.

References

- Canadian Mines Handbook 1979-1980.
Giblin et al. (1979).
Griffith (1967, p.102-124).
Robertson (1968a, p. 109-110).
Runnalls (1981).
The Northern Miner (November 6, 1975, p. 1).

GEMICO PROSPECT

Commodity

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°27'16"N, Longitude 82°40'06"W.
Bouck Township.
Map Reference: OGS Map 2419.

Geology

Diamond drill holes have traced the Cobalt, Quirke Lake, Hough Lake and Elliot Lake Groups down to the basement rocks. The Matinenda Formation of the Elliot Lake Group is composed of zones of quartz-pebble conglomerate in a matrix of arkose.

Economic Features

Based on a potential zone 15,000 feet by 4,000 feet by 36 feet at 0.7 pounds per ton, reserves are estimated at 180 million tons of ore and 126,000,000 pounds U₃O₈ (Annual Report, 1975, Canuc Mines Limited). Available assays for one sample show 0.5 pounds ThO₂ over 8 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto, Bouck Township, Drill Log Report No. 25, 1967-1969).

History of Development

1955-1956: Four drill holes totalling 4016 feet were drilled near the north shore of Banana Lake. Work by Kerr-McGee Corporation.

1966-1974: Airborne magnetometer, electromagnetic, radiometric surveys, and three drill holes totalling 14,932 feet were completed by Kerr-McGee Corporation.

References

- Canuc Mines Limited (1975, annual company report).
Giblin et al. (1979).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Bouck Township, Drill Log Reports No.
25 and 28, Kerr-McGee Corporation,
1967-1969.

Bouck Township, Drill Log Report No.
15, Buffana Uranium Mines Limited,
1955-1956.

Technical File No. 63.2802, Kerr-
McGee Corporation, 1970.

Technical File No. 63.2145, Kerr-
McGee Corporation, 1966.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 001106.

KERR-MCGEE 150-151 OCCURRENCE

Commodity

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°27'45"N, Longitude 82°36'16"W.
Bouck Township.

Map Reference: OGS Map 2419.

Geology

Diamond drill holes in this occurrence indicated formations of the Huronian Supergroup including the Cobalt, Quirke Lake, Hough Lake and Elliot Lake Groups. Oligomictic quartz-pebble conglomerates of the Matinenda Formation are mined by Rio Algom Mines Limited and Denison Mines Limited in the vicinity.

Economic Features

The Matinenda Formation was intersected at the 4656-foot level and extended to 4967 feet. Average assays are 0.083 percent U₃O₈ and 0.123 percent ThO₂ over 3.0 feet.

History of Development

1966: Airborne electromagnetic surveys were completed by B.B. Scott and L.T. Chandler, and the Kerr-McGee Corporation.

1968: One diamond drill hole to 5002 feet was completed by Kerr-McGee Corporation.

References

Giblin et al. (1979).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Bouck Township, Drill Log Report No.
24, Kerr-McGee Corporation, 1968.

Technical File No. 63.1898, B.B.
Scott, 1966.

Technical File No. 63.2074, Kerr-
McGee Corporation, 1966.

QUIRKE MINE NO. 1 (RIO ALGOM LIMITED)

Commodity

Uranium, thorium, and rare earths.

Radioactive Minerals

Brannerite, uraninite, thorite, and uranothorite.

Location

Latitude 46°30'45"N, Longitude 82°39'48"W.
Bouck Township.

Map Reference: ODM Map 2114.

Geology

Uranium mineralization occurs within oligomictic quartz-pebble conglomerates at the base of the Matinenda Formation. The ore zone outcrops at the north end of the property and reaches a depth of 1600 feet at the south end. In the vicinity of the deposit are a series of conglomerate horizons called the "Upper" Reef and the "A" Reef. In the "A" Reef, ore occurs as lenses of uraniferous quartz-pebble conglomerate interbedded with quartzite. The strike length is 7300 feet, the average thickness is 12 feet and the down slope length is 2000 feet.

Economic Features

From 1956 to 1961 4,173,327 tons of ore were milled, producing 8,709,429 pounds of U₃O₈ at an average grade of 2.34 pounds U₃O₈ per ton.

In 1966, 102,515 pounds of U₃O₈ were recovered from the mine waters.

History of Development

1953: After surface trenching and sampling, geological surveys and airborne scintillometer surveys were completed. Two hundred and three drill holes totalling 87,548 feet were drilled along the strike of the conglomerate at 200-foot intervals.

1955: A shaft was sunk to 861 feet. A 77-foot adit was added.

1956: A 3000 ton-per-day mill was put into operation.

1961: In March of this year operations were suspended. Total underground development included 74,747 feet of drifts, and 19,924 feet of crosscuts. All of this and previous work was completed by Algom Uranium Mines Limited.

1965: The mine was dewatered and the waters were treated for recovery at the Nordic mill.

1967: Production resumed from the No. 1 shaft.

1968: The Quirke mill was operating at 3,700 tons per day. The Rio Algom Limited production data given in Table 2 includes clean-up for Nordic as well as Quirke No. 1 production. Leachings from Nordic are included in the 1969 and 1970 production. 1971: Operations at this mine ceased in the latter part of this year at which time operations were transferred to the New Quirke Mine (Quirke No. 2).

Total development as of December, 1971, was 98,893 feet of drifts and 19,938 feet of crosscuts. A further 1798 feet of diamond drilling were completed underground. The total from 1969 to 1971 was 19,895 feet.

All the work was completed by Rio Algom Mines Limited.

References

Griffith (1967).
Robertson (1968a).
Rio Algom Limited, annual company reports.

Energy, Mines and Resources Canada Files
Mineral Resources Branch, Department of
Energy, Mines and Resources, Ottawa.

File 41 J/10, U6, Quirke No. 1, May 1973.

NEW QUIRKE MINE NO. 2 (RIO ALGOM LIMITED)

Commodity

Uranium and thorium.

Radioactive Minerals

Uraninite, brannerite, and monzite.

Location

Latitude 46°30'16"N, Longitude 82°37'15"W.
Bouck Township.
Map Reference: OGS Map 2419.

Geology

Uranium mineralization occurs within oligomictic quartz-pebble conglomerates at the base of the Matinenda Formation.

The best development is in the Denison "C" Reef some 100 feet below the Quirke "A" Reef. The Denison Reef usually contains conglomerate zones each 6 to 12 feet thick separated by arkose 2 to 3 feet thick. Radioactive minerals usually constitute 10 to 15 percent of the conglomerate's pyritic, sericitic feldspathic quartz matrix.

Economic Features

In 1980 the average grade was 2.1 pounds per ton U_3O_8 , at a recovery rate of 4886 pounds U_3O_8 from a mill rate of 6975 tons of ore per day.

The production data for Quirke No. 2 is included in Table 2. In 1971, the Quirke mill also milled ore from Quirke No. 1. After the re-opening of the Panel mine, minor amounts of Panel minewater and contaminated of development ore from Panel were also treated.

History of Development

1953: This property was staked by Preston East Dome Mines Limited. The property was subsequently sold to Algom Uranium Mines Limited.

1960: The New Quirke shaft was begun. Operations were suspended. The company name was change to Rio Algom Limited.

1966: The shaft was deepened to 2260 feet. 1968: The No. 1 mine and mill were reopened. Production from the New Quirke Mine began.

1973: Total underground development amounted to 99,995 feet of drifts and 13,194 feet of crosscuts. Diamond drilling from 1968 to 1973 included 7261 underground drill holes totalling 104,260 feet and 10 surface drill holes totalling 8237 feet.

1974: Expansion program was initiated. The mill rate was to increase to 7000 tons per day by 1978.

1976: The mine expanded to the "A" Reef to the west of the New Quirke shaft. The milling rate was up to 6350 tons of ore per day. The Quirke No. 1 was dewatered and the mine waters leached. Plans included using the Quirke No. 1 mine as a ventilation shaft.

1978: Production came from the New Quirke "C" and "A" Reefs.

1979: The Quirke mill capacity was up to 7004 tons ore per day. The average recovery rate per ton was 2.3 pounds U_3O_8 per ton.

1980: Production declined due to lower grades.

References

Giblin et al. (1979).
Rio Algom Limited, annual company reports.
Rio Algom Mines Limited (1972).
Robertson (1968a, p.118,121).
Robertson (1976a).
Robertson (1981).

Energy, Mines and Resources Canada Files
Mineral Resources Branch, Department of
Energy, Mines and Resources, Ottawa.

File 41 J/10 U3, Quirke No. 2 (New Quirke) May 1973.

SPANISH AMERICAN MINE (RIO ALGOM LIMITED)

Commodity
Uranium.

Radioactive Minerals
Brannerite and monazite.

Location
Latitude 46°28'37"N, Longitude 82°35'16"W.
Bouck Township.
Map Reference: ODM Map 2114.

Geology

The Spanish American property is underlain by sparse conglomerates and feldspathic quartzite of the Gowganda Formation except for a narrow strip along the shore of Quirke Lake where white to pink rocks of the Serpent Formation are exposed. A northwest-striking lineament, partly filled with Nipissing diabase, marks the steeply dipping Spanish American Fault.

The ore zone is a conglomerate unit up to 20 feet thick, of which an average of 10 feet is ore grade. The body strikes N70W, dips 17°S and is 2100 feet long by 400 feet wide.

The conglomerate consists of closely packed, subrounded quartz pebbles and cobbles in a matrix containing pyrite, hematite, magnetite, and zircon.

The thorium to uranium ratio is approximately 3 to 1.

Economic Features

Little et al. (1972) gave the mill production as 764,000 pounds U₃O₈ from 422,000 tons of ore milled. This figure includes ore purchased from Algom Uranium Mines Limited and ore from the Buckles mine; full details were not published.

History of Development

1953: The 36 claim property was staked by P. Westfield. Two diamond drill holes were drilled by Kinloch Mining Company Limited. 1953-1959: Spanish American Mines Limited drilled 12 holes all of which intersected uranium-bearing zones. This drilling indicated about 6,000,000 tons grading 0.10 percent U₃O₈.

Spanish American Mines Limited was incorporated into Northspan Uranium Mines Limited in 1956. No. 1 and 2 shafts were completed to 3476 feet and 3163 feet. A level was established at a depth of 3100 feet, on which 7056 feet of drifting and 4286 feet of crosscutting were completed by February, 1959. The mine was closed in February, 1959, because of severe ground conditions.

From 1956 to 1959 diamond drilling included 33 surface holes totalling 4276 feet and 432 underground holes totalling 14,833 feet.

1961: Northspan amalgamated with three other companies to form Rio Algom Limited.

References

Griffith (1967, p.173-175).
Robertson (1968a, p.44).
Robertson (1968b, p.122).

Energy, Mines and Resources Canada Files
Mineral Resources Branch, Department of Energy, Mines and Resources, Ottawa.
File 41/J U 47, Spanish American Mine, May 1975.

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File No. 000110.

BUCKLES TOWNSHIP

CAN-MET MINE (DENISON MINES LIMITED)

Commodity
Uranium.

Radioactive Minerals
Brannerite, monazite, and uraninite.

Location
Latitude 46°28'51"N, Longitude 82°32'54"W.
Buckles Township.
Map Reference: ODM Map 2002.

Geology

The Can-Met deposit is located on the north limb of the Quirke Syncline. The orebody which consists of two beds of uraniferous quartz-pebble conglomerate, is in one of the fringe areas of the major ore shoot, located on the north side of the Quirke Lake Trough.

A major easterly striking, high angle reverse fault, a large 100-foot thick dike and many minor normal and reverse faults transect the orebody.

Large pockets of pyrrhotite, pyrite, and some chalcopyrite have been found. The average depth of the ore zone is 1750 feet and occurs in an area 1400 feet by 1800 feet. The average strike is N40W, with dips ranging between 0 and 15°S. The average thickness of the ore is 17 feet, but widths of up to 25 feet have been encountered. Two shafts about 500 feet apart were sunk to depths of 2127 and 2395 feet respectively (Griffith 1967, p. 98-101).

Economic Features

From the start of production in 1957 to September 30, 1959, 2,489,824 pounds of U_3O_8 were recovered from 1,477,000 tons of ore milled. Production from October, 1959, to closure in April, 1960, has not been published.

History of Development

1954-1960: Can-Met Explorations Limited did the following development work. Two shafts were sunk 2127 and 2395 feet respectively. Shaft No. 1 is a two-compartment service and ventilation shaft with a station at 2089 feet. Shaft No. 2 is a three-compartment production shaft with a station at 2080 feet.

Total diamond drilling consisted of 11 holes from surface totalling 24,426 feet and 68 holes from underground totalling 11,302 feet. Total accumulated lateral development consisted of 32,389 feet of drifting and 2591 feet of crosscutting.

1960: Can-Met Explorations Limited amalgamated with other companies to form Denison Mines Limited. In this year, operations ceased.

1978: Dewatering of the Can-Met Mine was begun.

1979: Can-Met Shaft No. 1 was enlarged and used as a ventilation shaft for the Stanrock Mine.

References

Griffith (1967, p.98-101).
Robertson (1961, p.48).
Financial Post Survey of Mines (1959, Can-Met Explorations Limited).

CONECHO PROSPECT

Commodity

Uranium.

Radioactive Minerals

Brannerite, uraninite, and monazite.

Location

Latitude $46^{\circ}29'09''N$, Longitude $82^{\circ}31'16''W$.
Buckles Township.
Map Reference: ODM Map 2002.

Geology

Surface exposures show the Mississagi Formation of the Hough Lake Group and a thick diabase sill intruding the Mississagi Formation between Quirke and Teesdale Lakes. Drilling intersected uraniferous oligomictic conglomerate bands.

Economic Features

Drilling indicated a minimum of 2,000,000 tons of material averaging 0.07 percent U_3O_8 across an average width of 6 feet.

History of Development

1953: Airborne magnetometer and scintillometer survey was completed by Technical Mine Consultants Limited.

1954-1955: Twenty drill holes totalling 13,273 feet and located on the northeast shore of Teasdale Lake were drilled by Conecho Mines Limited.

1975: Conecho Mines Limited amalgamated with Rio Algom Limited.

1980: This property is now considered part of the Panel property.

References

Robertson (1957, p.49).
Robertson (1968a, p.33).

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Buckles Township, Drill Log Report No. 11, Conecho Mines Limited, 1954.
Technical File No. 63.419, Technical Mine Consultants Limited, 1953.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File No. 000091.

PANEL MINE (RIO ALGOM LIMITED)

Commodity

Uranium and thorium.

Radioactive Minerals

Brannerite, uraninite, and monazite.

Location

Latitude $46^{\circ}29'54''N$, Longitude $82^{\circ}33'00''W$.
Buckles Township.
Map Reference: OGS Map 2419.

Geology

The Panel orebody underlying Quirke Lake is on the north limb of the Quirke Lake Syncline. The ore-bearing zone is a quartz-pebble conglomerate of the Matinenda Formation of the Elliot Lake Group. The orebody is thickest in the northern part of the mine. The grade becomes richer to the west and some narrow sections of very high grade ore have been reported. Some assays show 0.85 percent U_3O_8 per ton.

The main ore zone, varying in thickness from 6 to 32 feet and averaging 12 feet,

strikes N75W and dips 14°S. The depths at which the ore was intersected range from 1100 to 1700 feet. Locally there are high concentrations of thorium.

Economic Features

According to the Ontario Department of Mines and Northern Affairs statistical records, a total of 3,628,000 tons of ore was milled, producing 7,818,000 pounds (Little et al. 1972) of U₃O₈. Average assays from nine drill holes showed 0.10 percent U₃O₈ over 13.8 feet.

History of Development

1953: Airborne magnetometer and scintillometer surveys were completed by Technical Mine Consultants Limited.

1956-1959: Northspan Uranium Mines Limited was responsible for the work carried on during this time. Two shafts about 1900 feet apart were sunk to depths of 1836 and 1250 feet. The mine was developed on 12 levels. Production first began in 1958 and the mill reached a capacity of 3000 tons of ore per day in August of 1958.

Total diamond drilling consisted of 26 holes for 23,792 feet from the surface and 1037 holes underground totalling 35,208 feet. Total footage during underground development consisted of 29,587 feet of drifts and 19,231 feet of crosscuts.

1960-1961: Northspan amalgamated with other companies to form Rio Algom Mines Limited.

In 1961 the Panel Mine was closed.

1977: The Panel Mine was part of Rio Algom's Phase 2 expansion. The mine was reactivated. The mill rate was 2 990 tonnes per day.

1979: Commercial production commenced in November averaging 2419 tons of ore per day with an average grade of 1.2 pounds U₃O₈ per ton. A total of 394,000 pounds of U₃O₈ were produced (Rio Algom Limited, Annual Report 1980) (see Table 2).

References

Lang et al. (1962, p.141).
Giblin et al. (1979).
Griffith (1967, p.152-162).
Rio Algom Limited, annual company reports.
Robertson (1968a, p.33-34).
Western Miner (July 1956, Vol.29, No.7, p.139).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Technical File No. 63.419, Technical
Mines Consultants Limited, 1953.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.

File No. 000094.

Sault Ste. Marie Regional Geologist's Files,
SSM 561, SSM 376, SSM 712, SSM
1072.

ROCHE LONG LAC PROSPECT

Commodity

Uranium, thorium.

Radioactive Minerals

Brannerite, uraninite, and monazite.

Location

Latitude 46°28'34"N, Longitude 82°31'28".
Buckles Township.

Map Reference: ODM Map 2002.

Geology

Drilling intersected the Espanola Formation at the surface, then penetrated the Bruce, Mississagi, Pecors, Ramsay Lake and Matinenda Formations, and the Early Precambrian granite at depths of 1850 to 2467 feet.

Economic Features

Average assay results from the uraniferous oligomictic conglomerate intersected in three drill holes were 0.06 percent U₃O₈ over a core length of 5.6 feet (Leahy 1973; Mineral Development Section, Energy Mines and Resources Canada, Ottawa, File N9, 41/J/7).

In November, 1967, reserves were estimated at 1.6 million tons grading 0.07 percent U₃O₈.

History of Development

1954-55: A geological survey and seven drill holes totalling 12,465 feet were completed by Roche Long Lac Mines Limited. Later, this company's name changed to North Rock Explorations Limited.

1967: Stanrock Uranium Mines Limited acquired a majority of shares.

1973: Amalgamated with Denison Mines Limited.

References

Leahy (1973).
Robertson (1961, p.54).
Robertson (1968a, p.32).

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto
 Technical File No. 63A.232, Roche
 Long Lac Mines Limited, 1954
 Source Mineral Deposits Record, Ontario
 Geological Survey, Toronto.
 File No. 000093

Energy, Mines and Resources Canada Files
 Mineral Development Section, Department of
 Energy, Mines and Resources, Ottawa.
 National Mineral Inventory File U9, 41
 J/7, North Rock, January, 1968.

ROMAN ISLAND OCCURRENCE

Commodity
 Uranium.

Radioactive Minerals
 Unknown.

Location
 Latitude 46°29'17"N, Longitude 82°34'20"W.
 Buckles Township.
 Map Reference: ODM Map 2002.

Geology
 One drill hole intersected uraniferous
 oligomictic conglomerate of the Matinenda
 Formation of the Elliot Lake Group.

Economic Features
 Assays of samples range from 0.01 to 0.36
 percent U₃O₈ (Leahy 1973).

History of Development
 1955: One drill hole to 1950 feet was
 completed by Consolidated Denison Mines
 Limited.
 1960: Amalgamated with Denison Mines
 Limited.

References
 Leahy (1973).
 Robertson (1961, p.18,20).

STANROCK MINE (DENISON MINES LIMITED)

Commodity
 Uranium and yttrium.

Radioactive Minerals
 Uraninite, brannerite, monazite, and
 thucholite.

Location
 Latitude 46°28'19"N, Longitude 82°33'41"W.
 Buckles Township.
 Map Reference: ODM Map 2002.

Geology
 The Stanrock orebody strikes northwest and
 dips an average of 17 degrees to the
 southwest. The ore minerals are associated
 with pyrite in the matrix of the quartz-
 pebble conglomerate of the Matinenda
 Formation of the Elliot Lake Group. The
 orebody is displaced by a major thrust fault,
 which strikes northwest and dips 15°S. The
 dip slip is approximately 1200 feet and the
 strike slip is approximately 300 feet. The
 resulting duplication of the orebody adds
 considerably to the ore potential along a
 broad band paralleling the fault. One major
 diabase dike 80 feet thick strikes east across
 the property and cuts nearly vertically
 through the ore zone. The ore was
 intersected at a depth of 3200 feet in the No.
 1 shaft and about 2700 feet in the No. 2
 shaft. The beds of ore average 6 feet in
 thickness.

Economic Features
 From 1958 to July 1964, 6,446,800 tons of
 ore were milled, producing 11,402,746 pounds
 of U₃O₈ at an average of 1.77 pounds U₃O₈
 recovered per ton. Salvage leaching
 operations terminated in early 1970 and
 brought the total production to 12,362,000
 pounds U₃O₈.

As of December, 1968, estimated reserves
 were approximately 10,000,000 tons assured
 and probable averaging about 1.5 pounds
 U₃O₈ per ton (Financial Post Survey of
 Mines, 1971).

History of Development
 1955: Diamond drilling was completed by
 Stancan Uranium Mines Limited.
 1956-1970: Stanrock Uranium Mines Limited
 completed the following work.
 Two shafts were sunk 797 feet apart. The
 main three-compartment production shaft
 was deepened to 3379 feet and a two-
 compartment service shaft to 2953 feet.
 Main levels were established at 3277 and
 2909 feet.

In 1958 a mill to handle 3300 tons per day
 went into operation.

Conventional mining ceased in October
 1964. Thereafter production came from
 leaching. In 1965 yttrium oxide was produced.

In April 1970 operations were suspended.
 Total diamond drilling consisted of 15 holes
 from the surface totalling 40,417 feet and
 4668 holes underground totalling 50,119 feet.

Total development in the mine was 114,045 feet of drifts and 2861 feet of crosscuts.

1973: Stanrock Uranium Mines Limited amalgamated with Denison Mines Limited.

1978: Dewatering of the mine was begun.

1979: The shafts were improved and deepened.

1981: Underground development was to be completed by 1982.

References

Robertson (1961, p.59).

Robertson (1968a, p.34-35).

Financial Post Survey of Mines (1971, Stanrock Mines Limited).

Western Miner (July 1956, Vol.29, No.7, p.153-156).

Ontario Ministry of Natural Resources Files

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000094.

DAY TOWNSHIP

CULLIS LAKE OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°17'50"N, Longitude 83°24'10"W.
Day Township.

Map Reference: OGS Map 2419.

Geology

At the southern end of Cullis Lake 7 to 10 m of coarse, pink arkose and subarkose overlies grey subarkose of the Livingston Creek Formation. At the base of the pink arkose is about 50 to 75 cm of quartz-pebble conglomerate which is locally pyritic, particularly near the base. Samples of pyritic conglomerate were weakly radioactive.

References

Bennett (1976, p.111-113).

Frarey (1977).

Giblin et al. (1979).

GAIASHK TOWNSHIP

CORNER LAKE OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°26'30"N, Longitude 82°25'58"W.
Gaiashk Township.

Map Reference: ODM Map 2003.

Geology

A deep drill hole intersected the Middle Precambrian Huronian sequence from the Gowganda Formation at the surface, through the Serpent, Espanola, Bruce, Mississagi, Pecors, Ramsay Lake and Matinenda Formations, to the Early Precambrian basement. Radioactive sections of quartz pebbles with minor sulphides were intersected.

History of Development

1967-1968: Combined airborne magnetic and electromagnetic surveys and two drill holes totalling 6359 feet were completed by Cominco Limited.

References

Giblin et al. (1979).

Robertson (1962, p.45).

Shklanka (1969, p.88).

Ontario Ministry of Natural Resources Files: Assessment Files Research Office, Ontario Geological Survey, Toronto.

Gaiashk Township, Drill Log Report
No. 38, Gulf Minerals, 1968.

Technical File No. 63.2171, Cominco
Limited, 1967.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.

File No. 000025

ISO MINES OCCURRENCE (WHISKEY ZONE)

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°23'13"N, Longitude 82°24'35"W.
Gaiashk Township.
Map Reference: ODM Map 2003.

Geology

This occurrence is underlain by Middle Precambrian Huronian metasediments. Uranium mineralization occurs in thin sections of pyritiferous, uraniferous, oligomictic conglomerate of the Matinenda Formation.

Economic Features

One drill hole gave average assays of 0.013 percent U₃O₈ over 2.3 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto: Gaiashk Township, Drill Log Report No. 40, 1968).

History of Development

1953-1957: Geological, airborne magnetometer and radiometric surveys were performed, and nine holes were drilled by Algom Uranium Mines Limited.

1968: Two drill holes totalling 943 feet on claim S140707 were drilled by Iso Mines Limited.

1974: An airborne magnetometer survey was completed by North American Nuclear Limited.

References

Giblin et al. (1979).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.

Gaiashk Township, Drill Log Report No. 13, Preston East Dome Mines Limited, 1953.

Technical File No. 63.441, Algom Uranium Mines Limited, 1953

Gaiashk Township, Drill Log Report No. 40 Iso Mines Limited, 1968.

Technical File No. 2.1783, North American Nuclear Limited, 1974.

PECORS LAKE EAST PROSPECT (WHISKEY ZONE)**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°22'46"N, Longitude 82°25'26"W.
Gaiashk Township.
Map Reference: ODM Map 2003.

Geology

Uranium mineralization is located in quartz-pebble conglomerate and arkose of the Matinenda Formation of the Elliot Lake Group. The Matinenda Formation is intruded by a Nipissing diabase sill and the outcrop is repeated by a thrust fault on the hanging wall of the sill. The conglomerate is restricted to an old stream channel (Robertson 1962).

Economic Features

Shallow exploration and assessment drilling was carried out on two surface exposures of radioactive quartz-pebble conglomerate found on claim S67641 and S64478-64479 (Claim Map M1212 Gaiashk Township). The drilling delineated a very low grade, narrow conglomerate zone, which averages 4.5 feet in thickness with a strike length of about 1300 feet and has an average grade of 0.05 percent U₃O₈ (Robertson 1968a).

History of Development

1953-1957: Geological and airborne magnetometer surveys were carried out. Four drill holes totalling 15,873 feet were drilled by Algom Uranium Mines Limited.

1974: An airborne magnetometer survey was carried out by North American Nuclear Limited.

References

Robertson (1962, p.62-63).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.

Technical File No. 63.441, Algom Uranium Mines Limited, 1953.

Technical File No. 63.441, Algom Uranium Mines Limited, 1953.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000083.

WHISKEY PROSPECT (WHISKEY ZONE)**Commodity**

Uranium.

Radioactive Minerals

Brannerite, uraninite, and monazite.

Location

Latitude 46°24'00"N, Longitude 82°20'50"W.
Gaiashk Township.
Map Reference: ODM Map 2003.

Geology

The uranium mineralization occurs in the Matinenda Formation in the thin beds and lenses of oligomictic conglomerate within arkose quartzite.

Economic Features

Samples from one drill hole assayed 0.036 percent over 0.9 feet. Another drill hole drilled by Grand Chibougamau Mines Limited gave an assay of 0.03 percent U_3O_8 over a 10-foot width (Robertson 1962, p. 78)

History of Development

1954: One drill hole to 180 feet was drilled by British Columbia Explorers Limited.

One hole to 133 feet was drilled by Grand Chibougamau Mines Limited.

1955: One hole to 553 feet was drilled by Panel Consolidated Uranium Mines Limited.

References

Robertson (1962, p.73).

Robertson (1968a, p.28-29).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.

Gaiashk Township, Drill Log Report No. 25, British Columbia Explorers Limited, 1954.

Gaiashk Township, Drill Log Report No. 30, Panel Consolidated Mines Limited, 1955.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000079.

GILES TOWNSHIP**SOO-TOMIC OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Pitchblende.

Location

Latitude 47°28'33"N, Longitude 84°49'20"W.
Giles Township.

Map Reference: OGS Map 2419.

Geology

Radioactive mineralization occurs in a brecciated zone at the contact and in a Keweenawan diabase dike cutting an Early Precambrian granite-pegmatite complex. The breccia zone was traced for 80 feet along the footwall and averaged 5 inches in

width. A parallel zone 4 feet away is 6 inches wide and was traced for 60 feet.

Economic Features

Samples from this deposit sent by J.G. McCombe assayed 1.62 percent U_3O_8 (radiometric equivalent) and up to 0.56 percent U_3O_8 (chemical). Samples sent by W. Jenks showed up to 7.30 percent U_3O_8 .

History of Development

Pre-1950: Sampling and prospected by J.G. McCombe.

1951: Soo-Tomic Uranium Mines Limited drilled 21 shallow holes totalling 671 feet.

1975: This occurrence is now within Lake Superior Provincial Park.

References

Giblin et al. (1979).

Lang (1952, p.124).

Robertson (1968a, p.27).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.

Drill Log Report, Soo-Tomic Uranium Mines Limited, 1951.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000077

GREENWOOD TOWNSHIP**PHILLIPS OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 47°27'00"N, Longitude 84°32'00"W.

Greenwood Township.

Map Reference: OGS Map 2419.

Geology

The radioactive samples were taken from pegmatites within the Early Precambrian basement. The basement is cut by a northwest-trending Keweenawan diabase dike.

Economic Features

Two samples from this deposit assayed 0.025 and 0.68 percent U_3O_8 (radiometric equivalent).

History of Development

Pre-1952: Some prospecting and sampling by R.B. Phillips.

References

Giblin et al. (1979).
Lang (1952, p.132).
Robertson (1968a, p.53).

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000060.

GUNTERMAN TOWNSHIP**BUCKLES MINE (RIO ALGOM LIMITED)****Commodity**

Uranium.

Radioactive Minerals

Brannerite, uraninite, monazite

Location

Latitude 46°22'34"N, Longitude 82°35'20"W.
Gunterman Township.
Map Reference: OGS Map 2419.

Geology

The orebody is the up-dip section of the Algom-Nordic ore horizon. The ore-bearing zone is composed of two quartz-pebble conglomerate beds. The lower bed rests on the older basement rocks; the upper bed is approximately 20 feet above the lower bed.

The ore outcrops along the southern limits of the property and is indicated at a depth of 210 feet at the northern extremity. The orebody dips 25°W for 4000 feet.

Economic Features

The total production of 189,598 tons of ore was shipped to the Spanish American and Lacnor mines. Production prior to 1958 was 124,890 tons of ore. In October, 1958, the mine was closed.

History of Development

1953: Airborne scintillometer and magnetometer surveys were performed by Technical Mine Consultants Limited.

1954-1955: Geological mapping, 34 drill holes totalling 3224 feet, and an exploratory drift to 77 feet were undertaken by Buckles Algoma Uranium Mines Limited.

1956: Claim group acquired by Spanish American Mines Limited.

1956: Spanish American amalgamated with others to form Northspan Uranium Mines Limited.

1956-1958: Northspan Uranium Mines Limited did the following development work.

The mine was opened by a three-compartment vertical shaft sunk 239 feet with levels at 160 and 211 feet (Griffith 1967, p.125).

The ore was milled at the Spanish American Mine and mostly at the Lacnor mill.

Approximately 250,000 tons of ore were mined before October 1958 when the mine was sealed.

Total diamond drilling consisted of one hole from the surface for 120 feet and three underground for 91 feet. Total footage during underground development consisted of 2730 feet of drifting and 462 feet of crosscutting.

1960: Northspan Uranium Mines Limited amalgamated to form Rio Algom Limited.

References

Giblin et al. (1979).
Griffith (1967, p.126).
Robertson (1968a, p.36).
Robertson (1968b, p.113).

Ontario Ministry of Natural Resources Files
Regional Geologist's Files, Sault Ste Marie.
SSM 336.

CROTCH LAKE OCCURRENCE**Commodity**

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°25'10"N, Longitude 82°36'34"W.
Gunterman Township.
Map Reference: OGS Map 2419.

Geology

One drill hole intersected the Huronian sequence from the Gowganda Formation at the surface, through the Serpent, Espanola, Bruce, Mississagi, Pecors, Ramsay Lake and Matinenda Formations, to the Early Precambrian basement at 3819 feet.

Economic Features

The uraniferous oligomictic conglomerate bands of the Matinenda Formation gave an average assay value of 0.38 pounds U₃O₈ per ton over 1.3 feet (Leahy 1973).

History of Development

1965-1967: Two drill holes to 490 and 3833 feet respectively were drilled by Rio Tinto Canadian Exploration Limited.

References

Giblin et al. (1979).
Leahy (1973).

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 001118.

GENEX PROSPECT

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°22'31"N, Longitude 82°35'54"W.
Gunterman Township.
Map Reference: OGS Map 2419.

Geology

A narrow strip of Matinenda quartzite with uraniferous conglomerate lies along the north edge of claim SSM 396273. These beds dip north at 15-20 degrees and rest on the Early Precambrian basement. Six drill holes intersected one or more radioactive conglomerate beds.

Economic Features

The main ore bed was 6.5 feet thick. There is approximately 10,000-11,000 tons of material of ore grade indicated.

History of Development

1954-1955: Geneva Lake Mines Limited performed surface mapping and drilled 12 holes totalling 2222 feet.
1957: 16 drill holes totalling 664 feet on the north shore of Nordic Lake were drilled by Genex Minex Limited.
1958: The claims were sold to Chipman Lake Mines Limited.
1976: Claim SSM 396273 was held by R. MacGregor of Sault Ste. Marie.

References

Giblin et al. (1979).
Robertson (1968a, p.40).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Gunterman Township, Drill Log Report
No. 12, Geneva Lake Mines, 1954.

Gunterman Township, Drill Log Report
No. 33, Genex Mines, 1957.

Technical File No. 63A.242, Geneva
Lake Mines Limited, 1955.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000096.

KAMIS OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°24'35"N, Longitude 82°40'45"W.
Gunterman Township.
Map Reference: ODM Map 2113.

Geology

Sedimentary rocks of the Hough Lake Group strike east and dip north at 10 to 30 degrees. The northwest-striking Home Lake fault crosses the area.

Economic Features

In one hole the average assay was 0.03 percent U_3O_8 over 3 feet. In another, assay values were 0.07 percent U_3O_8 and 0.03 percent ThO_2 over 1.0 foot.

History of Development

1955-1956: Geological survey and two drill holes for 6205 feet were completed by Kamis Copper Mines Limited.
1965-1966: Rio Tinto Canadian Exploration Limited drilled two holes for 4494 feet.

References

Robertson (1968a, p.40-41)
Robertson (1968b, p.111,136,143).

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Gunterman Township, Drill Log Report
No. 15, Kamis Copper Mines Limited,
1956.

Gunterman Township, Drill Log Report
No. 34, Rio Tinto Canadian
Exploration Limited, 1965.

Gunterman Township, Drill Log Report
No. 35, Rio Tinto Canadian
Exploration Limited, 1966.

Technical File No. 63A.292, Kamis
Uranium Mines Limited, 1955.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000097.

LACNOR MINE

Commodity
Uranium.

Radioactive Minerals
Brannerite and uraninite.

Location
Latitude 46°23'44"N, Longitude 82°36'31"W.
Gunterman Township.
Map Reference: OGS Map 2419.

Geology

The surface exposure is Mississagi quartzite of the Hough Lake Group dipping 20°N. The Nordic diabase sill-like intrusion lies to the north and is cut by northwest-striking dikes and a fault.

The Lacnor deposit is on the south limb of the Quirke Lake Syncline. The ore bed consists of uraniferous quartz-pebble conglomerate.

Two principal beds of conglomerate ore were found. The average thickness of the upper is 11 feet and the lower 12 feet. They are separated by a 14-foot bed of pebbly quartzite that is radioactive but not of ore grade. The strata strike about N70E and dips about 18°N. The east-west ore width indicated by drilling is approximately 3300 feet. The conglomerate consists of quartz pebbles in a matrix of feldspathic quartzite, well mineralized with pyrite (5 to 15 percent) and frequently containing a little chalcopyrite.

Economic Features

Diamond drilling indicated 5,202,054 tons of ore grading 0.111 percent U₃O₈.

History of Development

1954-1956: 19 drill holes totalling 44,179 feet were drilled.

1956-1960: Northspan Uranium Mines Limited did the following development work.

Two vertical, four-compartment shafts were sunk in 1956. No. 1 production shaft went to 2394 feet with shaft stations at 500, 1000, 1500, 2000, and 2300 feet. No. 2 service shaft is 2512 feet deep with stations at 84, 500, 1000, 1500, 2000, 2300, and 2400 feet. In 1959, No. 1 shaft was extended to 3510 feet with new levels at 2638, 2715, 2785, 2855, 2925, 3065, and 3135 feet. No. 2 shaft was extended to 2970 feet with new levels at 2637, 2714, 2854, and 2924 feet.

In 1960, No. 1 shaft was extended to 3663 feet and No. 2 to 3380 feet. A mill to handle 3922 tons per day went into operation in September 1957. By March 1958, the milling

rate reached 3770 tons per day. The mine was closed in 1960 and allowed to flood. During the period 1958-1960, 3,432,000 tons were milled and 6,309,000 pounds of U₃O₈ were recovered (Little et al. 1972).

Up to its 1960 closing, the mine had been developed on 16 levels. Total diamond drilling consisted of 123 holes from underground totalling 6112 feet and 50,993 feet of test-hole drilling from underground. Total footage underground consisted of 51,204 feet of drifting and 11,976 feet of crosscutting. 1960-1964: During 1964, mine waters were bled off through the Milliken Mine and the dissolved uranium was recovered. In July 1964, prior to the closing of the Milliken Mine, a raise was driven into the Lacnor workings above the water level. This work was by Rio Algom Mines Limited.

References

Giblin et al. (1979)
Robertson (1968b, p.81, 113-114, 116)

MILLIKEN MINE (RIO ALGOM LIMITED)

Commodity
Uranium.

Radioactive Minerals
Brannerite and uraninite.

Location
Latitude 46°24'09"N, Longitude 82°37'31"W.
Gunterman Township.
Map Reference: ODM Map 2113.

Geology

On surface, the property is underlain by Mississagi quartzite of the Hough Lake Group, dipping about 20°N. The Nordic diabase sill-like intrusion lies east and north of the mine site.

The Milliken Lake orebody is on the south limb of the Quirke Lake trough and is considered to be an extension of the Nordic and Lacnor beds. The ore zone is a quartz-pebble conglomerate of the Matinenda Formation.

Economic Features

The production shaft intersected two ore beds which are separated by 5 feet of quartzite at a depth of 3008 feet. The upper bed averaged 7 feet in thickness and contained 0.07 percent U₃O₈. The lower bed was 12 feet thick and averaged 0.12 percent U₃O₈. The service shaft intersected the upper ore bed at 2892 feet where it assayed 0.18 percent U₃O₈ over 7.5 feet. The lower

bed (separated from the upper by 2 feet of quartzite) was 6.6 feet thick and assay values were 0.14 percent U_3O_8 . The strike of the orebeds is N70W and the dip is 10 to 14°N (Lang et al. 1962, p.139).

Up to 1964, 14,204,530 pounds of U_3O_8 were produced from 6,325,960 tons of ore milled (Kelly 1960, 1961; Kelly and Riddell 1962, 1963, 1964; Riddell 1965, 1966, 1968).

History of Development

1954: Sixteen diamond drill holes totalling 45,210 feet and two shafts 550 feet apart and 300 feet deep were drilled by Milliken Lake Uranium Mines Limited.

1956-1965: The Rio Tinto Group took over in 1956. The shafts were extended to 3071 and 3400 feet. Mill construction was completed in March 1958. Plant capacity was 3200 tons of ore a day. The output of uranium concentrates was about 200,000 pounds per month.

Operations at the Milliken Mine ceased on June 30, 1964. An unsuccessful water leaching program was terminated the next year after yielding only 80,000 pounds of U_3O_8 .

During its operation the mine was developed on nine levels. Total diamond drilling consisted of 10,468 holes from underground totalling 134,301 feet. Total underground development consisted of 48,215 feet of drifting and 8,776 feet of crosscutting.

1975: This property was taken over by Rio Algom Limited.

1976: Rio Algom Limited planned to dewater the mine and widen the shafts.

References

- Canadian Mining Journal (October 1976, p.17)
 Griffith (1967, p. 130-137)
 Kelly (1960, 1961)
 Kelly and Riddell (1962, 1962, 1964)
 Lang et al. (1962, p.139)
 Riddell (1965, 1966, 1968)
 Robertson (1968b, p. 114-117)

Ontario Ministry of Natural Resources Files
 Source Mineral Deposits Record, Ontario
 Geological Survey, Toronto.
 File No. 00009.

NORDIC MINE (RIO ALGOM LIMITED)

Commodity

Uranium, thorium, and yttrium.

Radioactive Minerals

Brannerite, pitchblende, uraninite, and thucholite.

Location

Latitude 46°22'48"N, Longitude 82°35'20"W.
 Gunterman Township.
 Map Reference: OGS Map 2419.

Geology

The Nordic ore body is situated in a shallow valley structure on the south limb of the Quirke Lake trough. The Huronian units, over which a dip and scarp topography has developed, strike east and dip 10-25°N.

The depth of the orebody below the surface is 50 feet at the south end of the property and 2000 feet at the north boundary. The uraniferous conglomerates are confined to the basal 200 feet of the Matinenda Formation of the Elliot Lake Group.

Economic Features

Approximately 47,000 feet of diamond drilling in 109 holes have been completed in the upper portion of the orebody which represents about 20 percent of the mine's potential. The orebody strikes N80E and has an average dip of 17°N. The average thickness is 9.5 feet.

The basal conglomerate reef is composed of poorly-sorted quartz pebbles set in an impure quartz matrix. Here grades up to 0.08 percent U_3O_8 and thicknesses up to 15 feet are found. The Nordic reef is the major reef. It consists of fairly well-rounded, well-sorted, densely-packed, quartz pebbles. Another, unnamed, reef occurs 20 feet above the Nordic reef. It has similar composition.

Proven reserves as of January, 1966, were 831,000 tons grading 0.109 percent U_3O_8 ; possible reserves were 10,108,000 tons grading 0.106 percent U_3O_8 .

History of Development

1954-1959: One hundred and nine surface holes were drilled, totalling 47,000 feet, spaced 250 feet along strike and 450 feet along dip.

The No. 1 shaft went down to 1331 feet with levels at 230, 330, 425, 515, 605, 712, 775, 894, 971, 1054, 1153, and 1230 feet. Nordic mill was designed to process 3400 ton of ore a day and milling commenced in January 1957. Total drifting was 51,897 feet and total cross-cutting 7608 feet. Rio Algom Limited completed the work.

1960-1968: In 1961 the Nordic shaft was deepened to 1780 feet to provide access to three new levels where grades were up to 0.12 percent U_3O_8 . By the end of 1961 a

small pilot plant was being readied to make nuclear grade ammonium diuranate. A thorium recovery unit with a capacity of 150 to 200 tons a year was constructed.

In 1964 the eastern and western limits of the mine were opened up by both surface and underground drilling. Surface drilling indicated that the Pardee reef contained economic reserves.

Production of yttrium concentrates from the waste liquors from the uranium circuit began.

In 1966 a 150 ton per year uranium refinery was completed. The Nordic mill capacity increased to 3700 tons per day.

In 1968 milling operations ceased.

In 1970 mining operations were being phased out.

Total drilling during this period consisted of 17 holes from the surface totalling 19,447 feet and 25 holes underground totalling 16,042 feet. Total underground development was 121,240 feet of drifts and 22,397 feet of crosscuts.

As of 1968 the total U_3O_8 recovered was 30,676,782 pounds from 13,181,780 tons of ore milled.

1981: The Nordic Mine contains substantial uranium resources which may be commercially recoverable.

References

- Airth and Olson (1958)
 Giblin et al. (1979)
 Griffith (1967, p.138-151)
 Robertson (1968a, p. 38-39; 1968b, p. 117-120; 1981)

Ontario Ministry of Natural Resources Files.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
 File No. 000100.

SILVERMAQUE OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Probably brannerite, uraninite, and monazite.

Location

Latitude 46°24'32"N, Longitude 82°40'10"W.
 Gunterman Township.
 Map Reference: ODM Map 2113.

Geology

The southern part of the property is underlain by the Upper Mississagi Formation

consisting of grey crossbedded feldspathic quartzite. The central part is underlain by a Nordic diabase sill-like intrusion. The northwest-striking Horne Lake fault crosses the southwest part of the property.

Economic Features

On the basis of drilling, Norsyncomaque Mining Limited estimated ore reserves at 2,500,000 tons grading 0.10 percent U_3O_8 or 2.2 pounds per ton on the east side of the property and 2,000,000 tons grading 1.08 pounds U_3O_8 per ton contained in the "F" reef.

History of Development

1955: Norsyncomaque Mining Limited carried out geological and geophysical surveys.

1965-1967: Silvermaque Mining Limited drilled on the western part of the property.

1976: This property was optioned to Long Lac Mining and Exploration Limited.

1978: Long Lac Mining and Exploration Limited carried out diamond drilling.

References

- Canadian Mines Handbook, 1979-1980.
 Robertson (1968b, p. 136)

STANLEIGH MINE (RIO ALGOM LIMITED)

Commodity

Uranium.

Radioactive Minerals

Brannerite, uraninite, and monazite.

Location

Latitude 46°24'35"N, Longitude 82°37'50"W.
 Gunterman Township.
 Map Reference: ODM Map 2113.

Geology

Surface exposures consist of sparse conglomerate and minor feldspathic quartzite of the Gowganda Formation, cut by northwest- and west-trending diabase dikes. Two beds of ore grade conglomerate, each about 10 feet thick, are separated by a quartzite bed 5 to 22 feet thick. These beds strike east and dip 8 to 10°N.

Economic Features

No. 1 shaft intersected ore at 3493 feet, and No. 2 shaft at 3639 feet. To the closing in 1960, 2,495,726 tons were milled, producing 4,889,744 pounds of U_3O_8 . All mining operations ceased November 30, 1960.

History of Development

1956: Two shafts were sunk to 3792 feet and 3650 feet, respectively. Construction of a 3000 ton per day mill began, and was completed in 1957.

1958: A total of 583,208 tons of ore were milled. In the final two quarters, 720,531 pounds of U_3O_8 were recovered from 372,648 tons of ore with an average millhead grade of 2.07 pounds per ton U_3O_8 .

1959: The No. 1 shaft was deepened to 3,846 feet. Total production was 949,119 tons of ore from which 1,939,099 pounds of U_3O_8 were recovered.

1960: Production was 293,311 tons of ore milled, at a grade of 2.11 pounds U_3O_8 per ton, from which 464,741 pounds of U_3O_8 were recovered.

Stanleigh Uranium Mining Corporation and Preston East Dome Mines Limited amalgamated to form Preston Mines Limited.

Production ceased in November, 1960.

1978: Rehabilitation proceeded at the mine and mill to be completed in 1983.

1980: Rio Algom Limited and Preston Mines Limited amalgamated.

References

Canadian Mines Handbook 1981-1982
Robertson (1968b, p. 111-113)

JOLLINEAU TOWNSHIP**GARDINER OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°52'10"N, Longitude 83°35'37"W.
Jollineau Township.
Map Reference: OGS Map 2419.

Geology
The Gardiner occurrence lies in the Superior Structural Province and is underlain chiefly by Early Precambrian granite and granite gneiss.

Economic Features
Assay results from 15 samples ranged from 0.01 to 2.78 percent U_3O_8 (radiometric equivalent) (Lang 1952).

References
Giblin et al. (1979)
Lang (1952, p. 128)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000036.

JOUBIN TOWNSHIP**ABETA OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°22'43"N, Longitude 82°33'50"W.
Joubin Township.
Map Reference: OGS Map 2419.

Geology
Drilling intersected several radioactive oligomictic conglomerate beds within the Matinenda Formation.

Economic Features
Assays of samples from two drill holes gave maximum grades of 0.09 percent U_3O_8 over 5 feet and 0.14 percent U_3O_8 over 4 feet.

History of Development
1953-1954: Fifteen drill holes totalling 4241 feet were drilled by the Mining Corporation of Canada.
1955: Two drill holes totalling 1250 feet were drilled by the Abeta Mining Corporation Limited.
1977: These claims were optioned by Long Lac Mineral Exploration Limited.
1978: The option was dropped.

References
Giblin et al. (1979)
Robertson (1961, p. 47; 1968a, p.29,30)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000081.

MILLIQUA PROSPECT (PARDEE REEF OF PARDEE ZONE)

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 46°22'54"N, Longitude 82°31'16"W.
Joubin Township.
Map Reference: ODM Map 2001.

Geology

This prospect is underlain by Middle Precambrian Huronian metasediments comprising the arkose and conglomerate of the Matinenda Formation of the Elliot Lake Group. The strike is N70E and the dip is 15 to 20°N.

Economic Features

Diamond drilling indicated the presence of 5,390,000 tons of ore averaging 0.07 percent U₃O₈ at a depth of 1455 feet. Assays from two drill holes averaged 0.45 percent U₃O₈ and 1.83 percent ThO₂ over 6.7 feet (Robertson 1961).

History of Development

1953: Airborne magnetometer and radiometric surveys were completed by Technical Mine Consultants Limited.
1954: Extensive drilling was done by Pardee Amalgamated Mines Limited. An adit was driven to explore the zone underground.
McIntyre Porcupine Mines Limited drilled 21 holes totalling 5664 feet.

References

Robertson (1961, p.50-51)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Joubin Township, Drill Log Report No. 14, Pardee Amalgamated Mines, 1954-1955.
Technical File No. 63.419, Technical Mine Consultants Limited, 1953.

NASCO OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°26'33"N, Longitude 82°32'02"W.
Joubin Township.
Map Reference: ODM Map 2001.

Geology

Three deep diamond drill holes intersected the Huronian sequence from the

Serpent Formation at the surface through the Espanola, Bruce, Mississagi, Pecors, Ramsay Lake and Matinenda Formations to the Early Precambrian basement.

Economic Features

One drill hole intersected several bands of quartz-pebble conglomerate over a thickness of 110 feet. The best assay was 0.06 percent U₃O₈ over 2 feet, and a wedged section showed 0.04 percent U₃O₈ over 5 feet. In one hole mineralization occurred between 3242 and 3420 feet with assays of 0.07 percent U₃O₈ over 2 feet (Robertson 1961).

History of Development

1955-1957: Three drill holes totalling 12,226 feet were drilled by Nasco Cobalt Silver Mines Limited.
1970: Airborne radiometric survey by Kerr-McGee Corporation.
1974: Airborne magnetometer survey by North American Nuclear Limited.

References

Robertson (1961, p.51, Map 2001; 1968a, p.31)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Joubin Township, Drill Log Report No. 18, Nasco Cobalt Silver Mines Limited, 1955.
Technical File No. 63.2802, Kerr-McGee Corporation, 1970.
Technical File No. 2.1783, North American Nuclear Limited, 1974.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File No. 000083.

PARDEE PROSPECT (PARDEE ZONE)**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°23'28"N, Longitude 82°32'10"W.
Joubin Township.
Map Reference: OGS Map 2419.

Geology

Radioactivity is associated with the Matinenda Formation of the Elliot Lake Group. The beds strike east and have a shallow northward dip. The main oligomictic, conglomerate layers average 10 feet in

thickness and lie 40 to 70 feet above the basement (Robertson 1961).

Economic Features

The main zone is the Pardee reef which extends over a basement high into the Nordic channel where locally it has been mined. The average grade from mine intersections gave 0.04 percent U_3O_8 . A number of holes also intersected a 9 foot thick basal conglomerate bed which pinches out 1000 feet down. A third bed lies some 80 to 100 feet above the basement, but is intermittent (Robertson 1961).

History of Development

1953: Airborne magnetometer and radiometric surveys were completed by Technical Mine Consultants Limited.
 1955: Drilling was carried out by Pardee Amalgamated Mines Limited.
 1961: Pardee Amalgamated Mines Limited amalgamated with Rio Algom Mines Limited.
 1965-1967: Rio Tinto Mines Limited drilled five holes totalling 14,703 feet.
 1975: Name changed to Rio Algom Limited.

References

Giblin et al. (1979)
 Robertson (1961; 1968a, p. 52)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto.

Joubin Township, Drill Log Report No. 14, Pardee Amalgamated Mines Limited, 1954-1955.
 Joubin Township, Drill Log Report No. 32 and 42, Rio Tinto Canadian Exploration Limited, 1965-1967.
 Technical File No. 63.419, Technical Mine Consultants Limited, 1953

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
 File No. 000084.

PECORS LAKE (WEST) PROSPECT (PARDEE ZONE)

Commodity
 Uranium.

Radioactive Minerals
 Unknown.

Location
 Latitude $46^{\circ}23'16''N$, Longitude $82^{\circ}29'34''W$.
 Joubin Township.
 Map Reference: ODM Map 2001.

Geology

This prospect is underlain by Middle Precambrian Huronian metasediments comprising quartz-pebble conglomerate and arkose of the Matinenda Formation of the Elliot Lake Group. East-striking diabase dikes cross the area.

Economic Features

Ten drill holes spaced out over an area 1800 feet by 1100 feet indicated a grade of 0.05 percent U_3O_8 over a width of 4.9 feet.

Two holes were drilled from the northeast shore of Pecors Lake. The more easterly hole intersected seven radioactive oligomictic conglomerate bands. A 1.3 foot thick band at 1310.7 feet contained 0.04 percent U_3O_8 and 0.03 percent ThO_2 . A 7.5 foot band at 1338.5 feet contained 0.03 percent U_3O_8 and up to 0.04 percent ThO_2 (Robertson 1961).

History of Development

1953-1959: Airborne magnetometer and radiometric surveys were undertaken. Seventeen drill holes totalling 11,197 feet were drilled. Fifteen holes were drilled at the western end of Pecors Lake and two others on the northeast shore of Pecors Lake. Work was completed by Algom Uranium Mines Limited.
 1976: This property is now owned by Rio Algom Limited.

References

Robertson (1961, p. 47-48)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto.
 Technical File No. 63.441, Algom Uranium Mines Limited, 1953.

STANCAN OCCURRENCE

Commodity
 Uranium.

Radioactive Minerals
 Unknown.

Location
 Latitude $46^{\circ}24'14''N$, Longitude $82^{\circ}32'48''W$.
 Joubin Township.
 Map Reference: ODM Map 2001.

Geology

At the surface, the Gowganda Formation lies unconformably on the Serpent and Espanola Formations and all three are cut by Nipissing Diabase. At depth, drilling indicates

the presence of lowermost uraniferous member of the Mississagi Formation.

Economic Features

Assayed samples indicate a grade of 0.648 pounds U_3O_8 per ton over 34 feet, 1.13 pounds U_3O_8 per ton over 6.2 feet and 1.34 pounds U_3O_8 per ton over 5.1 feet (Mineral Deposit Inventory Record, File A0391, Ontario Geological Survey, Toronto).

History of Development

1955: One drill hole to 2717 feet was drilled by Stancan Uranium Corporation.

1966-1969: Twenty drill holes totalling 2165 feet were located between the southeast end of Elephant Lake and the northeast arm of Flying Goose Lake. Work was finished by Stanrock Uranium Mines Limited.

1968: Combined airborne magnetic and electromagnetic surveys were carried out by Stanrock Uranium Mines Limited.

1973: Stanrock Uranium Mines Limited amalgamated with Denison Mines Limited.

References

Robertson (1961, p. 58; 1968a, p.29-30)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Joubin Township, Drill Log Report No. 15, Stancan Uranium Mines Limited, 1955.

Joubin Township, Drill Log Report No. 39, Stanward Corporation and Stanrock Uranium Mines Limited, 1966-1969.

Technical File No. 63.2414, Stanrock Uranium Mines Limited, 1968.

Mineral Deposit Inventory Record, Ontario Geological Survey, Toronto
File A0391.

ST. MARY'S OCCURRENCE (PARDEE ZONE)

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude $46^{\circ}23'42''N$, Longitude $82^{\circ}34'01''W$.

Joubin Township.

Map Reference: ODM Map 2001.

Geology

A deep drill hole intersected the Huronian sequence from the Mississagi Formation at the surface through the Pecors, Ramsay Lake and Matinenda Formations. The rocks exposed were east-striking and north-dipping.

Economic Features

Assays from one drill hole gave values of 0.05 percent U_3O_8 over a width of 30 feet (Robertson 1961).

History of Development

1954-1955: One drill hole was drilled to a depth of 2105 feet. Work was finished by St. Mary's Uranium Mines Limited and New Jersey Zinc Exploration Company Canada Limited.

1975: St. Mary's Explorations Limited now owned by Rio Algom Limited.

References

Robertson (1961, p. 54; 1968a, p.29-30)

VITE OCCURRENCE (PECORS ZONE)

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude $46^{\circ}23'11''N$, Longitude $82^{\circ}27'14''W$.

Joubin Township.

Map Reference: ODM Map 2001.

Geology

This occurrence is underlain by Middle Precambrian Huronian metasediments comprising argillite and siltstone of the Pecors Formation of the Hough Lake Group and conglomerate of the Matinenda Formation of the Elliot Lake Group. The strike is N75W and dip is $30^{\circ}N$.

Economic Features

The two easternmost drill holes intersected thin beds of uraniferous conglomerate with assay values of 0.05 percent U_3O_8 over 3.7 feet and 0.04 percent U_3O_8 over 4.7 feet (Robertson 1968b).

History of Development

1951-1953: Exploration and drilling indicated a small nickeliferous zone located about 3600 feet southwest of the conglomerate intersections. Work was completed by Teck Corporation Company Limited.

1955-1956: Four drill holes were drilled

totalling 3518 feet by Vite Uranium Mines Limited.

1956: Airborne magnetic and electromagnetic surveys and one drill hole to 487 feet were completed by Kerr McGee Corporation.

References

Robertson (1961, p. 60; 1968a, p. 30-31)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Joubin Township, Drill Log Report No. 10, Teck Corporation Company Limited, 1941-1945.

Joubin Township, Drill Log Report No. 37, Kerr McGee Corporation Limited, 1967.

Technical File No. 63.2074, Kerr-McGee Corporation Limited, 1966.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000089.

KAMICHISITIT TOWNSHIP

KEE NO. 2 OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°28'01"N, Longitude 82°57'33"W.
Kamichisitit Township
Map reference: OGS Map 2419.

Geology

Outcrops consist of polymictic conglomerate of the Gowganda Formation of the Cobalt Group. One drill hole intersected three 60-foot thick conglomerate beds.

Economic Features

In the lowest section the conglomerate bed shows radioactivity at six times background. Concentrations average 0.05 U₃O₈.

History of Development

1969: One drill hole to 3712 feet was drilled by Cominco Limited.

1975: A 51 percent interest acquired by Imperial Oil Limited.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Kamichisitit Township, Drill Log Report No. 18, Cominco Limited, 1969.

Energy, Mines and Resources Canada Files Mineral Development Sector, Department of Energy, Mines and Resources, Ottawa.

National Inventory File U53, 41J/7, Kee #2, November, 1975.

KINCAID TOWNSHIP

HATHAWAY OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Pitchblende.

Location

Latitude 47°08'30"N, Longitude 84°43'40"W.
Kincaid Township.
Map Reference: OGS Map 2419.

Geology

The western third of the property is underlain by interbedded lava and sedimentary rocks. These are cut by Early Precambrian granite and pegmatite which underlie the greater part of the property. At least four sets of lower Keweenawan diabase dikes are found on the property. Radioactivity is associated with fractures in or near the diabase contact zones.

Economic Features

On No. 1 showing, the radioactive zone is 20 feet wide by 80 feet long. Grab samples assayed (chemical) 0.53, 0.67 and 0.66 percent U₃O₈. On No. 2 showing, samples assayed 0.421 and 0.60 percent (radiometric equivalent) (Assessment Files Research Office, File 63A.73, Ontario Geological Survey, Toronto).

History of Development

1949: Danaray Mines Limited completed geological mapping, stripping and trenching.

1951: A scintillometer survey was carried out.

1966: Hathaway Metal Mines Limited partially restaked the property.

References

Giblin et al. (1979)
Robertson (1968a, p. 22-23)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.
Technical File No. 63A.73, Danaray Mines Limited, 1949.

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.
Technical File No. 63.1801, Jayco Mines Limited, 1965.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File 000008.
Resident Geologist's Files, Sault Ste. Marie. Files SSM-68 and SSM-67, Abconore Uranium Mines, 1958.

KIRKWOOD TOWNSHIP**ABCONORE OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°20'50"N, Longitude 83°29'10"W.
Kirkwood Township.
Map Reference: OGS Map 2419.

Geology
The eastern half of claim 49979 is underlain by a Nipissing Diabase sill and the western half by sedimentary rocks of the Gowganda Formation consisting of boulder conglomerate and greywacke. A quartz-chalcopyrite vein strikes east and has been exposed in four pits over a length of 481 feet.

Economic Features
Radioactivity was recorded in greywacke in the westernmost pit and two grab samples averaged 0.095 percent U_3O_8 . A scintillometer survey showed radioactivity 16 times the background.

History of Development
1958: Pitting, geological and scintillometer surveys were carried out by Abconore Uranium Mines Limited.
1965: Ground magnetic, ground electromagnetic and geochemical soil surveys were completed by Jayco Mines Limited.
1966: Jayco Mines Limited amalgamated with Consolidated Silver Belle Mines Limited to form Silver Belle Mines Limited.
1969: Silver Belle Mines Limited merged with Alchib Development Limited.

References
Giblin et al. (1979)
Robertson (1968a, p. 6, 7)

LABELLE TOWNSHIP**FRANZ OCCURRENCE**

Commodity
Uranium, niobium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°23'00"N, Longitude 84°42'00"W.
Labelle Township.
Map Reference: OGS Map 2419.

Geology
Radioactive mineralization occurs at or near the north contact of a Keweenawan diabase dike cutting an Early Precambrian granite-gneiss complex. The Franz dike strikes N35W and dips steeply northeast. Pitchblende and uraninite are associated with a breccia zone caused by faulting along the north contact of the Franz dike and by cross-faulting in the diabase. The pitchblende is associated with calcite and secondary hematite and is believed to be of hydrothermal origin. Mineralization is also associated with fractures in the diabase and occurs in narrow zones up to 6 inches wide.

Economic Features
A channel sample taken over 3.5 feet across a fracture zone showed 0.80 percent U_3O_8 (radiometric equivalent). Another sample over 2.5 feet assayed 0.0002 percent U_3O_8 (radiometric equivalent).

History of Development
1951: Trenching and geological and scintillometer surveys were performed by Highland Prospecting Syndicate.
1975: The occurrence lies within Lake Superior Provincial Park.

References

Giblin et al. (1979)
Robertson (1968a, p. 26)
Lang (1952)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.
Technical File No. 63A.107, Highland
Prospecting Syndicate, 1950.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000074.

OTTAWA ASSOCIATES OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°22'00"N, Longitude 84°39'00"W.
Labelle Township.
Map Reference: OGS Map 2419.

Geology
Radioactive mineralization occurs in fractures and fault breccia along the contact of a Keweenaw diabase dike with Early Precambrian granite. Pitchblende mineralization was found in a shear zone 2 to 8 inches wide and exposed for a length of 184 feet. Fractures up to 10 feet long branch from the main shear zone into the diabase and granite.

Economic Features
Assays of eight grab samples averaged 3.97 percent U₃O₈ (Assessment Files Research Office, Ontario Geological Survey, Toronto, Technical File No. 63A.78).

History of Development
1949: A geological survey combined with geiger-counter exploration and some stripping was undertaken for the Ottawa Associates.
1956: Nine holes totalling 390 feet were drilled by the Ottawa Associates.
This occurrence is now situated within Lake Superior Provincial Park.

References
Giblin et al. (1979)
Robertson (1968a, p. 26-27)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File 63A.78, The Ottawa
Associates.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000075.

LABONTE TOWNSHIP**MCCOMBE SYNDICATE OCCURRENCE**

Commodity
Uranium, thorium.

Radioactive Minerals
Unknown.

Location
Latitude 47°21'30"N, Longitude 84°31'05"W.
Labonte Township.
Map Reference: OGS Map 2419.

Geology
The radioactive anomaly occurs in a brecciated Keweenaw diabase dike that cuts through Early Precambrian granitic gneiss. The dike trends northwest and dips steeply northeast. The dike, 4 feet wide, is cut by the Keenan fault, which caused brecciation over a length of about 300 feet and the entire width of the dike. The brecciated zone is highly radioactive with the footwall contact giving the highest readings.

Economic Features
A sample from this zone assayed 0.07 percent U₃O₈.

History of Development
1950: Geological and spot radiometric surveys by McCombe Syndicate.

References
Giblin et al. (1979)
Robertson (1968a, p. 53)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.
Technical File No. 63A.97, McCombe
Syndicate, 1950.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000056.

NAPRAY OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°22'30"N, Longitude 84°31'16"W.
Labonte Township.
Map Reference: OGS Map 2419.

Geology
The deposit occurs in a Keweenawan diabase dike about 100 feet wide, cutting Early Precambrian granitic gneiss and pegmatite. There are two showings. The Cranston showing consists of radioactive fractures striking N40W. The radioactive mineral is believed to be pitchblende. The discovery showing consists of a fracture zone striking N35W and dipping 75°E.

Economic Features
Two surface samples from the discovery showing assayed 0.03 percent U₃O₈. Two samples from the Cranston showing assayed 0.17 percent U₃O₈ (Robertson 1968a).

History of Development
1949: Prospecting, geological survey and reconnaissance geiger-counter survey and sampling was undertaken by Napray Mining Company Limited.

References
Giblin et al. (1979)
Robertson (1968a, p. 19)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.
Technical File No. 63A.72, Napray Mining Company Limited, 1949
Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File No. 000058.

LARSON TOWNSHIP**MCDONOUGH OCCURRENCE**

Commodity
Uranium, thorium.

Radioactive Minerals
Unknown.

Location
Latitude 47°22'00"N, Longitude 84°22'50"W
Larson Township
Map Reference: OGS Map 2419.

Geology
Radioactivity is associated with pegmatitic patches in granitic and migmatitic rocks. Narrow veins of dark grey quartz, commonly carrying abundant pyrite, cut some of the pegmatite zones and are also radioactive.

Economic Features
Analyses of samples collected by E.P. McDonough from trenches returned 0.5 to 4.4 pounds U₃O₈ per ton and 0.2 to 3.8 pounds ThO₂ per ton.

References
Giblin and Leahy (1979)
Giblin et al. (1979)

LONG TOWNSHIP**LOCATION X OCCURRENCE**

Commodity
Iron and uranium.

Radioactive Minerals
Pitchblende and thorite.

Location
Latitude 46°11'42"N, Longitude 82°46'43"W.
Long Township.
Map Reference: ODM Map 2186.

Geology
This occurrence is underlain by Middle Precambrian Huronian metasediments. A pyrite-specularite-quartz stockwork cuts interbedded quartzite and shale of the Gowganda Formation adjacent to a diabase dike.

Economic Features
Six samples from Location X were reported (Lang 1952) to contain an average of 0.08 percent U₃O₈. Two specimens submitted by W.C. Baycroft were found to contain pitchblende and thorite.

History of Development
1911: Location X was held by Lake Superior Mining Corporation Limited.
1949: Following the discovery by Aime Breton of a radioactive conglomerate sample in the Sault Ste. Marie main office of the mining recorder, Breton and Karl Gunterman discovered the source in Long Township.

The site was staked. Breton and Gunterman brought the discovery to the attention of Franc Joubin.

Ground radioactivity was not matched by assays of samples and the ground was allowed to lapse.

1952: Location X was staked by J.H. Hirshorn.

References

Lang (1952)

Lang et al. (1962, p. 128)

Robertson (1970a, p. 75)

PRONTO MINE (RIO ALGOM LIMITED)

Commodity
Uranium.

Radioactive Minerals

Brannerite, uraninite, monazite, thucholite, and uranophane.

Location

Latitude 46°12'38"N, Longitude 82°42'22"W.

Long Township.

Map Reference: ODM Map 2186.

Geology

The orebody is in a reef of quartz-pebble conglomerate at the base of the Matinenda Formation of the Elliot Lake Group. It strikes east and dips 15 to 20°S and unconformably overlies the Early Precambrian granite.

Several large diabase dikes cut the orebody and are affected locally by albite, chlorite, and carbonate alterations.

The typical ore is pyritic, uraniferous, quartz-pebble conglomerate, generally 6 to 10 feet thick and locally 15 to 20 feet thick. The thorium content here is the lowest of all Elliot Lake Mines. The orebody has a strike length of about 3500 feet and an average thickness of 7.5 feet.

Economic Features

As of 1960, a total of 2,264,404 tons of ore at a recovered grade of 2.05 pounds per ton had been milled, from which 4,643,835 pounds of U₃O₈ were recovered.

History of Development

1949: A. Breton discovered radioactivity. Karl Gunterman discovered radioactivity in rocks east of Lauzon Lake.

1952: Franc R. Joubin restaked the ground.

1953: Franc R. Joubin, financed by Hirshorn, headed diamond drilling exploration of this discovery. Pronto Uranium Mines Limited began development of a mine.

1954: A three-compartment, five-level shaft was completed to a depth of 592 feet. Construction of a mill and surface plant commenced.

1955: The mill was completed and rated at 1000 tons per day.

1957: The shaft was deepened and a seventh level was added. The plant was expanded to a mill rate of 1500 tons per day.

1960: The mine was closed. The mill was converted to treat copper from the Pater Mine.

Total diamond drilling consisted of 410 holes totalling 116,932 feet and 363 holes from underground totalling 18,089 feet. Two hundred feet of trenching on the surface was completed. Total underground development consisted of 24,421 feet of drifting and 10,168 feet of crosscutting.

References

Robertson (1968a, p. 8-9; 1970, p. 74-86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto.

Long Township, Drill Log Report No.

12. Pronto Uranium Mines Limited, 1953-1954.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000012.

MCAUGHEY TOWNSHIP

GIMBY OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 47°19'30"N, Longitude 84°11'50"W.

McAugey Township.

Map Reference: OGS Map 2419.

Geology

Radioactive mineralization occurs in a pegmatite-gneiss environment near the contact with mafic metavolcanics.

Economic Features

Two samples, one pegmatitic and the other gneissic containing pyrrhotite, assayed 0.063 and 0.081 percent U₃O₈ (radiometric equivalent).

History of Development

Circa 1950: Prospecting and sampling were done by J.E. Gimby.

1962: Airborne electromagnetic and magnetic surveys were carried out by the Algoma Central Railway.

References

Giblin et al. (1979)
Robertson (1968a, p. 52)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Technical File No. 63A.506, Group C,
Algoma Central Railway, 1962.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000040.

MORIN TOWNSHIP**MCINTYRE PORCUPINE OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°32'58"N, Longitude 83°37'38"W.
Morin Township.
Map Reference: ODM Map 2272.

Geology
Feldspathic sandstone and conglomerate of the Matinenda Formation of the Elliot Lake Group dip 25 to 45°S.

Economic Features
Quartz-pebble conglomerate beds, 2 to 3 feet thick, were intersected in drilling. Average assays were 0.016 percent U₃O₈ over 2.1 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto, Morin Township, Drill Log Report No. 10).

History of Development
1955: Four holes totalling 953 feet were drilled by McIntyre Porcupine Mines Limited.

References
Chandler (1973)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.
Morin Township, Drill Log Report No.

10. McIntyre Porcupine Mine Limited,
1955.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000070.

NICHOLAS TOWNSHIP**CRAZY LAKE OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°34'46"N, Longitude 82°53'49"W.
Nicholas Township.
Map Reference: OGS Map 2419.

Geology
Surface outcrops consist of amygdaloidal andesite-basalt and polymictic conglomerate of the Elliot Lake Group.

Economic Features
Drilling intersected a few quartz pebble conglomerate bands. Average assays were 0.023 percent U₃O₈ over 6.0 feet from bands that ranged in depth from 80.0 to 1719.7 feet below the surface.

History of Development
1967: Ground magnetometer survey was carried out by G.E. Parsons.
1968-1969: Geological and ground radiometric surveys and 15 drill holes totalling 9910 feet were performed by the Hanna Mining Company and Hecla Mining Company of Canada Limited.
1974: A multi-sensor airborne geophysical survey was carried out by Fort Norman Exploration Incorporated.

References
Giblin et al. (1979)
Leahy (1973)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 001110.

PEEVER TOWNSHIP

DAMASCUS OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 47°14'17"N, Longitude 84°35'26"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology
The radioactive anomaly occurs within the sheared contact zone of a Keweenawan diabase dike known as the Roche Dike.

Economic Features
The radioactivity was traced for 600 feet along the contact of the dike. One part of the zone showed 17 times background while the average was 2 to 4 times background. Four bulk samples assayed 0.007 percent U_3O_8 .

History of Development
1950: Prospecting and a detailed radioactive survey were carried out by Damascus Mines Limited.

References
Giblin et al. (1979)
Lang (1952)
Robertson (1968a, p. 18)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario Geological Survey, Toronto.
File No. 000046.

DOLAN OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°18'40"N, Longitude 84°36'20"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology
Pitchblende was found at the contact of a Keweenawan diabase dike cutting Early Precambrian granite.

Economic Features

The 17 samples from the property contained from 0.003 to 6.6 percent U_3O_8 (radiometric equivalent).

History of Development

1950: Prospecting and sampling by J.P. Dolan.

References

Giblin et al. (1979)
Lang (1952, p. 127)
Robertson (1968a, p. 18)

HENNESSY OCCURRENCE

Commodity
Uranium and thorium.

Radioactive Minerals
Unknown.

Location
Latitude 47°14'20"N, Longitude 84°34'50"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology

The radioactive samples are believed to have originated in pegmatite dikes within Early Precambrian granite.

Economic Features

One sample from a pegmatite dike assayed 0.36 percent U_3O_8 . Another assayed 2.4 percent ThO_2 and 0.02 percent U_3O_8 (Lang 1952).

History of Development

1950: Trenching and stripping was carried out by Hennessy Uranium Explorations Incorporated.

References

Giblin et al. (1979)
Lang (1952, p.128)

J.G. MCCOMBE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Ellsworthite (betafite) and allanite.

Location

Latitude 47°16'10"N, Longitude 84°31'40"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology

The deposit consists of five zones. Three occur at or near the contact of Keweenawan diabase dikes and Early Precambrian granite. The other two occur in granites and are believed to be pegmatites.

Economic Features

The average assay for 13 samples was 0.026 percent U_3O_8 .

History of Development

Circa 1950: Prospecting and sampling was done by J.G. McCombe.

References

Giblin et al. (1979)
Lang (1952, p. 130)

PATRICK OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Pitchblende.

Location

Latitude 47°13'46"N, Longitude 84°35'06"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology

The radioactive anomaly occurs within the sheared contact zone between a Keweenawan diabase dike and Early Precambrian granite. It is associated with pitchblende-bearing, calcite-hematite veins located within the sheared contacts and in the tension cracks in the diabase.

Economic Features

The average assay from a bulk sample was 0.101 percent U_3O_8 (chemical).

History of Development

1949-1951: Trenching and geiger-counter testing along with 24 drill holes totalling 2007 feet were carried out by Patrick Uranium Mines Limited.

References

Giblin et al. (1979)
Nuffield (1956)
Robertson (1968a, p. 16-17)

RANWICK PROSPECT**Commodity**

Uranium.

Radioactive Minerals

Pitchblende.

Location

Latitude 47°13'58"N, Longitude 84°36'00"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology

This occurrence is located at or near the contact of a Keweenawan diabase dike and Early Precambrian granitic gneiss. The dike, called the Ranson Dike, averages 70 feet wide, strikes northwest and dips 70° northeast. There are two other dikes, the Roche Dike to the north and the Canagau Dike to the south.

Pitchblende occurs in three sets of closely spaced fractures 5 to 10 feet apart parallel to and near the south (footwall) contact. The fractures are filled with calcite-hematite gangue.

Economic Features

Assays along 1049 feet of the adit indicated that pitchblende occurs in 10 zones. Assays from these zones averaged 0.034 percent U_3O_8 over 3.7 feet in width by 30.2 feet in length.

History of Development

1948: Discovered by R. Ranson.
1949-1951: Ranwick Uranium Mines Limited stripped, trenched, sampled, drove an adit to 1044 feet with two 70-foot crosscuts, and diamond-drilled nine holes.
1961: The adit was opened for public tours. A rock shop and mineral display was also opened.

References

Giblin et al. (1979)
Nuffield (1956, p. 25-26)
Robertson (1968a, p. 17)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000052.

VAN LAKE OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Pitchblende.

Location

Latitude 47°14'29"N, Longitude 84°35'32"W.
Peever Township.
Map Reference: OGS Map 2419.

Geology

Radioactive anomalies occur at the contact of a Keweenawan diabase dike and Early Precambrian granite. The dike strikes N40W, dips approximately 60°NE and has a width of 200 feet. The pitchblende mineralization is associated with calcite and hematite that fill the fractures in the sheared contact zone and cracks in the diabase and granite.

Economic Features

Two samples from two trenches gave chemical assays of 0.08 percent and 0.10 percent U₃O₈.

History of Development

1950-1951: A scintillometer survey and some trenching were done by Van Lake Prospecting Syndicate.

References

Giblin et al. (1979)
Lang (1952, p. 136)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Technical File No. 63.213, Van Lake
Prospecting Syndicate, 1950.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000073.

PICHE TOWNSHIP**GOLDEN ARROW OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°39'03"N, Longitude 82°33'08"W.
Piche Township.
Map Reference: ODM Map 2305.

Geology

In the area, uranium/thorium mineralization is associated with sedimentary rocks of the Lorrain Formation of the Cobalt Group. The sedimentary rocks strike locally east and dip gently to the south at 10 to 15°.

Economic Features

Purple bands of hematite or specularite in the sedimentary rocks are radioactive. Assays from samples of a 4-inch band of the purple band in conglomerate averaged 0.01 percent U₃O₈.

History of Development

In 1968, Consolidated Golden Arrow Mining Limited completed ground radiometric and geological surveys over this area.

References

Wood (1975, p. 56)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Technical File No. 63.2358,

Consolidated Golden Arrow Mining
Limited, 1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto.
File No. 000028.

INSPIRATION OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Monazite.

Location

Latitude 46°39'04"N, Longitude 82°33'11"W.
Piche Township.
Map Reference: ODM Map 2305.

Geology

Mineralization occurs in the interbedded quartzite and conglomerate of the Lorrain Formation of the Cobalt Group.

Economic Features

Drilling intersected bands of radioactive conglomerate. The average assay was 0.01 percent U₃O₈ (radiometric equivalent) over 29.6 feet.

History of Development

1953: Six drill holes totalling 1822 feet, were drilled between the southeast shore of Rawhide Lake and the southwest shore of

Rosemarie Lake. Work was done by A. and W. Hanson.
1968: Geological survey was carried out by Weston & Company Incorporated.

References

Wood (1975, p. 58)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Piche Township, Drill Log Report No. 11, W. Hanson, 1953.

Technical File No. 63A.545, Weston and Company Incorporated, 1968.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 000029.

1957: Cobalt Consolidated was renamed Agnico Mines Limited.

1972: Agnico Mines Limited amalgamated to form Agnico-Eagle Mines Limited.

References

Robertson (1968a, p. 13, 14; 1970)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto.

Poulin Township, Drill Log Report No. 11, Cobalt Consolidated Mining Corporation, 1955.

Poulin Township, Drill Log Report No. 13, E.E. Campbell, 1954.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto.

File No. 001121.

POULIN TOWNSHIP

COBALT CONSOLIDATED OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°38'14"N, Longitude 82°54'08"W.

Poulin Township.

Map Reference: ODM Map 2346.

Geology

Surface exposures consist of ferruginous slightly radioactive quartzite of the Lorrain Formation of the Cobalt Group. The beds strike east and dip 10 to 15°S.

Economic Features

One drill hole intersected radioactive quartz-pebble conglomerate which give 0.023 percent (radiometric equivalent) U_3O_8 over 1.5 feet. Assays of samples from another drill hole located about 660 feet north of the reference point, were 0.0006 percent U_3O_8 (radiometric equivalent) over 1.3 feet.

(Assessment Files Research Office, Ontario Geological Survey, Toronto, Poulin Township, Drill Log Reports No. 11 and 13).

History of Development

1954: One hole to 72 feet was drilled by E.E. Campbell.

1955: One hole was drilled to 1880 feet by Cobalt Consolidated Mining Corporation Limited.

REILLY TOWNSHIP

RANGER OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°57'00"N, Longitude 83°29'30"W.

Reilly Township,

Map Reference: OGS Map 2419.

Geology

The showing is in a diabase dike (location unknown) about 60 feet wide where an aplite dike 8 inches in width extends diagonally across it for a length of 30 feet. The central part of the aplite dike has a 2- to 3-inch wide radioactive zone of carbonate and quartz which has given anomalies up to five times background (Lang 1952).

Economic Features

A selected sample taken from the most radioactive part of the dike showed 0.30 percent U_3O_8 (radiometric equivalent).

History of Development

Pre-1953: Prospecting was carried out by H. Evans.

1953-1954: Trenching, aeroradiometric survey, and six inclined drill holes totalling 2240 feet were completed by Ranger Lake Uranium Mining Company Limited.

1956: Ranger Lake Uranium Mining Company Limited merged into Century Mining and Development Corporation.

References

Giblin et al. (1979)
Lang (1952, p. 123-124)
Robertson (1968a, p. 52)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Sault Ste. Marie.
File SSM 228.

Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.649, Ranger
Lake Uranium Mining Company
Limited, 1954.

Reilly Township, Drill Log Reports No.
10 and 11, Ranger Lake Uranium
Mining Company Limited, 1954.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000039.

RIX TOWNSHIP**CANAGAU DIKE OCCURRENCE**

Commodity
Uranium

Radioactive Minerals
Pitchblende.

Location
Latitude 47°13'52"N, Longitude 84°36'58"W.
Rix Township.
Map Reference: OGS Map 2419.

Geology
Radioactive anomalies occur at the sheared contact of several Keweenaw diabase dike and Early Precambrian granite. The Canagau dike strikes northwest and dips steeply north. Patches of pitchblende up to 1/4 inch wide occur in fractures filled with a calcite-hematite gangue.

References
Giblin et al. (1979)
Nuffield (1956, p.20)
Robertson (1968a, p.24)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Sault Ste. Marie.
File SSM 998.

ROOT TOWNSHIP**AUBREY FALLS OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°54'50"N, Longitude 83°12'40"W.
Root Township.
Map Reference: OGS Map 2419.

Geology
The radioactive minerals occur in rusty weathering gossan enclosed in small quartz veins, which cut red granite close to a diabase dike (Harding 1950).

Economic Features
Assays results from a zone 2 to 5 feet wide, traced over 800 feet averaged 0.10 percent U₃O₈ (Robertson 1968a).

History of Development
1949-1950: Preston East Dome Mines Limited did trenching, geological mapping, and a geiger counter survey; five inclined holes were drilled for a total of 799 feet.

References
Giblin et al. (1979)
Robertson (1968a, p. 14)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Root Township, Drill Log Report No. 10.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000037.

SAGARD TOWNSHIP**CANEONTI PROSPECT**

Commodity
Thorium, uranium.

Radioactive Minerals
Monazite.

Location
Latitude 46°38'56"N, Longitude 82°47'11"W.
Sagard Township.
Map Reference: ODM Map 2346.

Geology

At the west end of Tenfish Lake, a 35-foot thick quartz-pebble conglomerate bed striking easterly and dipping gently south is exposed for approximately 1000 feet.

Economic Features

Trenches revealed three quartz-pebble conglomerate beds over a width of 35 feet; the bottom 3 to 4 feet are the most strongly radioactive.

A radioactivity survey of the old trenches in the conglomerate bed indicated an average uranium content of 0.02 percent and average thorium content of 0.09 percent. A few thin hematitic conglomerate beds with radioactivity two to three times the background were intersected in drilling (Assessment Files Research Office, Ontario Geological Survey, Toronto, Sagard Township Drill Log Report No. 13). Four samples of a bulk sample from the trenches revealed 0.002 percent U_3O_8 and 0.52 percent ThO_2 .

History of Development

1955: Six trenches were excavated and two drill holes totalling 1164 feet were drilled by Caneonti Mines Limited.

Two drill holes totalling 363 feet, located near the southwest shore of Tenfish Lake, were drilled by Maralgo Mines Limited.

1968: Geological and radiometric surveys and deepening of old trenches were performed by Armore Mines Limited.

References

Giblin et al. (1979)
Robertson (1968a, p. 13; 1977a)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Sagard Township, Drill Log Report No. 13, Maralgo Mines Limited, 1955.
Technical File No. 63.2350, Armore Mines Limited, 1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 000032.

GREY TROUT LAKE PROSPECT

Commodity

Thorium and uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°39'06"N, Longitude 82°42'49"W.
Sagard Township.
Map Reference: OGS Map 2419.

Geology

A gamma-ray spectrometer survey indicated anomalous readings along an east-trending strike length of approximately 10,000 feet. The radioactivity appears to have a thorium source.

Economic Features

The radioactive quartz-pebble conglomerate beds were tested by eight drill holes. The zone has an indicated grade of 0.035 percent Th over 14.7 feet. Most beds contain less than 0.01 percent U_3O_8 .

Another set of six down-dip drill holes located approximately 1 mile southwest of the first set of holes and 4000 feet northeast of Blue Sky Lake, intersected hematite-bearing arkose quartz-pebble conglomerate at depths ranging from 10 to 800 feet. Drilling indicated a southerly-dipping (12°) conglomerate grit zone with an average thickness of 16.2 feet which contains approximately 0.02 percent ThO_2 and less than 0.01 percent U_3O_8 .

History of Development

1968: Geological, ground magnetic and radiometric surveys were carried out. Trenching and eight drill holes totalling 2130 feet were drilled. The work was completed by International Bibis Tin Mines Limited.

Six drill holes totalling 2731 feet and located about 1 mile southwest of the first set of holes were drilled by Silver Men Mines Limited.

One hole, drilled to 301 feet and located about 3100 feet southwest of the first set of holes, was drilled by E. Karpela.

1973: International Bibis Tin Mines Limited was renamed Laurasia Resources Limited.

References

Giblin et al. (1979)
Robertson (1970b; 1977a)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Sagard Township, Drill Log Report No. 11, E. Karpela, 1968.

Sagard Township, Drill Log Report No. 18, Silver Men Mines Limited, 1968.

Sagard Township, Drill Log Report No. 19, International Bibis Tin Mines Limited, 1968.

Technical File No. 63.2455,
International Bibis Tin Mines Limited,
1969.

HARVARD PROSPECT

Commodity
Uranium and copper.

Radioactive Minerals
Unknown.

Location
Latitude 46°38'08"N, Longitude 82°46'02"W.
Sagard Township.
Map Reference: ODM Map 2346.

Geology
Sedimentary rocks of the Lorrain Formation have been intruded by Nipissing diabase which contains numerous small quartz veins. Chalcopyrite, pyrite and uranium mineralization is associated with late stage alteration of granophyric sections of diabase and quartz veins. The principal alteration zones strike east, appear to dip vertically, have widths ranging from 10 to 100 feet and have strike lengths up to 600 feet.

Economic Features

Best assays average 0.14 percent U₃O₈ and 1.22 percent Cu over 6.0 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto, Sagard Township, Drill Log Report No. 17).

References

Robertson (1970b; 1977a)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto
Sagard Township, Drill Log Report No. 17, Harvard Uranium Mines Limited, 1955.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000033.

SHEDDEN TOWNSHIP

DENVIC LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 46°13'38"N, Longitude 82°19'22"W.
Shedden Township.
Map Reference: ODM Map 2314.

Geology

The Middle Precambrian Huronian metasediments strike N85E and dip 75°S, and they are bounded on the north by Early Precambrian granite. Minor radioactivity was associated with argillite of the Pecors Formation of the Hough Lake Group (Robertson 1977b).

Economic Features

A 10-foot wide shale band is strongly radioactive at the surface, and gave readings of 5 to 12 times the background at depth (Assessment Files Research Office, Ontario Geological Survey, Toronto, Shedden Township, Drill Log Record No. 12).

A few thin pyritic quartzite beds of Matinenda Formation were intersected in drilling. Assays averaged 0.01 percent U₃O₈ over 22 feet.

History of Development

1954: A geological survey and two drill holes totalling 401 feet were completed by Peach Uranium and Metal Mining Limited.

1955: Additional drilling on the western extension of the zone and eight trenches totalling 347 feet were carried out by Panel Consolidated Uranium Mines Limited.

References

Robertson (1977b, p.57)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto
Shedden Township, Drill Log Report No. 12, 1954.
Technical File No. 63A.179, Panel Consolidated Uranium Mines Limited.

SLATER TOWNSHIP

BATCHAWANA URANIUM MINES OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 47°11'04"N, Longitude 84°37'04"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology

Mineralization is located within a fault zone in one of eight Keweenawan diabase dikes cutting an Early Precambrian granite pegmatite complex. The fault zone is filled with quartz plus hematite and calcite in which a pitchblende vein is located.

The mineralization consists of two pods, one 15 by 10 by 1-1/2 inches and another located with the fault zone is exposed over a length of 150 feet.

History of Development

1949-1951: Geological, magnetometer and scintillometer surveys, nine holes totalling 355 feet, and some trenching were carried out by Batchawana Uranium Mines Limited.

References

Giblin et al. (1979)
Nuffield (1956, p.13)
Robertson (1968a, p. 20)

**Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto**

Technical File No. 63.172, Batchawana Uranium Mines Limited, 1950.

Technical File No. 63.261, Batchawana Uranium Mines Limited, 1951.

Drill Log Report, 1951, Batchawana Uranium Mines Limited.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000061.

Resident Geologist's Files, Sault Ste. Marie.
SSM Files 285, 286.

BOBCAM OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location

Latitude 47°11'00"N, Longitude 84°40'12"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology

Pitchblende occurs in narrow carbonate veinlets in fractures within Keweenawan diabase dikes. The veinlets are usually perpendicular to the contact with the country rock.

History of Development

1949: Geological, magnetometer and radiometric surveys, and some trenching were carried out by Bobcam Mines Limited.

References

Giblin et al. (1979)
Nuffield (1956, p.13)
Robertson (1968a, p.20)

**Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto**

Technical File No. 63.179, Bobcam Mines Limited, 1949.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000062.

CAMRAY OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location

Latitude 47°10'56"N, Longitude 84°41'22"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology

Pitchblende veins occur in tension cracks within Early Precambrian granite near its sheared contact with the footwall of a Keweenawan diabase dike. This dike strikes northwest and dips steeply north. (See "Theano Point Prospect", below.)

History of Development

1949: Radioactivity and geological surveys and some stripping were carried out. An adit 230 feet long was driven under the surface showing. All work was carried out by Camray Mines Limited.

References

Giblin et al. (1979)
Robertson (1968a, p. 21-22)

**Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto**

Technical File No. 63A.79, Camray Mines Limited, 1949.

Technical File No. 63.177, Camray Mines Limited, 1949.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000063.

THEANO POINT PROSPECT

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°10'58"N, Longitude 84°42'36"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology
Pitchblende veins occur in tension cracks within Early Precambrian granite at its contact with the footwall of the Keweenawan diabase dike. The dike strikes N80W and dips about 70°N.

The pitchblende is associated with calcite and hematite in veins usually 1 inch wide or less. Some stringers are 6 inches wide or more. The pitchblende-bearing fractures are concentrated in two zones. The west zone extends 245 feet east from the lake, and the east zone, 230 feet to the east, is 85 feet long. They both strike N80W and dip 70°N.

Economic Features
Samples from a shaft sunk 150 feet in 1949 averaged 0.24 percent U₃O₈ over 3 feet for a length of 52 feet. Along the drift, 90 feet from the shaft, average muck samples were 0.044 percent U₃O₈ and average back samples were 0.043 percent U₃O₈. East of the shaft and extending for 110 feet, average muck values were 0.02 percent U₃O₈ and average back samples were 0.037 percent U₃O₈ (Nuffield 1956, p.18-19).

History of Development
1847: Samples were collected in the area by B.A. Stanard and described by J.L. LeConte.
1949: The following work was done by Camray Mines Limited.

Radioactivity was monitored and geological surveys and trenching were done.

A shaft inclined at 69 degrees was sunk to 150 feet. A level was established at 138 feet on which 496 feet of drifting was done.

Diamond drilling consisted of 43 holes totalling 3086 feet from the surface and 29 holes totalling 1552 feet from underground.

1964: Camray Mines Limited's charter was cancelled.

References
Nuffield (1956)
Robertson (1968a, p. 21)
Giblin et al. (1979)

MOSHER-BYLES OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°08'33"N, Longitude 84°42'32"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology
Pitchblende occurs at the contact of a Keweenawan diabase dike with the Early Precambrian granitic country rock.

Economic Features
A sample from this occurrence assayed 0.42 percent U₃O₈ (chemical).

History of Development
Prior to 1952: Prospecting and sampling were done by A.C. Mosher and G. Byles.
1974: This property was restaked by Lucien Lacasse.

References
Giblin et al. (1979)
Lang (1952, p. 130)
Robertson (1968a, p. 53)

ROBB-MURMAC PROSPECT

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location
Latitude 47°12'26"N, Longitude 84°38'38"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology
The radiometric anomalies are located at or near the sheared contact between Keweenawan diabase dikes and Early Precambrian granite.

The "G" showing located in the south part of the property consists of a shear zone at the contact of the diabase and the granite. The shear zone, from a few inches to 5 feet in width, consists of thin veins and a fine network of fractures which are filled with hematite and calcite. Uranium bloom occurs on the surface. A sample from this showing registered 22.8 on the geiger counter.

The A-West showing consists of visible pitchblende exposed in a trench in the diabase about 3 feet from the granite contact. A geiger count of 8.1 was obtained over the showing.

The A-North showing consists of two zones where a geiger count reached 15. One zone is in the diabase approximately 15 feet from the south contact while the other zone is in the granite about 20 feet north of the diabase contact.

History of Development

1949: Geological and radiometric surveys, some stripping and trenching as well as 22 drill holes totalling 3322 feet were carried out by Murmac Lake Athabasca Mines Limited.

1976: A gamma ray spectrometer survey was carried out by Advance Murgor Explorations Limited.

References

Giblin et al. (1979)
Nuffield (1956, p.24)
Robertson (1968a, p. 23)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario
Geological Survey, Toronto
Drill Log Report, Murmac Lake
Athabasca Mines Limited, 1949.
Technical File No. 63.173, Robb-
Murmac Uranium Mines Limited, 1949.
Technical File No. 63A.85, J.P.
Arnott, 1949.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000007.

SOO-TOMIC OCCURRENCE

Commodity

Uranium and thorium.

Radioactive Minerals

Allanite and carnotite.

Location

Latitude 47°11'44"N, Longitude 84°36'21"W.
Slater Township.
Map Reference: OGS Map 2419.

Geology

Radioactivity occurs in a complex of granite, gneiss, and pegmatite of Early Precambrian age cut by northwest- and northeast-trending Keweenawan diabase dikes. Radioactivity was due to allanite. Some carnotite staining was also present.

Economic Features

Samples from oxidized sheared material gave assays of 0.02 and 0.04 percent U_3O_8 .

History of Development

1949: A geological survey and some sampling were done by Soo-Tomic Uranium Mines Limited.

References

Giblin et al. (1979)
Robertson (1968a, p.16)

Ontario Ministry of Natural Resources Files

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000045.

SMILSKY TOWNSHIP

FAUSTEN OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 47°13'29"N, Longitude 84°36'24"W.
Smilsky Township.
Map Reference: OGS Map 2419.

Geology

The Fausten occurrence consists of three radioactive fractures in Early Precambrian granite at the contact with a northwest-trending Keweenawan diabase dike.

Economic Features

Six samples from the deposit ranged from 0.026 to 0.20 percent U_3O_8 (radiometric equivalent) and four other samples showed 0.068 to 3.0 percent U_3O_8 (chemical).

History of Development

1949: Geological, geiger and geophysical surveys were completed by Ranrouyn Mines Limited.

1951: Geological and geophysical surveys completed by the Skaphe Group.

1952: Prospecting and sampling were done by Fausten Exploration Limited.

References

Giblin et al. (1979)
Robertson (1968a, p.52)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000042.

JAY-DEE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 47°13'20"N, Longitude 84°30'30"W.
Smilsky Township.
Map Reference: OGS Map 2419.

Geology

The radioactivity is associated with the pegmatite phase of an Early Precambrian granitic intrusion.

Economic Features

Drill core analyses ranged from 0.012 percent U_3O_8 over lengths ranging from 6 inches to 6 feet.

History of Development

1957: Geological and scintillometer surveys were completed. Fifteen holes were drilled totalling 3081 feet. Work was done by Jay-Dee Enterprises Incorporated.

References

Giblin et al. (1979)
Robertson (1968a, p.15)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Smilsky Township, Drill Log Report,
Jay-Dee Enterprises Incorporated,
1957.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000041.

LABINE-MCCARTHY OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende.

Location

Latitude 47°13'28"N, Longitude 84°35'06"W.
Smilsky Township.
Map Reference: OGS Map 2419.

Geology

Mineralization occurs near the south contact of a Keweenawan diabase dike with Early Precambrian granite. This dike strikes N80W and dips steeply to the north. The McCarthy fault cuts off the dike to the east.

The radioactivity is found mostly in a brecciated zone 12 to 15 inches wide within the diabase dike, parallel to and within several feet of the south contact. The fractures are filled with a calcite-hematite gangue and pitchblende. This zone gave readings of four to five times background.

History of Development

1949-1950: Prospecting and trenching was performed. A 6-mile road from the highway was built and a power transmission line was erected.

1951: An adit was driven with 1628 feet of drifting along the dike and the fault. Ten crosscuts totalling 192 feet and a 164-foot raise were driven. Eight holes totalling 1100 feet were drilled from underground. All work was performed by Labine-McCarthy Uranium Mines Limited.

1967: Some prospecting was done by Primrock Mining and Exploration Limited.

1969: Four holes were drilled, totalling 528 feet, by Primrock Mining and Exploration Limited.

References

Giblin et al. (1979)
Nuffield (1956)
Robertson (1968a, p. 15-16)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Smilsky Township, Drill Log Report
No. 10, Primrock Mining and
Exploration Limited, 1969.
Technical File No. 63A.546, Primrock
Mining and Exploration Limited, 1967.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000043.

Energy, Mines and Resources, Canada Files
Mineral Resources Branch, Department of
Energy, Mines and Resources, Ottawa, File
41 N/2, U1, McCarthy Dyke (Collins
Property), August 1969.

SPRAGGE TOWNSHIP

SPRAGGE CREEK OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°13'13"N, Longitude 82°41'07"W.
Spragge Township.
Map Reference: ODM Map 2186.

Geology
The Spragge Creek occurrence lies in the
Southern Structural Province and is underlain
by Middle Precambrian Huronian
metasediments.

Economic Features
A grab sample from a surface outcrop of
conglomerate at the fork in Spragge Creek,
approximately 1900 feet east of the township
boundary assayed 0.11 percent U₃O₈
(Robertson 1970a, p. 86).

History of Development
1953-1959: Airborne magnetic,
electromagnetic and radiometric surveys and
drilling were carried out by Pronto Uranium
Mines Limited.
1960: The claims became part of the Rio
Algom Mines Limited as the Pronto Division.
1975: Owner's name was changed to Rio
Algom Limited.

References
Robertson (1970a, p. 86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 63.419, Pronto
Uranium Mines Limited, 1953-1955.
Source Mineral Deposit Record, Ontario
Geological Survey, Toronto
File No. 000013.

TIMMERMANS TOWNSHIP

ANABAR OCCURRENCE

Commodity
Uranium.

Location
Latitude 46°22'46"N, Longitude 82°55'42"W.
Timmermans Township.
Map Reference: OGS Map 2419.

Geology
Drilling intersected a few thin pyritized
radioactive conglomerate bands from 0.4 to
3.5 feet thick.

Economic Features
The best assays were 0.05 percent U₃O₈
over 0.4 feet and 0.18 percent over 3.5 feet
(Robertson 1963, p. 56).

History of Development
1955: Fourteen drill holes totalling 5532 feet
were drilled by Anabar Mining and
Development Company Limited.
1967: Airborne geophysical and
magnetometer surveys were carried out by
Denison Mines Limited.

References
Giblin et al. (1979)
Robertson (1963, p.53, 55; 1968a, p. 48)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Timmermans Township, Drill Log
Report No. 20, Anabar Mining and
Development Company Limited, 1955.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000121.
Resident Geologist's Files, Sault Ste. Marie.
SSM File 423.

BIG GAME OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°23'21"N, Longitude 82°49'31"W.
Timmermans Township.
Map Reference: OGS Map 2419.

Geology

The Big Game occurrence lies in the Southern Structural Province and is underlain by Middle Precambrian Huronian metasediments of the Matinenda Formation. The area is crossed by the North and South Rossmere Faults.

Economic Features

A 2.5-foot thick bed of pyritiferous, radioactive, oligomictic conglomerate intersected in one drill hole gave an assay value of 0.019 percent U_3O_8 (Assessment Files Research Office, Ontario Geological Survey, Toronto, File No. 63A.303).

History of Development

1953: Airborne magnetometer and scintillometer surveys were carried out by Technical Mines Consultants Limited.
1954-1955: Geological, ground scintillometer surveys, soil sampling, and drilling of nine holes totalling 5434 feet were carried out by New Jersey Zinc Exploration Company (Canada) Limited. Drill hole BG-1 was completed jointly with Big Game Mines Limited to a depth of 917 feet.
1957: The claims came under the option of Consolidated Fredrick Mines Limited (controlled by Rio Tinto Group). This company allowed the claims to lapse.

References

Giblin et al. (1979)
Robertson (1968a, p. 54)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63A.303, New Jersey Zinc Exploration Company (Canada) Limited, 1954-1955.
Timmermans Townships, Drill Log Report No. 21, New Jersey Zinc Exploration Company (Canada) Limited, 1955.
Technical File No. 63.419, Technical Mine Consultants Limited, 1954.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000123.
Resident Geologist's Files, Sault Ste. Marie
SSM Files 393, 399.

BUFFANA PROSPECT

Commodity
Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°24'38"N, Longitude 82°49'52"W.
Timmermans Township.
Map Reference: OGS Map 2419.

Geology

The Matinenda Formation of the Elliot Lake Group outcrops east of Little Moon Lake. The Ramsay Lake and Pecors Formations of the Hough Lake Group outcrop at the north end of the lake. These sedimentary rocks lie unconformably on Early Precambrian granite which was intersected by drilling at depths ranging from 460 feet to 1025 feet.

Economic Features

A few thin radioactive quartz-pebble conglomerate beds with minor pyrite were intersected by drilling at 184.5 feet and 642.0 feet. The average assay was 0.048 percent U_3O_8 over 2.75 feet.

History of Development

1954-1955: Geological mapping was performed and six holes totalling 4867 feet were drilled by Rochester and Pittsburg Coal Company (Canada) Limited.
1974: Airborne magnetometer, electromagnetic, radiometric and resistivity surveys were carried out by Fort Norman Exploration Incorporated.
1975: Airborne magnetometer and electromagnetic surveys were done by the Fort Norman Group.

References

Giblin et al. (1979)
Robertson (1963, p.53,59; 1968a, p. 48)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Timmermans Township, Drill Log Report No. 26, Rochester & Pittsburg Coal Company (Canada) Limited, 1955.
Technical Report 2.1575, Fort Norman Exploration Incorporated, 1974.
Technical Report 2.1850, Fort Norman Group of Companies, 1975.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000124.
Resident Geologist's Files, Sault Ste. Marie
SSM File 431.

COFFEE LAKE OCCURRENCE**Commodity**

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°25'42"N, Longitude 82°56'32"W.
Timmermans Township.

Map Reference: OGS Map 2419.

Geology

Surface exposures consist of Mississagi Formation of the Hough Lake Group. At depth, rock types belonging to Pecors, McKim, and Matinenda Formations lie unconformably on Early Precambrian granite. The basement was reached at 3193 feet in a drill hole. In this drill hole, two thin sections of pyritiferous, uraniferous, oligomictic conglomerate were intersected at depths from 2963.5 to 2996.5. The average assay was 0.054 percent U₃O₈ and 0.092 percent ThO₂ percent over 1.0 foot.

History of Development

1967: Combined airborne magnetometer and electromagnetic surveys were carried out by Denison Mines Limited.

1969: Drill hole DM 69-3 was drilled to 3212 feet by Denison Mines Limited.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log
Report No. 13, Denison Mines Limited,
1969.

Technical File No. 63.2201, Denison
Mines Limited, 1967.

Technical File No. 63.2264, Denison
Mines Limited, 1967.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001112.

DENISON OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°24'58"N, Longitude 82°51'22"W.
Timmermans Township.

Map Reference: ODM Map 2026.

Geology

Surface exposures consist of rock types belonging to the Ramsay Lake Formation of the Hough Lake Group. These sedimentary rocks strike east and dip at 10 to 25°N.

Economic Features

Drill hole DM-68-1 intersected pyritiferous, radioactive, oligomictic conglomerate from 820 to 916 feet and 0.036 percent ThO₂ over 1.5 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto, Timmermans Township, Drill Log Report No. 10, 1968).

History of Development

1954: Airborne scintillometer survey was done by Sapphire Petroleum Limited.

1967-1968: Combined airborne magnetometer and electromagnetic surveys and drilling of one hole to 1004 feet were carried out by Denison Mines Limited.

References

Leahy (1973)

Robertson (1963)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log
Report No. 10, Denison Mines Limited,
1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001120.

DOMINION PROSPECT**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°23'00"N, Longitude 82°53'47"W.
Timmermans Township.

Map Reference: ODM Map 2026.

Geology

Fourteen holes, drilled on the south shore of Bakers Bay, southeast of Graveyard Island, intersected occasional thin oligomictic

pebble bands. The average assay was 0.016 percent U_3O_8 over 7.0 feet

History of Development

1954-1955: Geological survey and 14 drill holes totalling 5709 feet were completed by Dominion Uranium Corporation.

1974: Airborne magnetometer, electromagnetic, radiometric and resistivity surveys on former claim SSM 32628 were carried out by Fort Norman Explorations Limited.

References

Robertson (1963; 1968a, p. 49)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log Report No. 32, Dominion Uranium Corporation, 1955.

Technical Report No. 2.1575, Fort Norman Explorations Incorporated, 1974.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000125.

JEANETTE PROSPECT

Commodity

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°24'46"N, Longitude 82°51'40"W.
Timmermans Township.

Map Reference: ODM Map 2026.

Geology

The Jeanette prospect lies in the Superior Structural Province and is underlain by the Matinenda Formation of the Elliot Lake Group. These sedimentary rocks strike east and dip gently to the north.

Economic Features

Two drill holes intersected pyritiferous, radioactive, oligomictic conglomerate with average assays of 0.029 percent U_3O_8 and 0.02 percent ThO_2 over 28 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto, Timmermans Township, Drill Log Report No. 24).

History of Development

1953: 14 pits over a distance of 1 mile were dug by Moneta Mining Company Limited.

1954: Geological and airborne scintillometer surveys and five drill holes were completed by Sapphire Petroleums Limited.

1955: Five holes totalling 2406 feet were drilled by Jeanette Minerals Limited in conjunction with Sapphire Petroleums Limited.

1967: Combined airborne magnetometer and electromagnetic surveys were completed by Denison Mines Limited.

References

Robertson (1963, p. 66-67)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log Reports No. 24 and 25, Sapphire Petroleums Limited, 1954-1955.

Technical File No. 63.466, Sapphire Petroleums Limited, 1954.

Technical File No. 63.2201, Denison Mines Limited, 1967.

Technical File No. 63.2264, Denison Mines Limited, 1967.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000127.

MARTIN OCCURRENCE

Commodity

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°23'18"N, Longitude 82°51'58"W.
Timmermans Township.

Map Reference: ODM Map 2026.

Geology

The Martin occurrence is underlain by feldspathic quartzite, arkose and conglomerate of the Matinenda Formation of the Elliot Lake Group which lies unconformably on Early Precambrian granite. Two drill holes intersected the basement at 477 and 408 feet respectively.

Economic Features

Two drill holes intersected thin radioactive conglomerate beds with sulphide mineralization at depths ranging from 125.6

to 565.3 feet. Average assays were 0.021 percent U_3O_8 and 0.040 percent ThO_2 over 1.2 feet.

History of Development

1955: Two holes totalling 1111 feet were drilled by Darwin R. Martin.

1974: Airborne magnetometer, electromagnetic radiometric, and resistivity surveys were carried out by Fort Norman Explorations Incorporated.

References

Robertson (1963, p. 49-50)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Timmermans Township, Drill Log Report No. 41, Darwin R. Martin Property, 1955.

Technical File No. 2.1575, Fort Norman Explorations Incorporated, 1974.

1957: New Jersey Zinc Exploration Company (Canada) Limited merged with others in the Rio Tinto Group to form Consolidated Fredrick Mines Limited.

References

Robertson (1963, p. 70)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Timmermans Township, Drill Log Report No. 22, New Jersey Zinc Exploration Company (Canada) Limited, 1954-1955.

Technical File No. 63A.303, New Jersey Zinc Exploration Company (Canada) Limited, 1954-1955.

Technical File No. 2.1575, Fort Norman Explorations Incorporated, 1974.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 001113.

MOON LAKE OCCURRENCE

Commodity

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude $46^{\circ}23'10''N$, Longitude $82^{\circ}50'30''W$.

Timmermans Township.

Map Reference: ODM Map 2026.

Geology

The area lying along the north shore of Baker's Bay is underlain by Matinenda Formation quartzite and arkose.

Economic Features

Twenty-four holes were drilled in an area stretching along the south shore of Matinenda Lake, between the south end of Little Moon Lake and the west end of Baker's Bay and in the vicinity of Caribou Lake.

Two thin sections of pyritiferous uraniferous oligomictic conglomerate were intersected at 293.8 and 556.5 feet. Average assays were 0.020 percent U_3O_8 and 0.040 percent ThO_2 over 1.9 feet.

History of Development

1954-1955: Geological, ground scintillometer and soil surveys and 24 drill holes were all completed by New Jersey Zinc Exploration Company (Canada) Limited.

PICTON OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude $46^{\circ}26'20''N$, Longitude $52^{\circ}55'05''W$.

Timmermans Township.

Map Reference: ODM Map 2026.

Geology

At the surface, Mississagi quartzite of the Hough Lake Group is well exposed, striking east and dipping gently north.

At depth, Matinenda arkose and quartzite lie unconformably on "Keewatin greenstone" which was intersected at 2309 feet in hole No. 1. Radioactive pyritic conglomerate pebble bands were intersected in the same hole at 2059, 2070, and 2115 feet for an average thickness of 3.5 feet.

Economic Features

At depths of 1049.5 to 1052 feet, chalcocopyrite, pyrite and pyrrhotite mineralization assayed 1.2 percent copper over 2.5 feet; trace amounts of gold and cobalt were also indicated.

History of Development

1954-55: Drill hole p-1 went to 2309 feet. The drilling was done jointly by Picton

Uranium Mines Limited and Brunette Porcupine Gold Mines Limited.

Drill holes p-2 and p-3, totalling 1444 feet and located 330 feet and 1650 feet northwest of p-1 respectively, were completed by Picton Uranium Mines Limited.

1967: Claims owned by Denison Mines Limited.

References

Robertson (1963, p. 72-73)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical Report No. 63.2264, Denison Mines Limited, 1967

Drill Log Report No. 38, Picton Uranium Mines Limited, 1955

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000129

PISTOL LAKE PROSPECT

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°22'00"N, Longitude 82°51'22"W.
Timmermans Township.
Map Reference: ODM Map 2026.

Geology

Minor radioactive, quartz-pebble conglomerate bands of the Matinenda Formation of the Elliot Lake Group were intersected in three drill holes.

Economic Features

Average assay results were 0.026 percent U₃O₈ over 1.0 feet. The radioactive beds were intersected at 139.0, 464.5, and 488.0 feet (Assessment Files Research Office, Ontario Geological Survey, Toronto, Timmermans Townships, Drill Log Report No. 19).

History of Development

1954-1958: Geological and geophysical surveys and 13 drill holes totalling 4682 feet were completed by Algom Uranium Mines Limited.

1958: The claims were allowed to lapse.

References

Robertson (1963, p. 54)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Drill Log Report No. 19, Algom Uranium Mines Limited, 1954.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000130.

ZENMAC OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°24'00"N, Longitude 82°53'10"W.
Timmermans Township.
Map Reference: ODM Map 2026.

Geology

The central portion of the area is underlain by massive dioritic gabbro at the east end of the Matinenda sill-like intrusion. The southern part shows well-bedded Matinenda quartzite dipping at about 10°S. The northern part is made up of water claims.

Three drill holes, located on the north shore of Baker's Bay, intersected radioactive oligomictic conglomerate beds at 467.5, 483.5 and 506.0 feet. The beds average 6.3 feet in width and 5.1 times the background radioactivity.

History of Development

1954: Two drill holes totalling 1153 feet were drilled by Zenmac Metal Mines Limited.

1955: One drill hole, 752 feet in depth and located about 3000 feet west of Zenmac hole II-2, was completed by Stancan Uranium Corporation. Stanrock Uranium Mines Limited later acquired Stancan Uranium Corporation.

1973: Stanrock Uranium Mines Limited amalgamated with Denison Mines Limited.

References

Robertson (1963, p. 78)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log

Report No. 27, Zenmac Metal Mines Limited, 1954.

Timmermans Township, Drill Log Report No. 30, Stancan Uranium Corporation, 1955.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000131.

TUPPER TOWNSHIP

PITCHORE PROSPECT

Commodity

Uranium, copper.

Radioactive Minerals

Unknown.

Location

Latitude 46°52'30"N, Longitude 84°13'30"W.
Tupper and Shields Townships.
Map Reference: OGS Map 2419.

Geology

Uranium mineralization is associated with quartz veins containing chalcopyrite and traces of pyrite, hematite and carbonate. The veins occur in diabase and granite. The mineralized zone extends over a strike length of 460 feet.

Economic Features

Assays from drill core samples averaged 0.076 percent U_3O_8 over 5.7 feet (Shklanka 1969, p. 69).

History of Development

1954-1955: Trenching, magnetic and scintillometer surveys, and 25 drill holes totalling 6133 feet were completed by Pitchore Uranium Mines Limited.

References

Giblin et al. (1979)
Robertson (1968a, p. 51)
Shklanka (1969, p.69)
Financial Post Survey of Mines (1956, p.319; 1957, p.339)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63.666, Pitchore Uranium Mines Limited, 1955.

VANKOUGHNET TOWNSHIP

VANKOUGHNET OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°46'00"N, Longitude 84°47'45"W.
VanKoughnet Township.
Map Reference: OGS Map 2419.

Geology

The radioactive zones consist of narrow limonite-stained fractures which contain quartz and feldspar, and tiny breccia fragments of quartzite. They range in thickness from 5 to 15 cm and can be traced along strike for about 6 m. Three parallel zones occur on a steep hillside across a horizontal distance of about 12 m. The zones strike N70W and dip 30 to 35°N.

Economic Features

Three samples collected from the widest zone were analysed and found to contain 40, 67 and 87 ppm U_3O_8 with less than 10 ppm ThO_2 in each.

References

Giblin and Leahy (1979, p.89-99)
Giblin et al. (1979)

VIEL TOWNSHIP

GAITWIN OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°40'00"N, Longitude 82°40'32"W.
Viel Township.
Map Reference: OGS Map 2305.

Geology

Sedimentary rocks of the Lorrain Formation of the Cobalt Group strike locally northeast and dip 45°SE. Uranium mineralization occurs in ferruginous feldspathic conglomerate occurring in a bed approximately 5 feet wide and extending 1100 feet.

Economic Features

Geiger counter readings showed radioactivity up to five times background along this bed (Assessment Files Research Office, Ontario Geological Survey, Toronto, Technical File No. 63A.171).

History of Development

1953-1955: A geological survey and one drill hole to 900 feet were completed by Gaitwin Explorations Limited.

1969: Combined airborne magnetic and electromagnetic surveys were carried out by A. St Denis.

1976: This occurrence was open for staking.

References

Wood (1975, p. 58)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Viel Township, Drill Log Report No. 12, 1955.

Technical File No. 63A.171, Gaitwin Explorations Limited, 1953.

Technical File No 63.2477, A. St Denis, 1969.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000030.

MATTAINI OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°38'53"N, Longitude 82°37'58"W.
Viel Township.

Map Reference: ODM Map 2305.

Geology

The Mattaini occurrence lies in the Southern Structural Province and is underlain by Middle Precambrian Huronian metasediments. A differentiated Nipissing Diabase sill intrudes interbedded quartzite and hematitic radioactive conglomerate of the Lorrain Formation (Robertson 1968a). Granodiorite zones in the diabase contain scattered chalcopyrite, pyrite, bornite, and specularite. Some of the zones intersected in drilling are radioactive up to 15 times the background.

Economic Features

Assays from 10 radioactive zones, averaging 2.5 feet thick, were 0.09 percent U_3O_8 (radiometric equivalent) (Assessment Files Research Office, Ontario Geological Survey, Toronto, Viel Township, Drill Log Report No. 13).

History of Development

1954: An airborne radiometric survey was carried out by Carl and Blanche Mattaini.

1955: Eight pits, four inclined drill holes totalling 2048 feet, and two vertical drill holes totalling 2480 feet were completed by Belfast Mines Limited.

References

Robertson (1968a, p. 12)

Wood (1975, p. 56, 58)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Viel Township, Drill Log Report No. 13, Belfast Mines Limited, 1955.

RAWHIDE OCCURRENCE**Commodity**

Thorium and uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°38'18"N, Longitude 82°38'05"W.
Viel Township.

Map Reference: ODM Map 2305.

Geology

Sedimentary rocks of the Lorrain Formation of the Cobalt Group have been intruded by Nipissing Diabase. Two areas of mineralization, 200 to 250 feet apart, were located near the western boundary of former claim S150104. The quartzite in this area has been hematitized near the contact with a diabase sill.

Economic Features

Drilling intersected feldspathized diabase. Assays of samples taken here were 0.02 percent U_3O_8 over 1.4 feet and 0.02 percent ThO_2 over 5 feet (Energy, Mines and Resources Canada, Ottawa, File U2, 41 J/10, February, 1969).

History of Development

1954: Airborne radiometric survey by
Aerosint Exploration Syndicate.

1968: Geological, ground radiometric surveys,
and four drill holes totalling 1,002 feet were
completed by Rawhide "U" Mines Limited.

References

Wood (1975, p. 55, 58)

Ontario Ministry of Natural Resources Files

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001114.

Energy, Mines and Resources Canada Files

Mineral Development Sector, Department of
Energy, Mines and Resources, Ottawa
National Mineral Inventory File U2, 41
J/10, Rawhide "U" Mines Limited,
February, 1969).

MINOR URANIUM AND THORIUM OCCURRENCES OF ALGOMA DISTRICT

ALBANEL TOWNSHIP

ARCO TRILLER NO. 2

Location

Latitude 46°33'54"N, Longitude 82°57'50"W
Albanel Township.

Remarks

Drill hole No. 2 by Arco Triller intersected the Huronian Espanola, Bruce, and Mississagi Formations, and Early Precambrian basement at 4082 feet. Sections between 3010 and 3012 feet assayed 0.03 percent U₃O₈ (Siemiakowska et al. 1975).

In 1968 a ground magnetometer survey and two drill holes totalling 4577 feet were completed by Arco Triller Explorations Limited.

In 1974 a multi-sensor airborne geophysical survey was completed by Fort Norman Explorations Incorporated.

References

Giblin et al. (1979)
Leahy (1973)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2213, Arco Triller Explorations Limited, 1967.
Albanel Township, Drill Log Report No. 16, Atlantic Richfield Company, 1968.

Technical File No. 2.1575, Fort Norman Explorations Limited, 1974.

BOLGER TOWNSHIP

PEERLESS URANIUM MINING CORPORATION LIMITED (EXPLORATION AREA)

Location

Latitude 46°22'43"N, Longitude 82°47'16"W.
Bolger Township.

Remarks

Two northwest-striking faults, one the southeast continuation of the Moon Lake fault, and a north-striking fault, lying close to the east boundary, cross the area. Drilling indicated a maximum of 135 feet of arkose considered to be the equivalent of the Matinenda Formation of the Elliot Lake

Group. A number of thin oligomictic conglomerate bands were intersected but were only slightly radioactive.

References

Canadian Mines Handbook (1968-1969, p. 268)
Robertson (1963, p. 72)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.244, Peerless Uranium Mining Corporation Limited.
Bolger Township, Drill Log Report No. 12, Peerless Uranium Mining Corporation Limited, 1955.

DAY TOWNSHIP

SOWERBY OCCURRENCE

Location

Latitude 46°18'30"N, Longitude 83°23'15"W.
Day Township.

Remarks

Spotty mineralization occurs in Nipissing Diabase and the Gowganda Formation adjacent to the contact or as brecciated inclusions in the diabase. The mineralization includes Cu, Co, Ni, and U. Radioactive minerals include pitchblende and uranophane.

References

Giblin et al. (1979)
Robertson (1968a, p. 4)

DUNCAN TOWNSHIP

DUNCAN TOWNSHIP OCCURRENCE

Location

Latitude 46°37'30"N, Longitude 84°07'00"W.
Duncan Township.

Remarks

The Duncan Township uranium occurrence is situated in the quartz-pebble conglomerate of the Matinenda Formation of the Elliot Lake Group.

References

Frarey (1977)
Giblin et al. (1979)

MAUD LAKE OCCURRENCE**Location**

Latitude 46°36'30"N, Longitude 84°08'50"W.
Duncan Township.

Remarks

There are several showings of uraniferous quartz-pebble conglomerate along the east shore of Maud Lake. This unit consists of conglomerate, arkose and greywacke. The individual beds are no thicker than 2 feet and total thickness is 15 feet.

References

Giblin et al. (1979)
Frarey (1977)
Robertson (1968a, p. 5)

GAIASHK TOWNSHIP**CANUC OCCURRENCE****Location**

Latitude 46°26'04"N, Longitude 82°20'32"W.
Gaiashk Township.

Remarks

The Precambrian country rock consists mainly of Huronian sedimentary rocks of the Quirke syncline and Keweenawan diabase.

In 1954, assays of drill core from 105.5 to 106.7 feet averaged 0.01 percent U_3O_8 . Assays of core from 527 to 527.5 feet were 0.017 percent U_3O_8 .

Between 1954 and 1968, one hole was drilled to 574 feet and numerous geophysical surveys were carried out.

References

Giblin et al. (1979)
Robertson (1962, p. 71)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Gaiashk Township, Drill Log Report
No. 17, British Columbian Explorers,
1954.

Technical File No. 63.419, Pronto
Uranium Mines Limited, 1957.

Technical File No. 63.2286, Canuc
Mines Limited, 1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001132.

Energy, Mines and Resources Canada Files
Mineral Development Sector, Department of
Energy, Mines and Resources, Ottawa
National Inventory File U2, 41, J/8,
Anticline Point, September 1962.

GAUDETTE TOWNSHIP**ALUR OCCURRENCE****Location**

Latitude 46°49'10"N, Longitude 83°59'00"W.
Gaudette Township.

Remarks

There are several fractured Keweenawan diabase dikes in the area that are marked by the presence of iron carbonate. One dike on the southeast corner of claim AC4121 is crossed by a carbonated shear which is radioactive. A sample assayed 0.07 percent U_3O_8 (radiometric equivalent). Prospecting and geologic mapping were carried out by Alur Mines Limited in 1949.

References

Giblin et al. (1979)
Lang (1952, p. 135)
Financial Post Survey of Mines (1953, p.225)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.75, 1949.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000003.

GUNTERMAN TOWNSHIP**INTERNATIONAL OCCURRENCE****Location**

Latitude 46°22'50"N, Longitude 82°39'43"W.
Gunterman Township.

Remarks

One outcrop of Lower Mississagi Formation arkose with oligomictic pebble bands extends into the area from Rio Algom Limited's Nordic claim group. The arkose assayed 0.02 and 0.104 percent U_3O_8 and the pebble bands (6 inches to 1 foot thick) 0.003 and 0.07 percent U_3O_8 .

References

Prasad (1981, p. 74)
Robertson (1968b, p. 132)

**THORNCREST RADIOACTIVITY
OCCURRENCE****Location**

Latitude 46°25'21"N, Longitude 82°40'27"W.
Gunterman Township.

Remarks

Radioactivity was found in the conglomerate of the Gowganda Formation.

References

Prasad (1981, p. 73)
Robertson (1968b, p. 141-143)

JARVIS TOWNSHIP**RESERVE LAKE OCCURRENCE****Location**

Latitude 46°39'00"N, Longitude 84°10'30"W.
Jarvis Township.

Remarks

Mineralization occurs in shear zones in granite. Two samples taken by M.C. Gardiner of Haileybury, Ontario, assayed 0.034 and 1.35 percent U₃O₈ (radiometric equivalent).

References

Robertson (1968a, p. 5-6)

JOHNSON TOWNSHIP**DESBARATS RADIOACTIVITY
OCCURRENCE****Location**

Latitude 46°23'06"N, Longitude 83°55'32"W.
Johnson Township.

Remarks

Radioactive veins in diabase reportedly occur on the northeast shore of Desbarats Lake.

References

Frarey (1962)
Giblin et al. (1979)
Financial Post Survey of Mines (1957, p.203)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Johnson Township, Drill Log Report
Nos. 10 and 11, Desbarats Mining
Company Limited, 1956.

JUILLETTE TOWNSHIP**FANO RADIOACTIVITY OCCURRENCE****Location**

Latitude 46°21'34"N, Longitude 82°57'26"W.
Juillette Township.

Remarks

This occurrence is underlain by feldspathic quartzite, conglomerate and arkose of the Matinenda Formation of the Elliot Lake Group. A 6-inch thick band of oligomictic conglomerate was intersected in one of the drill holes by Fano Uranium Mines Limited in 1955, and indicated a radioactivity count of 2 to 3 times the background.

In 1970 D.S. Robertson and Associates completed an airborne electromagnetic and magnetic survey over the area.

References

Robertson (1963, p. 63)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Juillette Township, Drill Log Report
No. 14, Fano Uranium Mines Limited,
1955.

Technical File No. 63.2800, D.S.
Robertson and Associates Limited,
1970.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000133.

**MATINENDA RADIOACTIVITY
OCCURRENCE****Location**

Latitude 46°22'00"N, Longitude 83°03'40"W.
Juillette Township.

Remarks

Thin seams and beds of conglomerate with minor pyrite and chalcopyrite, and with radioactivity up to five times background, were intersected in two drill holes. A pyritized seam at 2020 feet gave a radioactivity count of nine times the background. Two drill holes totalling 4851 feet were drilled by Matinenda Uranium Mines Limited in 1955.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Juillette Township, Drill Log Report
Nos. 12 and 13, 1955.

Technical File No. 63.2394, Radex
Syndicate, 1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000134.

KAMICHISITIT TOWNSHIP

BLIND RIVER OCCURRENCE

Location

Latitude 46°29'20"N, Longitude 83°01'50"W.
Kamichisitit Township.

Remarks

Surface exposures consist of conglomerate, quartzite and arkose of the Gowganda Formation of the Cobalt Group. A deep drill hole intersected a 90-foot bed of radioactive conglomerate, the best section of which assayed 0.017 percent U_3O_8 over 18.6 feet.

In 1968 airborne magnetic and electromagnetic surveys were completed by Pacific Petroleums Limited.

In 1975 airborne magnetic, electromagnetic, radiometric, and resistivity surveys and two diamond drill holes totalling 4965 feet were completed by Superior Northwest Incorporated.

References

Giblin et al. (1979)
The Northern Miner (Sept. 23, 1976, p. 17;
Oct. 21, 1976, p.11)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2303, Pacific
Petroleum Limited, 1968.

Technical File No. 2.1576, Superior
Northwest Incorporated, 1974.

Kamichisitit Township, Drill Log
Report No. 30, Superior Northwest,
1976.

KIRKWOOD TOWNSHIP

KIRKWOOD TOWNSHIP RADIOACTIVITY OCCURRENCE

Location

Latitude 46°20'26"N, Longitude 83°28'24"W.
Kirkwood Township.

Remarks

A bed of quartz-pebble conglomerate and arkosic grit, about 30 cm thick, occurs within coarse-grained pink-to-grey arkose 2 000 m

south of Kirkwood Lake. The conglomerate contains about 15 percent fine pyrite and is radioactive up to ten times the background. The conglomerate is located within mafic volcanic rocks near the base of the Thessalon Formation.

References

Bennett (1976)

LABONTE TOWNSHIP

HUCLIF PORCUPINE RADIOACTIVITY OCCURRENCE

Location

Latitude 47°19'16"N, Longitude 84°29'20"W.
Labonte Township.

Remarks

Radioactivity is associated with pegmatites within Early Precambrian granite gneiss. The granitic gneiss is cut by numerous southwest-trending Keweenawan diabase dikes. These pegmatites gave geiger counter readings of 2 to 2.5 times the background (Assessment Files Research Office, File 63.587, Ontario Geological Survey, Toronto).

In 1954, geophysical surveys were completed by Matachewan Canadian Gold Limited. In 1955, geophysical surveys, sampling, and drilling were completed by Huclif Porcupine Mines Limited.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.587, Huclif
Porcupine Mines Limited, 1955.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000053.

JALORE OCCURRENCE

Location

Latitude 47°21'40"N, Longitude 84°35'05"W.
Labonte Township.

Remarks

Radioactivity is associated with a zone of calcite-hematite stringers within a northeast-trending Keweenawan diabase dike. The radioactive zone, 10 to 12 feet wide, trends nearly at right angles to the diabase

dike and extends for about 15 feet into granite on each side. Uraninite was identified as the radioactive mineral. This deposit is of hydrothermal origin (Nuffield 1955).

References

Giblin et al. (1979)
Nuffield (1955)
Robertson (1968a, p. 52)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Labonte Township, Drill Log Report,
Jalore Mining Company, Jupiter
Group, 1949.

Technical File 63A.137, Jalore Mining
Co. Limited, 1952.

Technical File 2.2270, R.A.

MacGregor, 1976.

Technical File 2.2588, R.A.

MacGregor, 1978.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000054.

LAPASKA OCCURRENCE

Location

Latitude 47°20'16"N, Longitude 84°35'00"W.
Labonte Township.

Remarks

Radioactivity is present in a fracture zone at or near the contact of a Keweenawan diabase dike and the Early Precambrian granite complex. The diabase dike, approximately 70 feet wide strikes N40E and dips vertically. The 3-foot wide fracture zone strikes N50W and dips 38°SW. The fractures are narrow and filled with calcite, hematite and probably pitchblende. Radioactivity readings were up to five times the background.

References

Giblin et al. (1979)
Robertson (1968a, p. 53)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.76, Lapaska
Mines Limited, North Group, 1949.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000055.

MCDEVITT OCCURRENCE

Location

Latitude 47°19'50"N, Longitude 84°35'00"W.
Labonte Township.

Remarks

Radioactive anomalies are located in pegmatites within an Early Precambrian granite and syenite complex. All the rocks are exposed in a 100 to 200-foot high vertical scarp.

No. 1 showing consists of a branching fracture, striking N75E and dipping 70°N in red pegmatite. The fracture is exposed for a length of 20 feet and is from 12 to 18 inches wide. A sample across 18 inches assayed 0.10 percent U₃O₈.

No. 2 showing consists of a single, flat dipping fracture in grey pegmatite. Near the fracture there is a band of coarse-grained biotite mica that is partly chloritized and is radioactive.

No. 3 showing consists of uranium oxide stain on light grey pegmatite.

References

Giblin et al. (1979)
Robertson (1968a, p. 18)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.93, F. Joubin,
1949.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000057.

TOOKER RADIOACTIVITY OCCURRENCE

Location

Latitude 47°19'40"N, Longitude 84°31'40"W.
Labonte Township.

Remarks

This occurrence is situated in Superior Provincial Park. Two samples from claim SSM169442, sent to the Ontario Department of Mines by E.O. Tooker, showed 0.024 and 0.75 percent U₃O₈ (radiometric equivalent).

References

Giblin et al. (1979)
Lang (1952, p. 53)

LECARON TOWNSHIP

KIRKPATRICK LAKE RADIOACTIVITY OCCURRENCE

Location

Latitude 46°39'44"N, Longitude 83°01'10"W.
Lecaron Township.

Remarks

This occurrence is underlain by an outlier of the Lorrain Formation which consists of quartzite and thin, hematitic, weakly radioactive conglomerate, overlying the Gowganda Formation and Early Precambrian granite. Radioactivity in two thin conglomerate beds was less than twice background.

In 1954, geological mapping and two drill holes totalling 1149 feet were completed on claim SSM27552 and 27555 by Blue Lake Mining Syndicate Limited.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Lecaron Township, Drill Log Report
Nos. 10 and 11, Blue Lake Mining
Syndicate Limited, 1954.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000035.

LEFROY TOWNSHIP

FRYE RADIOACTIVITY OCCURRENCE

Location

Latitude 46°17'16"N, Longitude 83°42'16"W.
Lefroy Township.

Remarks

This occurrence is underlain chiefly by the Mississagi Formation of the Hough Lake Group. Feldspathic quartzite with a few thin radioactive pebble beds occurs in the upper 50 feet. In 1955, a drill hole totalling 1405 feet was drilled by H.E. Frye.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Lefroy Township, Drill Log Report No.
10, 1956.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000009.

WEBB OCCURRENCE

Location

Latitude 46°20'50"N, Longitude 83°41'25"W.
Lefroy Township.

Remarks

A sheared quartzite lies adjacent to the Murray Fault. The shear zone is exposed over a length of 5 to 10 feet. An assay of the outcrop material gave 0.02 percent U₃O₈.

References

Giblin et al. (1979)

Robertson (1968a, p. 7)

LEHMAN TOWNSHIP

BRACEMAR RADIOACTIVITY OCCURRENCE

Location

Latitude 46°27'41"N, Longitude 82°26'01"W.
Lehman Township.

Remarks

Quartz-pebble conglomerate bands with radioactivity up to 2 times the background were intersected at 3031 and 3066 feet, and had an average thickness of 3 feet.

In 1955, one drill hole totalling 3216 feet was drilled on claim S67166 by Bracemar Mines Limited.

In 1966 and 1968, airborne magnetic and electromagnetic surveys and one drill hole to 4946 feet were completed by Kerr-McGee Corporation.

References

Giblin et al. (1979)

Robertson (1962, p.65)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Lehman Township, Drill Log Report
No. 18, Bracemar Mines Limited,
1955.

Lehman Township, Drill Log Report
No. 20, Kerr-McGee Corporation,
1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000080.

MACK TOWNSHIP**BLACK LAKE RADIOACTIVITY OCCURRENCE****Location**

Latitude 46°16'45"N, Longitude 82°52'55"W.
Mack Township.

Remarks

Radioactivity up to four times the background has been reported from sheared argillite of the Pecors Formation of the Hough Lake Group.

References

Giblin et al. (1979)
Robertson (1968a, p. 9)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Mack Township, Drill Log Report No. 15, Harico Mining and Development Company Limited, 1954.
Technical File No. 63.2201, Denison Mines Limited, 1967.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000014.

MAECK TOWNSHIP**SEABROOK LAKE CARBONATITE COMPLEX****Location**

Latitude 47°00'00"N, Longitude 83°17'00"W.
Maek Township.

Remarks

The complex consists of carbonatite and ijolite. The carbonatite occurs in the northern part of the complex. The ijolitic rocks in the southern portion vary in composition from pyroxenite to pegmatitic segregations of urtite. The main part of the complex is enveloped by an inner zone of fenitized granite breccia and an outer zone of fenitized granite.

References

Sage (1976)
Thurston et al. (1977)

MCMURRAY TOWNSHIP**FIRESAND RIVER CARBONATITE COMPLEX****Location**

Latitude 47°59'00"N, Longitude 84°39'40"W.
McMurray and Lastheels Townships.

Remarks

The Firesand River carbonatite complex consists of a dolomite-rich core and a calcite-rich border. The complex consists almost exclusively of carbonatite rocks, ijolite and nepheline syenite.

References

Parson (1961)
Robertson (1981)
Sage (1977)

SURLUGA OCCURRENCE**Location**

Latitude 47°57'55"N, Longitude 84°44'03"W.
McMurray Township.

Remarks

Radioactive mineralization occurs in a pegmatite dike intruding an Early Precambrian granite complex. A sample from the dike assayed 0.10 percent U₃O₈.

References

Lang (1952, p. 135)
Milne et al. (1972)
Robertson (1968a, p. 53)

MONESTIME TOWNSHIP**RUSSIAN LAKE RADIOACTIVITY OCCURRENCE****Location**

Latitude 46°45'44"N, Longitude 82°12'00"W.
Monestime Township.

Remarks

This occurrence is underlain by Early Precambrian granite which is intruded by Keweenawan diabase dikes. Radioactivity three times the background count was obtained near Russian Lake.

References

Sherran (1952)

MONTGOMERY TOWNSHIP**DEMAREST LAKE OCCURRENCE****Location**

Latitude 46°22'28"N, Longitude 83°05'18"W.
Montgomery Township.

Remarks

Two thin bands of uraniferous, pebbly, feldspathic quartzite of the Matinenda Formation of the Elliot Lake Group were intersected at 3562 feet below the surface. Assays gave the following results:
3516 - 3517 feet 0.11 percent U₃O₈, 0.22 percent ThO₂;
3562 - 3563.8 feet 0.07 percent U₃O₈, 0.14 percent ThO₂.

Wakefield Uranium Mines Limited drilled a hole to 1358 feet which Rio Tinto Canadian Explorations Limited deepened to 3946 feet in 1966.

References

Giblin et al. (1979)
Robertson (1968a, p. 10)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000015.

MORIN TOWNSHIP**SUTHERLAND OCCURRENCE****Location**

Latitude 46°32'22"N, Longitude 83°36'10"W.
Morin Township.

Remarks

Locally, rocks of the Matinenda Formation are overlain by rocks of the Ramsay Lake and Mississagi Formations.

In 1968, W.D. Sutherland and Associates drilled six holes and encountered uranium in quartz-pebble conglomerates.

References

Chandler (1973, p. 60)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Morin Township, Drill Log Report No. 10, W.D. Sutherland and Associates, 1968.

Technical File No. 63.2490, 1969.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000016.

RAIMBAULT TOWNSHIP**BOYMAR OCCURRENCE****Location**

Latitude 46°33'08"N, Longitude 82°47'58"W.
Raimbault Township.

Remarks

Uranium mineralization is associated with narrow pyritic pebble bands found in the green arkose of the Mississagi Formation of the Hough Lake Group. The best assay values from samples taken by Hollinger Mines Limited were 0.01 percent U₃O₈ (radiometric equivalent).

In 1954, 15 drill holes totalling 4590 feet were drilled by Boymar Gold Mines Limited.

In 1968 and 1969, ground and airborne magnetic, electromagnetic, and radiometric surveys were completed by Hollinger Mines Limited.

References

Robertson (1977b)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Raimbault Township, Drill Log Report No. 13, Boymar Gold Mines Limited, 1954.

Technical File No. 63.2469, Hollinger Mines Limited, 1968.

Technical File No. 63.2491, Hollinger Mines Limited, 1969.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000119.

GOD'S LAKE MINES OCCURRENCE**Location**

Latitude 46°33'38"N, Longitude 82°49'08"W.
Raimbault Township.

Remarks

Drilling intersected slightly radioactive, thin pebble beds of the Mississagi Formation. The beds occur within an arkosic matrix and contain sparse pyrite and chalcopyrite. The best geiger counter reading was 460 counts per minute over 1.5 feet.

God's Lake Mines Limited completed radiometric surveys and six drill holes in Nicholas and Raimbault townships.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Raimbault Township, Drill Log Report
No. 11, God's Lake Mines Limited,
1954.

Technical File No. 2.1575, Fort
Norman Explorations Incorporated,
1974.

Technical File No. 63.2581, Astonish
Uranium Mining Corporation Limited,
1969.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000132.

ZENMAC RADIOACTIVITY OCCURRENCE

Location

Latitude 46°32'39"N, Longitude 82°47'16"W.
Raimbault Township.

Remarks

Slightly radioactive pebble bands were
found in the Mississagi Formation. At
approximately 300 feet depth, radioactivity
was four to eight times the background.

In 1954, two drill holes were drilled
totalling 1373 feet by Zenmac Metal Mines
Limited.

References

Giblin et al. (1979)
Robertson (1977b)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Raimbault Township, Drill Log
Reports No. 10 and 12, Zenmac Metal
Mines Limited, 1954.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000120.

Regional Geologists' Files, Ontario Ministry
of Natural Resources, Sault Ste. Marie
Raimbault Township, File No. SSM
421, Zenmac Metal Mines Limited,
1954.

RIX TOWNSHIP

FALCONBRIDGE OCCURRENCE

Location

Latitude 47°14'26"N, Longitude 84°37'08"W.
Rix Township.

Remarks

Radioactive anomalies are located along
the sheared contacts of a Keweenawan
diabase dike and Early Precambrian granite.
Pitchblende is associated with calcite-

hematite gangue that fills fractures in the
shear zone and in the diabase.

References

Giblin et al. (1979)
Nuffield (1956, p.25)
Robertson (1968a, p.25)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Rix Township, Drill Log Report,
Falconbridge Nickel Mines Limited.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000071.

ROCHE OCCURRENCE

Location

Latitude 47°14'21"N, Longitude 84°36'24"W.
Rix Township.

Remarks

The radioactive anomaly occurs in
fractures at the contact of a Keweenawan
diabase dike and Early Precambrian granite.
The fractures occurring in the diabase are
filled with a calcite-hematite gangue
containing some pitchblende.
In 1949, prospecting and stripping were
carried out for F.D. Roche.

References

Giblin et al. (1979)
Nuffield (1956)
Robertson (1968a, p. 25)

ROLLINS TOWNSHIP

JAMES OCCURRENCE

Location

Latitude 46°56'00"N, Longitude 83°16'00"W.
Rollins Township.

Remarks

In the northwest portion of Rollins
Township there is an exposure of alkalic
syenite carbonatite and related rocks of the
Middle and Late Precambrian age. A few
diabase dikes of Keweenawan age are also
present. Two samples contained microscopic
grains of sulphide minerals and showed 0.103
percent U₃O₈.

References

Giblin et al. (1979)
Lang (1952, p.129)

SAGARD TOWNSHIP**MOUNT LAKE AREA**

- a. Quebec Uranium Mining Occurrence
- b. Wakman-Grinnicke Occurrence
- c. Soderstrom-Hammerstrom Occurrence
- d. Silver Men Occurrence

Location

- a. Latitude 46°41'00"N, Longitude 82°45'25"W
 - b. Latitude 46°41'40"N, Longitude 82°44'00"W
 - c. Latitude 46°41'10"N, Longitude 82°43'50"W
 - d. Latitude 46°38'00"N, Longitude 82°43'00"W.
- Sagard Township.

Remarks

All thorium deposits in the Mount Lake area are associated with the Lorrain Formation. It consists of interbedded quartzite and conglomerate. The conglomerate is thinly-bedded and hematitic. Radioactivity is usually 2 or 3 times the background.

References

Robertson (1968a, p. 13; 1977a)

SALTER TOWNSHIP**SHUNCK OCCURRENCE****Location**

Latitude 46°15'10"N, Longitude 82°04'00"W.
Salter Township.

Remarks

The highest uranium assay obtained was 0.012 percent U₃O₈ from a 4-foot thick bed of argillaceous quartzite.

References

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto
Technical File No. 2.2409, Report for R. Shunck, 1977.

SLATER TOWNSHIP**MANWOOD RADIOACTIVITY OCCURRENCE****Location**

Latitude 47°08'12"N, Longitude 84°38'57"W.
Slater Township.

Remarks

Radioactive anomalies occur at the contact of Keweenawan diabase dikes and Early Precambrian granite. The occurrence consists of four different showings. All are small and have low geiger counter readings.

In 1949, geological and radiometric surveys, stripping, and trenching were carried out by Manwood Mining Corporation Limited.

References

Robertson (1968a, p. 53)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto
Technical File No. 63A.80, Manwood Mining Corporation Limited, 1949.

PENNWOOD OCCURRENCE**Location**

Latitude 47°13'30"N, Longitude 84°37'40"W.
Slater Township.

Remarks

Radioactive anomalies occur at the sheared contacts of Keweenawan diabase dikes and Early Precambrian granite. The contacts are filled with quartz and calcite stringers with a bright red stain due to the presence of hematite. Some radioactivity of up to 8 times the background was found in the granite and associated pegmatites. Uraninite and ellsworthite were identified.

References

Robertson (1968a, p. 24)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto
Technical File No. 63A.67, Pennwood Gold Mines Limited, 1949.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000066.

SMILSKY TOWNSHIP**RANROUYN OCCURRENCE****Location**

Latitude 47°12'59"N, Longitude 84°34'14"W.
Smilsky Township.

Remarks

The radioactive anomalies occur in the Early Precambrian granites at or near the contact with northeast-trending Keweenawan diabase dikes. Radioactivity was also associated with pegmatite veins and pegmatite boulders that contained small amounts of uraninite.

During 1949 and 1950, geological, ground magnetometer, and geiger counter surveys were completed by Ranrouyn Mines Limited.

References

Giblin et al. (1979)
Lang (1952, p. 132)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.71, Ranrouyn
Mines Limited, 1949.

Technical File No. 63A.215, Ranrouyn
Mines Limited, 1950.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000044.

STRAIN TOWNSHIP**MADAWANSON LAKE OCCURRENCE****Location**

Latitude 46°36'10"N, Longitude 82°10'50"W.
Strain Township.

Remarks

This occurrence is underlain by Early Precambrian granite which is intruded by Keweenawan diabase dikes. At the contact radioactivity is three times the background. A grab sample showed 0.05 percent U₃O₈ (radiometric equivalent).

References

Giblin et al. (1979)
Sheeran (1952)

STRIKER TOWNSHIP**CYR OCCURRENCE****Location**

Latitude 46°11'45"N, Longitude 82°55'48"W.
Striker Township.

Remarks

The arkose and conglomerate of the Matinenda Formation of the Elliot Lake Group rest on the Early Precambrian granite and dip 35°S. The maximum thickness of the conglomerate is 2 feet. A grab sample assayed 0.02 percent U₃O₈.

In 1959, exploration was carried out on this property in conjunction with the Cane property (to the south) by Pronto Uranium Mines Limited.

References

Robertson (1964; 1968a, p.10)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000018.

TARBUTT TOWNSHIP**TARBUTT MINES OCCURRENCE****Location**

Latitude 46°23'16"N, Longitude 83°58'04"W.
Tarbutt Township.

Remarks

Two carbonate veins, 1.5 feet apart and 80 feet long, occur in diabase. The veins strike N60W and dip 87°N. Pitchblende is contained within the veins.

References

Giblin et al. (1979)
Robertson (1968a, p. 11)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 000020.

THESSALON TOWNSHIP**WEST THESSALON OCCURRENCE****Location**

Latitude 46°16'25"N, Longitude 83°29'30"W.
Thessalon Township.

Remarks

Weakly radioactive quartz-pebble conglomerate beds occur at the top of the Livingstone Creek Formation and in the basal part of the Thessalon Formation.

References

Frarey (1977, p. 79)
Giblin et al. (1979)

TIMMERMANS TOWNSHIP**ANUWON OCCURRENCE****Location**

Latitude 46°23'52"N, Longitude 82°51'18"W.
Timmermans Township.

Remarks

Minor intersections of radioactive quartz-pebble conglomerate beds with minor pyrite and radioactivity up to eight to ten times the background were encountered in drilling. In one drill hole a 6.8-foot conglomerate bed gave a radiometric reading of 0.013 percent U_3O_8 . The beds are typically less than 8 inches thick.

References

Giblin et al. (1979)
Robertson (1963, p. 56-58)
The Financial Post Survey of Mines (1956, p.357)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log
Report No. 23, Anuwon Uranium Mines
Limited, 1955.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000122.

Regional Geologists Files, Ontario Ministry
of Natural Resources, Sault Ste. Marie
File No. SSM 424.

FANO RADIOACTIVITY OCCURRENCE**Location**

Latitude 46°22'19"N, Longitude 82°54'34"W.
Timmermans Township.

Remarks

This occurrence is underlain by feldspathic quartzite, conglomerate and arkose of the Matinenda Formation of the Elliot Lake Group. The sedimentary rocks lie unconformably on Early Precambrian granite

at depths ranging from 471 to 584 feet below the surface. The sedimentary rocks strike northeast and dip gently south.

Scintillometer readings of oligomictic, uraniferous quartzite intersected in five drill holes averaged three times the background.

References

Robertson (1963, p. 63)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Timmermans Township, Drill Log
Report No. 37, Fano Uranium Mines
Limited, 1955.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000126.

FORT NORMAN RADIOACTIVITY OCCURRENCE**Location**

Latitude 46°25'50"N, Longitude 82°51'34"W.
Timmermans Township.

Remarks

Radioactivity is associated with conglomerate and quartzite of the lower Matinenda Formation of the Elliot Lake Group.

Scintillometer readings on sections of drill cores of the Matinenda Formation, by Fort Norman Explorations Incorporated, averaged 1.5 times the background. Chemical analyses showed less than 0.005 percent U_3O_8 and less than 0.005 percent ThO_2 .

Between 1967 and 1977, geophysical surveys were carried out by Cominco Limited, Denison Mines Limited, and Fort Norman Explorations Incorporated. Fort Norman also drilled three holes totalling 4374 feet.

References

Giblin et al. (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2171, Cominco
Limited, 1967.

Technical File No. 2.1575, Fort
Norman Explorations Incorporated,
1974.

Technical File No. 2.1850, Fort
Norman Explorations Incorporated,
1975.

Timmermans Township, Drill Log

Report No. 44, Fort Norman Explorations
Incorporated, 1976-77.

VIEL TOWNSHIP

WESTON RADIOACTIVITY OCCURRENCE

Location

Latitude 46°39'32"N, Longitude 82°36'24"W.
Viel Township.

Remarks

A radioactive zone in red to purple-coloured quartz-pebble conglomerate is approximately 40 feet long and 10 to 12 feet wide. Radioactivity was reported to be up to ten times the background.

References

Wood (1975)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.545, Weston &
Company Incorporated, 1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001115.

WHITMAN TOWNSHIP

CRESTLAND OCCURRENCE

Location

Latitude 46°43'50"N, Longitude 83°53'00"W.
Whitman Township.

Remarks

A carbonated, rusty shear zone was traced for 350 feet. A series of trenches up to 18 feet deep was dug. Assays ranged from 0.046 percent to 0.05 percent U_3O_8 (chemical).

In 1948, trenching was performed by Consolidated Northland Mines Limited.

References

Giblin et al. (1979)

Robertson (1968a, p. 11)

URANIUM AND THORIUM DEPOSITS OF COCHRANE DISTRICT

PITT TOWNSHIP

CALMOR OCCURRENCE

Commodity

Uranium and thorium.

Radioactive Minerals

Pitchblende and monazite.

Location

Latitude 50°09'50"N, Longitude 81°37'03"W.

Pitt Township.

Map Reference: ODM Map P.370.

Geology

The uranium mineralization occurs predominantly in a carbonate vein averaging 0.5 feet wide and 240 feet long. It strikes north and dips 75°W. This vein cuts Early Precambrian gneiss, pegmatites and diabase. A second vein, 10 feet to the east strikes due north and dips 65-75°W, and is exposed for 70 feet. Both veins contain some specularite, pyrite, and chalcopyrite. This occurrence is on the escarpment that marks the northern edge of the exposure of Precambrian rocks and the southern edge of the Paleozoic rocks of the James Bay Basin.

Economic Features

Three samples taken from the main vein assayed 0.03 percent, 0.12 percent, and 0.055 percent U₃O₈ (radiometric equivalent) (Hogg 1948, p.9).

History of Development

1947: The vein was discovered by A. Mosher for Calmor Mines Limited.

1948: Moneta Porcupine Mines Limited carried out a detailed geiger counter survey and a geological reconnaissance survey.

References

Hogg (1948, p.5-10)

Lang et al. (1962, p.276)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.65, D.J.

McDougall, 1948.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto

File No. 0000141.

MINOR URANIUM AND THORIUM OCCURRENCES OF COCHRANE DISTRICT

HABEL TOWNSHIP

SOWESKA RIVER OCCURRENCE

Location

Latitude 50°24'16"N, Longitude 82°33'04"W.
Habel Township.

Remarks

Traces of uranium mineralization are associated with lignite beds.

References

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 83-1-44, Algoma
Central Railway and Hudson's Bay
Railway, 1966.

PITT TOWNSHIP

CORAL RAPIDS OCCURRENCE

Location

Latitude 50°09'45"N, Longitude 81°37'15"W.
Pitt Township.

Remarks

The Sextant Formation here forms the Precambrian basement. An anomalous layer within this formation, 20 m thick with 0.05 to 0.1 pounds per ton U_3O_8 , was intersected in drillholes by Kerr Addison Mines Limited and Hudson Bay Mining Limited along the margin of a 10 km wide basinal structure.

In 1978, assays of diamond drill core revealed 0.10 pounds per ton U_3O_8 over 0.20 m in coarse arkosic beds at the southwest edge of a basin trending north-northwest.

References

Bennett et al. (1968)
Tihor and Hunt (1979)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 83.1-127, Licence
of Occupation for Kerr Addison Mines,
Limited, 1977-1979.

SHACKLETON TOWNSHIP

PROVENCHER OCCURRENCE

Location

Latitude 49°17'21"N, Longitude 81°52'24"W.
Shackleton Township.

Remarks

Mineralization is found in pegmatites.

References

Lang (1952, p. 277)

URANIUM AND THORIUM DEPOSITS OF KENORA DISTRICT

DEPOSITS ON UNSURVEYED LAND

AEROBUS LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Uranophane.

Location
Latitude 50°21'30"N, Longitude 93°21'07"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology
Uranium mineralization is found in medium-to coarse-grained quartz monzonite near the contact with quartz-feldspar-biotite metasediments. Yellow staining is common.

Economic Features
Of two drill holes, one showed 0.5 percent U_3O_8 (radiometric equivalent) over 5.5 feet while the other assayed 0.07 percent U_3O_8 (radiometric equivalent) over 4.5 feet (Assessment Files Research Office, Drill Log Report No. 10, 1955, Ontario Geological Survey, Toronto).

Grab samples taken by ministry staff assayed 0.03 and 0.006 percent U_3O_8 (Beard and Scott 1976).

History of Development
1955: Trenching, geological and scintillometer surveys, and nine drill holes totalling 678 feet were completed by S. Duggan and C. St. Paul.

References
Beard and Scott (1976)
Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto
Aerobus Lake Area, Drill Log Report No. 10, 1955.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
Aerobus Lake Area, File 000298.
Resident Geologist's Files, Ontario Ministry of Natural Resources, Kenora

BAMAJI LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Pitchblende, uraninite and brannerite.

Location
Latitude 51°10'31"N, Longitude 91°26'07"W.
District of Kenora (Patricia Portion).
Map Reference: ODM Map 2247.

Geology
The "Main" showing is just south of a small lake, 1 km south of the western end of Moosetegon Lake. Radioactivity was detected in a series of trenches over a strike length of 125 m. The uranium and thorium occurs in a stratigraphic unit of calc-silicate material up to 3 m wide striking approximately N75E, consisting of actinolite, calcite, dolomite and magnetite. A second area is located approximately 700 m west of the main occurrence. In a series of shallow trenches, narrow veinlets (2 to 4 cm) of actinolite-biotite cut porphyritic biotite trondhjemite. Grab samples collected by Sage and Breaks (1982) of the veinlet material and of the surrounding trondhjemite gave values of 0.04 percent and 0.06 percent U_3O_8 respectively (Sage and Breaks 1982, p.77). A third showing found by R. Knappett by reconnaissance prospecting with a scintillometer, is located approximately 600 m south of the eastern end of North Bamaji Lake. It consists of a thin (less than 400 m) unit of calc-silicate material, predominantly actinolite. This unit is similar to the host rock in the main showing, 1000 m along the regional strike to the east.

At least two other small uranium-thorium occurrences lie between North Bamaji Lake and Moosetegon Lake. All may occur within the same units or units in close proximity. Further evidence suggests that the uranium-bearing units may be of considerable lateral extent.

Economic Features
The highest values obtained by Urangesellschaft Canada Limited were in the range 0.060-0.290 percent U_3O_8 .

History of Development
1954: Trenching, stripping and 11 diamond drill holes were completed by McCombe Mining and Exploration Limited.

1968: An airborne spectrometer survey was carried out by the Kirkland Townsite Gold Mines Limited.

1975: Magnetometer and electromagnetic survey of the Bamaji Lake claim group was completed by Dome Exploration (Canada) Limited.

1979: Exploration activities were carried out by Urangesellschaft (Canada) Limited.

References

Bond and Breaks (1978)

Sage and Breaks (1982)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2510, Kirkland
Townsite Gold Mines Limited, Fry
Lake Area, 1968.

Technical File No. 2.3258,
Urangesellschaft (Canada) Limited,
Fry Lake Area, 1979.

Technical File No. 2.1705, Dome
Exploration (Canada) Limited, Fry
Lake Area, 1975.

Fry Lake Area, Drill Log Report No.
10, McCombe Mining and Explorations
Limited, 1954.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 000302, Kirkland Townsite.

BEARHEAD LAKE PROSPECT

Commodity

Uranium.

Radioactive Minerals

Uranophane, uranothorite, and zircon.

Location

Latitude 52°46'42"N, Longitude 93°44'56"W.

Showing	Latitude	Longitude
1	52°46'25"	93°44'00"
1A	52°46'30"	93°44'30"
1B	52°46'45"	93°45'00"
1C	52°47'25"	93°47'25"
1D	52°48'00"	93°49'50"
1E	52°45'45"	93°43'35"
1F	52°45'15"	93°42'00"
1G	52°47'35"	93°48'35"
1H	52°43'50"	93°38'00"
1I	52°41'30"	93°38'00"
1J	52°39'40"	93°38'30"
1K	52°39'20"	93°36'00"
1L	52°41'00"	93°34'00"
1M	52°39'00"	93°29'00"
1N	52°40'00"	93°28'00"
1O	52°39'00"	93°26'00"

1P	52°40'00"	93°23'10"
1Q	52°48'40"	93°51'20"
1R	52°37'30"	93°20'00"
1S	52°44'00"	93°39'00"
1T	52°12'30"	93°32'10"

District of Kenora (Patricia Portion).
Map Reference: ODM Map 2262.

Geology

Uranium mineralization occurs in pegmatite-bearing zones between biotite-granite gneiss to the south and a migmatite-granite gneiss-pegmatite complex to the north. The strike of the contact is N65-70W. All of the rocks are of Early Precambrian age.

The "East" showing (1H on Map P.2425, Robertson 1982) is approximately 500 feet long, 20-100 feet wide, trending N65W, dipping 60-90°S; assays ranged from 0.01 to 0.10 percent U₃O₈ (radiometric equivalent).

The "Camp" showing (1A on Map P.2425) is a continuation of the "East" showing. It strikes N67-73W, dips steeply south, is 250 feet long, and 25-30 feet wide. Assays ranged from 0.005 to 0.018 percent U₃O₈ (radiometric equivalent).

The "Bear Creek" showing (1B on Map P.2425) strikes N68-70W, is 200 feet long and 25-30 feet wide. Assays ranged from 0.009 to 0.209 percent U₃O₈ (radiometric equivalent).

The "Father" showing (1C on Map P.2425) strikes N70-87W, dips 70-90°S, is 600 feet long, and 75-150 feet wide. Assays show 0.131 percent U₃O₈ (radiometric equivalent).

The "Crag" showing (1D on Map P.2425) is approximately 1500 feet long.

The dimensions of the "Mother" showing (1Q on Map P.2425) are unknown.

Assays from the remaining showings are as follows:

"Zone A" (1F, 1D, 1C, and 1G on Map P.2425) 0.045 percent U₃O₈;

"Zone B" 0.09 percent U₃O₈;

"Zone C" 0.070 percent U₃O₈;

"Zone D" (1E on Map P.2425) 0.025 percent U₃O₈;

"Zone E" 0.20 percent U₃O₈;

"Zone F" 0.03 percent U₃O₈;

"West" showing 0.03 percent U₃O₈.

The other showings correspond with numbers on Map P.2425 (Robertson 1982) as follows:

"Zone H" 1J, 1K, 1I

"Zone I" 1L

"Zone K" 1M, 1N, 1O

"Zone M" 1P

Economic Features

In 1977 Kerr Addison Mines Limited and Dolores Bench Resources drilled 29,871 feet in 66 holes. The indicated reserves are 978,810 tons averaging 0.06 percent U_3O_8 to a depth of approximately 500 feet.

History of Development

1929: Ontario Department of Mines survey by M.E. Hurst (1930).

1955: Surface mapping, aerial and ground radiometric surveys, trenching and seven diamond drill holes totalling 1423 feet were completed on claims KRL546636, KRL154640 and KRL60453 by Cam Mines Limited.

Favourable Mines drilled three diamond drill holes for a total of 1250 feet on claims KRL5987-KRL6009.

1969: Stripping, trenching, blasting, scintillometer surveys, and eight diamond drill holes totalling 7084 feet were completed by the Keevil Mining Group Limited.

1971: Twenty-six trenches were dug on Zones A,B,C,D,E,F, and West by the Keevil Mining Group Limited.

1977: A total of 29,871 feet of drilling was completed by Kerr Addison Mines Limited and Dolores Bench Resources.

References

Ayres et al. (1972)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63.797, Sigasco Explorations Limited, 1957.

Technical File No. 63.2278, Cam Mines Limited, 1969.

Setting Net Lake Area, Drill Log Report Nos. 10-17, 1956-1959.

Technical File No. 2.286, Keevil Mining Group Limited, 1971.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 00304.

CAMERON-ALCOCK OCCURRENCE

Commodity

Uranium, thorium.

Radioactive Minerals

Monazite, uraninite.

Location

Latitude $50^{\circ}00'00''N$, Longitude $94^{\circ}31'32''W$.

District of Kenora.

Map Reference: ODM Map 2175.

Geology

Radioactive mineralization occurs in pegmatite dikes that intrude metavolcanic remnants that are altered by granite and pegmatites intrusive along a schistosity. The dike strikes N15W, is exposed over a length of 30 feet, and has a width of 9 feet.

Economic Features

Counts of up to eight times the background were obtained in a pit on the dike. Grab samples assayed 0.17 percent U_3O_8 (radiometric equivalent) and 0.14 percent U_3O_8 (chemical equivalent). Chip samples were taken by Minorex Limited in existing trenches. Their assays averaged 0.22 pounds U_3O_8 per ton.

History of Development

1952: Pits were dug by M. Cameron and C. Alcock.

1967: An airborne radiometric survey was completed by Headvue Mines Limited.

1979: Radiometric, magnetometric and geological mapping was undertaken by Minorex Limited.

References

Chisholm (1951, p. 2-3)

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 2.3136, Minorex Limited, 1979.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File 000297

CAN-FER OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Uranophane.

Location

Latitude $50^{\circ}21'07''N$, Longitude $94^{\circ}41'52''W$
District of Kenora.

Map Reference: ODM Map 2175.

Geology

Uranium mineralization is associated with light grey and pink granitic pegmatite dikes which trend east. The highest radioactivity occurs along minor biotite slips and shears.

Economic Features

In 1968, samples collected by ministry staff from two showings 15 and 5 feet wide averaged 0.065 percent and 0.38 percent U_3O_8 respectively (Resident Geologist's files, Ontario Ministry of Natural Resources, Kenora).

In 1975, six trenches along strike of a zone of high radiation 20 to 40 feet wide averaged 0.002 percent to 0.006 percent U_3O_8 (Beard and Scott 1976).

History of Development

1968: Trenching, scintillometer survey and assaying were carried out by Steve Lesavage.

1970: A geological survey was carried out by Ontario Department of Mines staff.

1975: Trenching and assaying were undertaken by the Ministry of Natural Resources' regional staff.

References

Beard and Scott (1976)

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2634, Can Fer
Mines Limited, 1969.

Technical File No. 2.169, Bralorne Can
Fer Resources Limited, 1970.

Technical File No. 2.2212, Noranda
Exploration Company Limited, 1976.

Resident Geologist's Files, Kenora.

DAVIDSON LAKE OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°29'26"N, Longitude 95°08'50"W.

District of Kenora

Map Reference: ODM Map 2175.

Geology

This area is underlain by medium-grained, grey granitic rocks with occasional sections and remnants of well-bedded grey, quartz-biotite metasediments. Radioactivity is confined to coarse-grained pink pegmatite phases with coarse biotite within the granite.

Economic Features

Grab samples taken by Ministry of Natural Resources staff assayed 0.05, 0.03,

and 0.006 percent U_3O_8 (Beard and Scott 1976).

History of Development

1955-1956: Airborne radiometric, magnetometer, scintillometer, and ground electromagnetic surveys, and diamond drilling were carried out by Anglo Barrington Mines Limited.

1970: Geophysical and geological reports were completed by Consolidated Manitoba Mines Limited.

References

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.751, Anglo
Barrington Mines Limited, 1955-1956.

Technical File No. 2.331, Consolidated
Manitoba Mines Limited, 1970.

Technical File No. 2.652, Consolidated
Manitoba Mines Limited, 1971.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto

File No. 000196.

Resident Geologist's Files, Ontario Ministry
of Natural Resources, Kenora

Kenora Files, Davidson Lake.

HALLIDAY OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°36'32"N, Longitude 95°04'04"W.

District of Kenora.

Map Reference: ODM Map 2175.

Geology

This area is underlain by biotite gneiss cut by narrow dikes of granite and granodiorite. Moderate radioactivity is associated with the granite dikes.

Economic Features

Chemical analyses returned abundances of 0.01 percent U_3O_8 and 0.01 percent ThO_2 . Radiometric equivalent values averaged 0.02 percent U_3O_8 in dikes 0.06 to 4.9 m wide.

History of Development

1969: Six diamond drill holes totalling 305 feet were drilled by L.B. Halladay.

1979: A radiometric survey was undertaken by W.G. Wahl Limited.

References

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Snowshoe Lake Area, Drill Log Report
No. 10, 1969.

Technical File No. 2.2943, W.G. Wahl,
Limited, 1979.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 001129.

HEADVUE OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Uraninite and monazite.

Location

Latitude 50°01'19"N, Longitude 94°32'13"W.
District of Kenora.

Map Reference: ODM Map 2175.

Geology

Uranium mineralization is associated with
pegmatite intruding strongly foliated Early
Precambrian gneiss.

Economic Features

Unknown

History of Development

1967: Aeroradiometric and ground
magnetometer surveys as well as trenching
and sampling were completed by Headvue
Mines Limited.

1979: Radiometric and magnetometer
surveys, geological mapping and sampling
were carried out by Minorex Limited.

References

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 2.3136, Minorex
Limited, 1979.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 001077

Resident Geologist's Files, Ontario Ministry
of Natural Resources, Kenora

JENSON OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Uraninite and thorite.

Location

Latitude 49°50'57"N, Longitude 93°25'54"W.
Langton Township.

Map Reference: OGS Map 2443.

Geology

Uranium mineralization occurs in two
lenses of gneissic pegmatitic granite. They
are 3 to 5 feet wide by 50 feet long and have
intruded Early Precambrian lavas that
contain much magnetite.

Economic Features

A radioactive survey showed counts up to
18 times the background. A bulk sample
assayed by the Ministry's regional staff had
an average content of 0.054 percent U_3O_8
(radiometric equivalent). The average of the
grab samples from two channel samples was
0.016 percent U_3O_8 (radiometric equivalent)
(Lang 1952, p.118).

History of Development

1955: Four diamond drill holes totalling 806
feet were drilled by A.O. Lantz on claims
K17455 and K17460.

References

Blackburn (1981)

Lang (1952)

Robertson (1968a, p. 57)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Langton Township, Drill Log Report
No. 12, M. Jenson, 1955.

Resident Geologist's Files, Kenora.

PANCER OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°19'30"N, Longitude 94°35'22"W.
District of Kenora.

Map Reference: ODM Map 2175.

Geology

Uranium mineralization occurs in five east-trending pegmatite dikes, cutting granite. Radioactivity is erratic, but highest near biotite concentrations. There are three main showings with dimensions as follows: 400 feet by 60 to 150 feet, 200 feet by 30 feet, and 12 feet by 6 feet.

Economic Features

Two samples averaged 0.13 percent U_3O_8 (chemical). Grab samples gathered by Ontario Department of Mines staff averaged 0.004 percent U_3O_8 (Assessment Files Research Office, Ontario Geological Survey, Toronto, Technical File No. 63.633, 1955).

History of Development

1955: Geological and scintillometer surveys were completed by S. Ciglen.
1970: Ten diamond drill holes totalling 3010 feet were drilled by P. Davidson.
1979: A magnetic survey was completed over this area.

References

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63.633, Paterson Lake Area, 1955.

Paterson Lake Area, Drill Log Report No. 12, 1970.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File 000299.

PERCH LAKE OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°04'57"N, Longitude 94°33'04"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology

Folded metasediments and metavolcanics have been intruded by later Precambrian granites, quartz monzonites and pegmatites. Biotite-rich pegmatites have the strongest radioactivity. A 30-foot long trench exposed coarse-grained pegmatite with local concentrations of biotite.

Economic Features

Grab samples gave assays of 0.086 percent U_3O_8 and 0.03 percent ThO_2 . Grab samples from surface test pits assayed by the Ontario Geological Survey ranged from 0.44 to 1.72 pounds per ton (Beard and Rivett 1978).

History of Development

1950: Trenching by C. Alcock.

1969: Airborne radiometric work was completed by C. Morton.

1976-1977: The property was optioned to Anschutz Oil Company, which in conjunction with the owner, D. Loudon, completed ground radiometric and geological surveys.

References

Ferguson et al. (1967)

Beard and Rivett (1978)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 2.2405, Anschutz Uranium Corporation, 1977.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File 001125

PIRSON OCCURRENCE (MEDICINE STONE LAKE OCCURRENCE)**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°54'38"N, Longitude 94°10'09"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology

The radioactivity is associated with a mass of sheared hornblende granite containing many "greenstone" inclusions. Thorium accounts for a large portion of the radioactivity.

Economic Features

The greatest radioactivity occurs in areas approximately 2 to 8 feet in width, underlain by shear zones in "greenstone" inclusions. Grab samples which have been assayed show 0.047 to 0.079 percent U_3O_8 .

History of Development

Unknown.

References

Ferguson et al. (1967)
Robertson (1968a, p. 62)

SCHRYBURT LAKE OCCURRENCE**Commodity**

Uranium and niobium.

Radioactive Minerals

Unknown.

Location

Latitude 52°36'00"N, Longitude 89°36'10"W.
District of Kenora (Patricia Portion).
Map Reference: ODM Map P. 2236.

Geology

This occurrence is situated within the Schryburt Lake carbonatite complex. It is composed of calcitic carbonatite. Bands of nearly pure actinolite, apatite, magnetite, biotite, or pyrrhotite alternate with pink to pinkish-white carbonatite. The texture is equigranular and medium-grained.

Economic Features

In 1961, E.I. Dupont and De Nemours Company Limited and Many Lakes Exploration Company Limited completed 28 pits and trenches. The uranium and niobium values are in uranium-bearing pyrochlore. A sample of weathered carbonatite assayed 0.039 percent U_3O_8 and 0.08 percent Nb_2O_5 (assay by Geoscience Laboratories, Ontario Geological Survey)

References

Sage and Wright (1979)

SNOOK LAKE OCCURRENCE**Commodity**

Uranium, thorium.

Radioactive Minerals

Uranophane.

Location

Latitude 50°12'35"N, Longitude 94°40'05"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology

Uranium mineralization is associated with homogeneous coarse-grained pink quartz

monzonite having a vague but distinctive porphyritic appearance. There are occasional coarse spots of magnetite.

Economic Features

Chip samples collected by Beard and Rivett in 1977 across a width of 40 m, were assayed by the Ontario Geological Survey. One 20 m sample (a chip every 15 cm) assayed 0.54 pounds U_3O_8 and 0.24 pounds ThO_2 per ton. The second sample over a width of 21 m, assayed 0.12 pounds U_3O_8 and 0.26 pounds ThO_2 per ton (Beard and Rivett 1978).

History of Development

1976: Trenching and one diamond drill hole were completed by J. Harrison.
1977: Trenching was carried out by J. Harrison.

References

Beard and Rivett (1978)
Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora.

TOURIST LAKE PROSPECT**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°16'04"N, Longitude 94°40'08"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology

Uranium mineralization is associated with the narrow and highly sheared portions of two narrow northeasterly trending bands of biotite schist and gneiss.

Economic Features

Assays of samples taken in 1969 averaged 1.25 pounds per ton U_3O_8 over an average width of 7.5 feet.

In 1976, assays by Huronian Mines returned values of 0.9 pounds U_3O_8 per ton over 9 feet. In three holes completed in one 23-foot section, assay values averaged 0.56 pounds U_3O_8 per ton.

History of Development

1968-1970: Airborne radiometric survey, geological, ground magnetometer, scintillometer surveys, 31 trenches totalling 721 feet, and three Winkie drill holes totalling 113 feet were completed by Bralorne Can-Fer Resources Limited.

1975: Sampling of the westerly trenches was done by the Hanna Mining Company and Denison Mines Limited.

1976: Nine drill holes totalling 1635 feet were drilled by Consolidated Summit Mines Limited.

References

Ferguson et al. (1967)
Prospectus, Consolidated Summit Mines Limited, 1976.
The Northern Miner, Vol. 62, No. 22 and 39, 1976.

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto
Paterson Lake Area, Drill Log Report Nos. 13 and 14, 1976.
Technical File No. 2.307, 1971.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 001126.

TREELINED LAKE OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°08'04"N, Longitude 94°40'00"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology

Metasediments in this location are generally siliceous, contain sulphides and are highly metamorphosed and locally recrystallized. Interspersed are zones of biotite-rich granitic rocks. Pegmatites are also present. The highest radioactivity is associated with biotite-rich granitic rocks.

Economic Features

Grab samples assayed 0.008 to 0.02 percent U_3O_8 and 0.28 percent ThO_2 .

History of Development

1968: Linklatter dug trenches and pits.
1976: This area was restaked by Harrison and Perkins.

1976: Sampling was carried out by T. Skimming.

References

Beard and Rivett (1977)
Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora

VERMILION LAKE OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 50°01'37"N, Longitude 94°30'00"W.
District of Kenora.
Map Reference: ODM Map 2175.

Geology

The geology is similar to that of the Cameron-Alcock occurrence where radioactivity is associated with a pegmatite dike which has intruded metavolcanic remnants.

Economic Features

Unknown.

History of Development

1967: Airborne radiometric survey was completed by Headvue Mines Limited.

References

Ferguson et al. (1967)
Robertson (1968a, p. 6)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto
Technical File No. 63.2344, Headvue Mines, Limited, 1967.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File 000296.

WOLF ISLAND OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 49°39'40"N, Longitude 94°32'36"W.
District of Kenora.
Map Reference: OGS Map 2443.

Geology

Mineralization is associated with a 6-inch, buff coloured quartz-carbonate hydrothermal vein exposed in pillowed, basaltic lava for 20 feet. The vein consists of fine-grained quartz and carbonate and is sparsely mineralized with fine pyrite and chalcopyrite.

Economic Features

A grab sample taken by E.O. Chisholm assayed 0.10 percent U₃O₈ (radiometric equivalent) and a trace of gold.

History of Development

1950: Samples were taken by E.O. Chisholm.

References

Blackburn (1981)
Robertson (1968a, p.62)

Energy, Mines and Resources, Canada Files
Mineral Development Sector, Department of Energy, Mines and Resources, National Mineral Inventory File U1, 52/E/10, Wolf Island, Sept. 1963.

BRIDGES TOWNSHIP**COULEE OCCURRENCE (KIMBER LAKE - WEST OCCURRENCE)**

Commodity
Uranium.

Radioactive Minerals

Uraninite, uranophane, and beta-uranotile.

Location

Latitude 49°50'33"N, Longitude 93°37'12"W.
Bridges Township.
Map Reference: ODM Map 2303.

Geology

The property is underlain by Early Precambrian metasediments which have been intruded by felsic granitic rocks. These intrusions vary in composition from medium- to coarse-grained granodiorite to pink pegmatite.

Uranium mineralization is associated with pegmatite. Secondary minerals are concentrated along the discontinuous and randomly oriented fractures near the surface.

Economic Features

Samples taken from the trenches and pits by Coulee Lead and Zinc Mines Limited showed assays running from 0.02 to 0.42 percent U₃O₈. The highest radiometric value obtained from a drill core by Noranda Mines Limited was 0.08 percent U₃O₈ over 5 feet. Most values were less than 0.02 percent U₃O₈.

History of Development

1955: Trenching and two drill holes totalling 90 feet were completed by F. Mallery.

1967: An airborne radiometric survey followed by ground radiometric and magnetometer surveys, trenching and pitting were completed by Coulee Lead and Zinc Mines Limited.

1968: The property was optioned to Noranda Mines Limited who completed a geological survey, blasted numerous trenches and drilled four diamond drill holes totalling 1384 feet.

1974: A geological report was completed by Imperial Oil Limited.

1977: Anschutz Uranium Mines Limited completed a geological evaluation.

References

Pryslak (1976, p. 37-38)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Bridges Township, Drill Log Report No. 17, 1968.

Technical File No. 63.2267, Coulee Lead and Zinc Mines Limited, 1968.

Technical File No. 2.1665, Imperial Oil Limited, 1974.

Technical File No. 2.2451, Anschutz Uranium Mines Limited, 1977.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000281.

FAIRSERVICE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals

Uraninite, uranophane, and allanite.

Location

Latitude 49°50'00"N, Longitude 93°41'22"W.
Bridges Township.
Map Reference: ODM Map 2303.

Geology

Radioactivity was associated with the pegmatitic phases of the granitic rocks of Early Precambrian age. One of several radioactive pegmatite dikes in the area can be traced over a length of approximately 1550 feet, and is well exposed along the rock-cut for the natural gas pipeline. This dike is approximately 25 feet wide at this point and contains abundant secondary uranium minerals along fracture planes.

Economic Features

Grab samples collected by the Ministry's regional staff assayed 0.051 percent and 0.025 percent U_3O_8 (Beard and Scott 1976, p.6).

Samples taken by Sherritt Gordon Mines Limited assayed 1.52 to 2.8 pounds U_3O_8 per ton. Molybdenum is also present here.

History of Development

1967-1968: Three pits were dug by Noranda Mines Limited.

1976: Two drill holes totalling 243 feet were drilled by Sherritt Gordon Mines Limited. This company also completed scintillometer and magnetometer surveys.

1977: Robert Fairservice drilled and sampled. Geological mapping was undertaken by D. Pike.

References

Beard and Scott (1976)
Pryslak (1976, p.6)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 2.2506, R.

Fairservice, 1977.

Bridges Township, Drill Log Report No. 20, Sherritt Gordon Mines Limited, 1976.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 001122.

HEADWAY OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Uraninite.

Location

Latitude $49^{\circ}50'37''N$, Longitude $93^{\circ}40'28''W$.
Bridges Township.
Map Reference: ODM Map 2303.

Geology

The highest radioactivity occurs in association with the biotite-rich zones within pegmatites.

Economic Features

Two grab samples from two showings assayed 4.2 pounds U_3O_8 per ton (Northern Miner 1967, Oct.26).

History of Development

1967: Grab samples were taken by Headway Red Lake Gold Mines Limited.

An aeroradiometric survey was completed by Coulee Lead and Zinc Mines Limited.

1969: One diamond drill hole was completed by Noranda Exploration Company.

1974: Imperial Oil Limited completed a geological survey in the area.

References

Northern Miner (1967, Oct.19, Oct.26)
Pryslak (1976)
Robertson (1968a, p. 56)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Bridges Township, Drill Log Report No. 14, Noranda Mines Limited, 1969.
Technical File No. 2.1665, Imperial Oil Limited, 1974.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000282.

DOCKER TOWNSHIP**KIMBER LAKE (EAST) OCCURRENCE (C.S. STEPHENS)****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude $49^{\circ}50'36''N$, Longitude $93^{\circ}34'15''W$.
Docker Township.
Map Reference: ODM Map 2303.

Geology

An easterly trending radioactive pegmatite dike occurs along the north shore of Kimber Lake, near the east end of the lake. The dike ranges from 100 to 250 feet in width and can be followed for 4000 feet. The highest radioactivity is associated with

biotite-rich and apatite-rich phases of the pegmatite.

Economic Features

Chip samples taken across 10-foot sections from a trench contained an average of 0.024 percent U_3O_8 .

History of Development

1968: Pitting and trenching by C.S. Stephens.
1974: A geological survey was completed by Imperial Oil Limited.

References

Pryslak (1976, p.50)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File 2.1665, Imperial Oil
Limited, 1974.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 001124.

Resident Geologist's Files, Kenora.

DROPE TOWNSHIP

BLUETT LAKE OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Allanite and uranophane.

Location

Latitude 49°58'01"N, Longitude 92°38'09"W.
Drope and Breithaupt Townships.
Map Reference: OGS Map 2443.

Geology

Uranium mineralization is contained in a sill of pegmatite 35 feet wide, bounded by quartz-biotite metasediments, and striking N45E.

Economic Features

Thirteen grab samples taken from a 3-foot exposure assayed from 0.25 to 5.4 percent U_3O_8 . Molybdenite is also present.

History of Development

1968: Prospecting, trenching and sampling were done by Conwest Exploration Company Limited.

References

Blackburn (1981)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Sioux Lookout

LANGTON TOWNSHIP

LOUDON PROSPECT

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 49°51'46"N, Longitude 93°21'39"W.
Langton Township.
Map Reference: OGS Map 2443.

Geology

Radioactive showings occur in an area underlain by granite containing quartz monzonite and amphibolite.

Economic Features

In 1974 R.C. Beard, Regional Geologist, Ontario Ministry of Natural Resources, Kenora, carried out scintillometer surveys in the area. Radioactive readings were 8000 counts per minute, compared to a background of 1000 counts per minute.

In 1975 Kerr Addison Mines Limited assayed some drill core samples. The average value was 0.25 percent U_3O_8 over 30 to 40 feet.

References

Blackburn (1981)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Langton Township, Drill Log Report
No. 20, Kerr Addison Mines Limited,
1975.

Resident Geologist's Files, Kenora
Tew and Loudon Prospect, Kenora.

PRESTON OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 49°51'09"N, Longitude 93°25'45"W.
Langton Township.
Map Reference: OGS Map 2443.

Geology

The mineralized zone consists of radioactive pegmatite cutting biotite gneiss and granite.

Economic Features

Drilling by Preston East Dome Mines Limited, in 1955, cut 25 feet of radioactive pegmatite that averaged 0.28 percent U_3O_8 .

History of Development

1954-1955: One diamond drill hole was drilled by J. McLeod on claim K17346.

1955: Twenty-eight drill holes totalling 2132 feet and five trenches were completed by Preston East Dome Mines Limited on claim K17437.

References

Blackburn (1981)
The Northern Miner, Vol. 62, No. 28, Sept. 23, 1976, p. C.18.

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Langton Township, Drill Log Report No. 14, 1955.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000285.

MACNICOL TOWNSHIP**FOOT LAKE OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 49°49'12"N, Longitude 93°59'50"W.
MacNicol Township.
Map Reference: ODM Map 2302.

Geology
Radioactive mineralization occurs in sections in granite within mafic metavolcanics.

Economic Features
One 10-foot section assayed 0.07 percent U_3O_8 .

History of Development
1955: One diamond drill hole was drilled by Kerr Addison Mines Limited.

References

Pryslak (1974a)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora.

HAWK LAKE OCCURRENCE (BYBERG-CAMPBELL-MACFARLANE OCCURRENCE)

Commodity
Uranium.

Radioactive Minerals

Uraninite, euxenite, thorite, uranophane, and beta-uranotile.

Location

Latitude 49°48'47"N, Longitude 94°00'16"W.
MacNicol Township.
Map Reference: ODM Map 2302.

Geology

This area is underlain by mafic to intermediate volcanic rocks (hornblende schist) which are intruded by numerous irregular dikes and masses of pegmatites. Uranium mineralization is associated with masses and stringers of coarse magnetite within the pegmatite.

Economic Features

Twenty-three grab samples from the vicinity of the main showing averaged less than 0.05 percent U_3O_8 .

Grab samples taken by Kerr Addison Mines Limited in 1975 assayed 0.41 to 1.9 pounds U_3O_8 per ton.

History of Development

Up to 1954: The main showing, on claim K16596, had been explored by stripping and trenching, one open cut over a length of 130 feet, and a geiger counter survey.

1954: Trenching was done by the Great Lakes Uranium Corporation.

1975: Kerr Addison Mines Limited performed diamond drilling.

1977: CANMET (Canada Centre for Mineral and Energy Technology) indicated a new mineral resembling allanite. It assayed 67 percent U_3O_8 , 21 percent PbO_2 , 8.5 percent ThO_2 , and 0.5 percent cerium.

References

Beard and Scott (1976)
Pryslak (1974a)
Robertson (1968a, p.57-58, 1981)
Satterly (1955, p.1-5)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
MacNicol Township, Drill Log Report
No. 20, Kerr Addison Mines Limited,
1975.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000288.

KENORATOMIC OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Uraninite, monazite, allanite, kasolite,
uranophane, and beta-uranolite.

Location
Latitude 49°50'36"N, Longitude 93°55'23"W.
MacNicol Township.
Map Reference: ODM Map 2302.

Geology
The area is underlain by foliated to gneissic metavolcanics, which are mafic to intermediate in composition, and have been intruded by widespread dikes and irregular bodies of pegmatite.
Uranium mineralization is associated with pegmatite, magnetite, biotite, and occasional molybdenum. This property is on the eastern extension of the deposit held by New Campbell Island Mines Limited.

Economic Features
An assay of a sample from this zone was 0.098 percent over 3.5 feet. Another sample taken at the north end of Richard Lake assayed 0.087 percent U_3O_8 across 2.5 feet and 0.064 percent U_3O_8 across 5 feet (Pryslak 1976).

History of Development
1955: Trenching and two diamond drill holes totalling 394 feet were completed by Burning Rock Uranium Mines Limited.

Trenching and four diamond drill holes totalling 1082 feet were completed by Acko Mines Limited at the west end of the property.

1956-1957: Geological, scintillometer and magnetometer surveys, and 12 diamond drill holes totalling 2500 feet were completed by Kenoratomic Mines Limited.

1978: Geological and geophysical surveys were completed on the Richard Lake property by Golden Standard Mines Limited.

References
Pryslak (1976, p.45-46)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 63.916,
Kenoratomic Mines Limited, 1957.
MacNicol Township, Drill Log Report
No. 17.
Technical File No. 2.2905, Golden
Standard Mines Limited, 1978.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000287.

RICHARD LAKE PROSPECT (NEW CAMPBELL ISLAND)

Commodity
Uranium.

Radioactive Minerals
Uraninite, uranothorite, allanite, and beta-uranolite.

Location
Latitude 49°50'52"N, Longitude 93°55'00"W.
MacNicol Township.
Map Reference: ODM Map 2302.

Geology
Uranium mineralization is associated with a series of parallel magnetite-bearing pegmatite dikes cutting highly recrystallized, foliated to gneissic, Early Precambrian metavolcanics.

The prospect consists of four zones. The two southernmost contain dikes of low grade material 5 to 7 feet in width. The two central zones are the widest. The dikes here range from 10 to 20 feet in width and extend for at least 700 feet. Radioactivity is, in places, associated with magnetite-rich sections within the pegmatites.

Economic Features
A.S. Bayne and Company estimated reserves at 650,000 tons grading 0.10 percent U_3O_8 based on an explored length of 700 feet, a depth of 1000 feet, and an average width of 10 feet (Pryslak 1976).

History of Development
1954 - 1956: Seventeen diamond drill holes, totalling about 5000 feet, are located over a strike length of 1300 feet. Underground exploration included two adits totalling 1100 feet, and horizontal drill holes totalling 255 feet (Pryslak 1976).

1976: Rollmac Exploration Limited remapped the underground workings and carried out ground magnetic and scintillometer surveys.
 1977: Golden Standard Mines Limited drilled 10 diamond drill holes totalling 2000 feet. The drilling doubled the strike length of the zone. Magnetometer and radiometric surveys were also carried out.
 1978: International Standard Resources Limited, formerly Golden Standard Mines Limited, performed geological and geophysical surveys.

References

Beard (1977)
 Prysak (1976, p.45-46)
 Robertson (1968a, p.58)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario Geological Survey, Toronto
 Technical File No. 2.2480, Rollmac Exploration Limited, 1977.
 Technical File No. 2.2905, Rollmac Exploration Limited, 1978.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 000289.

TEMPLE TOWNSHIP

BOTTLE BAY OCCURRENCE

Commodity
 Uranium.

Radioactive Minerals
 Uranophane.

Location
 Latitude 49°47'11"N, Longitude 93°16'05"W.
 Temple Township.
 Map Reference: OGS Map 2443.

Geology
 This property is underlain by metasediments (arkose) and hornblende granite intruded by masses and dikes of coarse-grained granite pegmatite. Radioactivity is associated with apatite in the granite pegmatite where uraniferous staining is reported.

Economic Features
 Radiometric surveying indicates values up to 1.0 pound U₃O₈ per ton. Ontario Division of Mines assays recorded 0.005 and 0.083 percent U₃O₈ (Beard and Scott 1976).

History of Development

1975-1976: Trenching, radiometric surveys and one diamond drill hole were completed by F.O.B. Mining and Exploration Limited.
 1977: F.O.B. Mining and Exploration Limited completed a geological report.

References

Beard and Scott (1976)
 Blackburn (1981)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario Geological Survey, Toronto
 Temple Township, Drill Log Report, F.O.B. Mining and Exploration Company Limited, 1976.
 Technical File No. 2.2261, F.O.B. Mining and Exploration Limited, 1976.
 Technical File No. 2.2668, F.O.B. Mining and Exploration Company Limited, 1977.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 000669.

TUSTIN TOWNSHIP

ASCOT OCCURRENCE

Commodity
 Uranium.

Radioactive Minerals
 Uraninite and uranophane.

Location
 Latitude 49°50'37"N, Longitude 93°52'12"W.
 Tustin Township.
 Map Reference: ODM Map 2302.

Geology
 Uranium mineralization is associated with pegmatite that intrudes a belt of Early Precambrian biotite paragneiss. The pegmatite is 1 mile wide and crosses the claim group at a strike of N80E and dip of 70°N.

Economic Features
 Three radioactive zones (A, B, and C) are associated with pegmatite dikes. Radiation ranges from 2 to 4 times the background.

History of Development
 1955-1956: Geological, magnetic and radiation surveys were carried out by the Ascot Metals Corporation Limited. One diamond drill hole reached 51 feet. Eight

trenches and some stripping in Zone A, extensive stripping in Zone B and some blasting and stripping in Zone C were completed.

Some surface work was completed by Burning Rock Uranium Mines Limited.

1956-1957: Unspecified work was carried out by Kenoratomic Mines Limited.

References

Pryslak (1976, p.49)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.601, Ascot
Metal Corporation, Hawk Lake Area,
R.C. Coutts, 1955.

Technical File No. 63.916,
Kenoratomic Mines Limited, 1956.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000286.

PETURSSON LAKE OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Allanite and uraninite.

Location

Latitude 49°50'13"N, Longitude 93°49'01"W.

Tustin Township.

Map Reference: ODM Map 2302.

Geology

Uranium mineralization occurs in a zone 360 m by 75 m. This zone consists of coarse- to medium-grained pink pegmatite and quartz monzonite which parallel the contact between the granite and metavolcanics.

Economic Features

Assays average 0.56 pounds per ton U₃O₈.

History of Development

1977: A radiometric survey and five Winkie diamond drill holes totalling 365 feet were carried out by Sherritt Gordon Mines Limited.

1978: Ground magnetometer, radiometric and spectrometer surveys were completed by Sherritt Gordon Mines Limited.

References

Pryslak (1974a)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora, Petursson Lake.

Assessment Files Research Office, Ontario
Geological Survey, Toronto

Tustin Township Drill Log Report No. 14, 1977.

Technical File No. 2.2699, Sherritt
Gordon Mines Limited, 1978.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 001134.

WABIGOON TOWNSHIP

BENCHMARK OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Uranophane.

Location

Latitude 49°53'00"N, Longitude 93°21'00"W.

Wabigoon Township.

Map Reference: OGS Map 2443.

Geology

Radioactive pegmatite was exposed in a roadcut on Highway 105, approximately 800 feet north of the contact between the Wabigoon Belt and the English River Subprovince. Outcrops consist of coarse-grained pegmatite and quartz monzonite which intrude mafic trondhjemite. The pegmatite - quartz monzonite contains coarse magnetite masses up to 30 mm in diameter.

Economic Features

Radioactivity is associated with the magnetite. Scintillometer readings range up to 20 000 counts per minute. A grab sample assayed 0.022 percent U₃O₈ and 15 ppm thorium.

History of Development

1977: Geological and scintillometer surveys were completed by the Ontario Geological Survey.

References

Blackburn (1981)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora.

MINOR URANIUM AND THORIUM OCCURRENCES OF KENORA DISTRICT

OCCURRENCES ON UNSURVEYED LAND

BELAIR LAKE RADIOACTIVITY
OCCURRENCE**Location**

Latitude 50°36'30"N, Longitude 94°35'35"W.
District of Kenora (Patricia Portion)

Remarks

The Belair Lake occurrence is situated in the Sydney Lake Cataclastic Zone which represents the contact between the Uchi Subprovince to the north and the English River Subprovince to the south (Breaks et al. 1975).

In the area of the occurrence is found white, homogeneous diatexite, medium-grained with small spots of relic biotite throughout and 40 percent paleosome material, occasional pegmatite sections, with moderate cataclasis. Radioactivity was found to be three times the background.

References

Beard and Rivett (1978)
Breaks et al. (1975)

CHASE LAKE RADIOACTIVITY
OCCURRENCE**Location**

Latitude 50°35'30"N, Longitude 94°53'00"W.
District of Kenora (Patricia Portion)

Remarks

This occurrence is situated in the Sydney Lake Cataclastic Zone which represents the contact between the Uchi Subprovince and the English River Subprovince (Breaks et al. 1975).

The occurrence is underlain by homogeneous diatexite with 1 percent widely scattered, thin relic bands of sedimentary rock up to 15 cm wide. The diatexite is slightly cataclastic. Radioactivity is three times the background.

References

Beard and Rivett (1978)
Breaks et al. (1975)

HORNBY LAKE OCCURRENCE

Location

Latitude 52°31'30"N, Longitude 93°37'30"W.
Latitude 52°31'00"N, Longitude 93°35'30"W.
Latitude 52°29'00"N, Longitude 93°36'20"W.
District of Kenora (Patricia Portion)

Remarks

Uranium mineralization is associated with several batholiths in the vicinity of Hornby Lake "greenstone" belt. In all cases mineralization tends to be near the batholiths.

References

Ayres et al. (1973)

HUSTON LAKE RADIOACTIVITY
OCCURRENCE**Location**

Latitude 50°23'45"N, Longitude 95°05'10"W.
District of Kenora (Patricia Portion)

Remarks

Anomalies in this area reflect a narrow to broad band of supracrustal rocks extending from the Manitoba border to the English River (Breaks et al. 1975). This unit consists of a sequence of metasediments, mostly greywacke with a few volcanic sections, all of which have been subjected to anatexis.

At this location, medium- to fine-grained, pink, nebulitic, quartz monzonite contains 10-15 percent biotite as disseminations and small clumps. Radioactivity is 2.5 times the background.

References

Beard and Rivett (1978)
Breaks et al. (1975)

JOHNSON LAKE RADIOACTIVITY
OCCURRENCE**Location**

Latitude 50°36'45"N, Longitude 95°37'15"W.
District of Kenora (Patricia Portion)

Remarks

This occurrence is situated in the Sydney Lake Cataclastic Zone which represents the contact between the Uchi Subprovince and the English River Subprovince (Breaks et al. 1975).

Grey, highly folded quartz-biotite metasediment contains (10 percent) narrow lenses and knots of leucosome, and shows little cataclasis. Radioactivity is 1.5 times the background.

References

Beard and Rivett (1978)
Breaks et al. (1975)

OAK LAKE OCCURRENCE

Location

Latitude 50°24'28"N, Longitude 93°56'18"W.
District of Kenora

Remarks

The uranium mineralization occurs where Early Precambrian pegmatite has intruded the metasediments. The uranium minerals are monazite and uraninite. In three diamond drill holes, less than 0.01 percent U₃O₈ (radiometric equivalent) was found.

References

Ferguson et al. (1967)
Lang et al. (1962, p.275)
Robertson (1968a, p.61)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Oak Lake Area, Drill Log Report No.
10.

OCTOPUS LAKE RADIOACTIVITY OCCURRENCE

Location

Latitude 50°37'45"N, Longitude 95°38'35"W.
District of Kenora (Patricia Portion)

Remarks

This occurrence is situated in the Sydney Lake Cataclastic Zone which represents the contact between the Uchi Subprovince and the English River Subprovince (Breaks et al. 1975).

Foliated and sheared granitoid rock is highly cataclastic, contains coarse augen feldspar, and is highly seritic.

Radioactivity here is fairly low.

References

Beard and Rivett (1978)
Breaks et al. (1975)
Ferguson et al. (1967)

PATERSON LAKE OCCURRENCE

Location

Latitude 50°18'00"N, Longitude 94°41'00"W.
District of Kenora

Remarks

The occurrence lies within the English River gneiss belt and all of the rocks are of Early Precambrian age. The uraniumiferous zones contain granodiorite, quartz monzonite, and granite that have intruded metavolcanics and greywacke.

References

Ferguson et al. (1967)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora.

PINENEEDLE LAKE RADIOACTIVITY OCCURRENCE

Location

Latitude 50°46'10"N, Longitude 94°34'45"W.
District of Kenora

Remarks

This occurrence is situated in the Sydney Lake Cataclastic Zone which represents the contact between the Uchi Subprovince and the English River Subprovince (Breaks et al. 1975). White, inhomogeneous diatexite has 50 percent paleosome, and is highly cataclastic. Radioactivity is up to two times the background.

References

Beard and Rivett (1978)
Breaks et al. (1975)

PINE ROAD OCCURRENCE

Location

Latitude 49°46'41"N, Longitude 93°49'01"W.
District of Kenora.

Remarks

The rocks in the area of the occurrence belong to the Feist Lake Group. Radioactivity is confined to the pegmatite bodies in contact with the gneiss and migmatite. The strike lengths of the anomalies are apparently controlled by the strike lengths of the lenses of supracrustal rock with which they are spatially associated.

References

Blackburn (1981)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File 2.2768, Sherritt Gordon
Mines Limited, 1978.
Mineral Deposits Inventory Record, Ontario
Geological Survey, Toronto
File K 0344.

REYNAR LAKE OCCURRENCE

Location

Latitude 50°28'22"N, Longitude 95°07'45"W.
District of Kenora

Remarks

Uranium mineralization is located in
pegmatite and hornblende-biotite gneiss.
Assays from a drill section averaged 1.8
percent U₃O₈ over 2.0 feet.

In 1955 five diamond drill holes were
drilled by E. Anderson.

References

Ferguson et al. (1967)

SYDNEY LAKE RADIOACTIVITY OCCURRENCE

Location

Latitude 50°36'45"N, Longitude 94°32'40"W.
District of Kenora (Patricia Portion)

Remarks

This occurrence is situated in the Sydney
Lake Cataclastic Zone which represents the
contact between the Uchi Subprovince and
the English Subprovince (Breaks et al. 1975).

There is a predominance of leucogranite
with a few sections of relic sedimentary
rock.

References

Beard and Rivett (1978)
Breaks et al. (1975)
Ferguson et al. (1967)

BRIDGES TOWNSHIP

PARTH OCCURRENCE

Location

Latitude 49°50'05"N, Longitude 93°44'18"W.
Tustin and Bridges Townships

Remarks

The main showing consists of pegmatite
lenses, averaging 8 feet wide and occurring

at intervals over a distance of 2 miles. One
assay showed 0.13 percent U₃O₈ (radiometric
equivalent).

References

Lang (1952, p.120)
Pryslak (1974b)
Robertson (1968a, p.59)

WILSON OCCURRENCE

Location

Latitude 49°50'24"N, Longitude 93°40'26"W.
Bridges Township.

Remarks

Uranium mineralization occurs in a pink,
medium- to coarse-grained pegmatite
intrusion. The radioactivity is erratically
distributed and is concentrated in biotite-rich
zones.

In 1955 one diamond drill hole to 146 feet
and two pits were completed by A.L. Wilson.

In 1967 an aeroradiometric survey was
completed by Coulee Lead and Zinc Mines
Limited.

References

Pryslak (1974b)
Robertson (1968a, p.62)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto.

Bridges Township, Drill Log Report
No. 13, 1955.

Source Mineral Deposit Record, Ontario
Geological Survey, Toronto
File 000283.

MCBRIEN TOWNSHIP

COAL CREEK OCCURRENCE

Location

Latitude 50°10'41"N, Longitude 82°58'11"W.
McBrien Township.

Remarks

Uranium occurs in lignite beds which are
associated with quartz and kaolin sands.
These are contained within the Mattagami
Formation.

References

Vos (1975)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 83.1-35, F.R.
Joubin and Associates, 1962.
Technical File No. 83.1-44, Algoma
Central Railway and Hudson's Bay
Railway, 1966.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000140.

MCMMASTER TOWNSHIP

ASARCO OCCURRENCE

Location

Splatter Lake: Latitude 48°56'10"N,
Longitude 88°38'40"W.
Eagle Mountain: Latitude 49°02'00"N,
Longitude 88°35'00"W.
McMaster Township.

Remarks

This area is underlain by red, fine-grained sandstone of the Sibley Group. At least one diabase dike and a diabase sill intrude these sandstones.

In 1977, Asarco Exploration Company of Canada Limited undertook a lake sediment sampling program in the Wolf River area. The two most important areas in the study were the Splatter Lake area and the Eagle Mountain area.

The average value of uranium found in the sediments was 4.02 parts per million.

References

Coates (1967b)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 2.3105, Asarco
Exploration Company of Canada
Limited.

MCNICOL TOWNSHIP

VICEROY OCCURRENCE

Location

Latitude 49°48'36"N, Longitude 93°59'54"W.
McNicol Township.

Remarks

Uranium mineralization occurs in vaguely foliated, grey to slightly pink granodiorite. Pegmatite is present. Some sections of

biotite gneiss resemble assimilated metasediments or metavolcanics.

In 1955 trenching and six diamond drill holes were completed by Viceroy Uranium Corporation Limited.

References

Pryslak (1976, p.51, 1974a)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora

TUSTIN TOWNSHIP

BEE LAKE OCCURRENCE

Location

Latitude 49°50'48"N, Longitude 93°50'39"W.
Tustin Township.

Remarks

Uranium mineralization is contained in an irregular pegmatite mass with a maximum thickness of 100 feet and a length of 2700 feet. The dike decreases in thickness with depth. The dike trends N80W and dips 30 to 60°N.

Radioactivity levels were up to 8 times the background.

References

Pryslak (1976, p.36)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.923, Tustin
Mines Limited, 1958.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000291.

PADDINGTON LAKE OCCURRENCE

Location

Latitude 49°49'58"N, Longitude 93°45'46"W.
Tustin Township.

Remarks

Radioactivity is widespread in a large mass of granitic pegmatite within pyroclastic metavolcanics. A chip sample assayed 0.5 percent U₃O₈ over 15 feet.

In 1977, the occurrence was prospected by R. Fairservice.

References

Blackburn (1981)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kenora
Property examination, Paddington
Lake.

WABIGOON TOWNSHIP

QUIBELL OCCURRENCE (MEEHAN OCCURRENCE)

Location

Latitude 49°57'21"N, Longitude 93°27'12"W.
Wabigoon Township.

Remarks

The uranium mineralization occurs in lenses and dikes of granite and pegmatite cutting metavolcanics which in places are gneissic or schistose. Graphite, molybdenite, and pyrite have been identified.

References

Blackburn (1981)
Robertson (1968a, p.59)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 000292.

URANIUM AND THORIUM DEPOSITS OF NIPISSING DISTRICT

PARDO TOWNSHIP

PICKLE CROW OCCURRENCE

Commodity

Uranium, gold and copper.

Radioactive Minerals

Pitchblende.

Location

Latitude 46°46'05"N, Longitude 80°15'40"W.
Pardo Township.
Map Reference: OGS Map 2361.

Geology

Uranium mineralization is found in a quartz-pebble conglomerate, 2 to 40 feet thick, which locally grades into quartzite. In this area, the radioactive layer forms a sinuous arc opening to the south.

Economic Features

In nine diamond drill holes, an average thickness of 16.3 feet of conglomerate gave analyses of 0.02 to 0.08 percent U₃O₈.

History of Development

1956-1957: Extensive surface work, geologic mapping, and 16 diamond drill holes totalling 10,882 feet were completed by Pickle Crow Mines Limited.
1969: Airborne electromagnetic and magnetic surveys were completed by Kennco (Canada) Explorations Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.63)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63.2445, Kennco (Canada) Explorations Limited, 1969.
Pardo Township, Drill Log Report No. 11, Pickle Crow Gold Mines Limited, 1956.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 00318.

VOGT TOWNSHIP

PROSCO PROSPECT

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°49'21"N, Longitude 80°05'10"W.
Vogt Township.
Map Reference: ODM Map 2048.

Geology

Uranium mineralization occurs in the basal beds of the Huronian Supergroup. In this area it is generally associated with the Matinenda and Mississagi Formations of the Elliot Lake Group. They are composed of quartzite and quartz-pebble conglomerate.

Drilling encountered uranium and gold at depths of 200 and 300 feet.

Economic Features

The best assays were obtained by Aubay Uranium Mines Limited in 1955 and 1956. Values were 0.02 to 0.038 percent U₃O₈ (Assessment Files Research Office, Ontario Geological Survey, Toronto, Technical File No. 63.2642)

History of Development

1955: Four diamond drill holes totalling 2100 feet were drilled by Algoma Gold Mines Limited.

1955-1956: Seven diamond drill holes totalling 1716 feet and a magnetic survey were completed by Aubay Uranium Mines Limited.

1958: Eight diamond drill holes totalling 1842 feet were drilled by Proscow Mines Limited.

1969: Geological, magnetometer, electromagnetic, and scintillometer surveys were carried out by the Keevil Mining Group.
1973: Surface prospecting was carried out by the Gowganda Syndicate.

1974: This property was owned by Rand Reef Mines Limited.

1975: Rand Reef Mines Limited carried out a geological survey.

1976: Rand Reef Mines Limited completed an aeromagnetic survey on the Lake Temagami property.

References

Robertson (1968a, p.63)
Thomson (1960a, p.36-37)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63.2642, Keevil Mining Group Limited, 1970.

Technical File No. 63.3287, Rand Reef
Mines Limited, 1974.

Technical File No. 2.1848, Rand Reef
Mines Limited, 1975.

Technical File No. 2.050, Rand Reef
Mines Limited, 1976.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000322.

MINOR URANIUM AND THORIUM OCCURRENCES OF NIPISSING DISTRICT

STRATHCONA TOWNSHIP

NEIL OCCURRENCE

Location

Latitude 47°02'00"N, Longitude 79°49'00"W.
Strathcona Township.

Remarks

Four small, highly radioactive areas were discovered. Three of these areas contain quartz veins.

References

Card and Lumbers (1977)
Robertson (1968a, p.64)

Ontario Ministry of Natural Resources Files

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
Strathcona Township File.

URANIUM AND THORIUM OCCURRENCES OF RAINY RIVER DISTRICT

MAINVILLE LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 48°49'14"N, Longitude 93°10'48"W.
Rainy River District
Map Reference: OGS Map 2443.

Geology

The Mainville Lake Group is underlain by a series of granite gneisses intruded by great numbers of pegmatite dikes.

A large radioactive pegmatite dike on the south shore of Otter Bay is underlain by a series of rusty weathering, biotite-hornblende gneisses. It varies from 200 to 250 feet in width and is 800 feet long.

Economic Features

A composite sample of pegmatite was taken by H.D. Carlson, Ontario Department of Mines. Typical assays were 0.01 to 0.017 percent U_3O_8 .

History of Development

1955-1957: Thirteen diamond drill holes totalling 1170 feet were drilled. Geological and radiometric surveys were completed by Rainy Lake Mining Company Limited.

References

Blackburn (1973; 1981)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.333, Rainy
Lake Mining Limited, 1957.

Drill Log Report No. 10, Rainy Lake
Mining Limited, 1957.

Resident Geologist's Files, Kenora.

MINOR URANIUM AND THORIUM OCCURRENCES OF RAINY RIVER DISTRICT

TROTIER TOWNSHIP

HEWARD LAKE OCCURRENCE

Location

Latitude 48°46'00"N, Longitude 91°09'00"W.
Trotier Township.

Remarks

This occurrence is situated in
metasedimentary migmatite in muscovite
graphic granite pegmatite.

References

Pirie (1978)

URANIUM AND THORIUM OCCURRENCES OF SUDBURY DISTRICT

BALDWIN TOWNSHIP

BROULAN REEF CENTRAL OCCURRENCE

Commodity

Uranium and thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°20'52"N, Longitude 81°46'23"W.
Baldwin Township.
Map Reference: OGS Map 2361.

Geology

Radioactive quartz-pebble conglomerate beds outcrop intermittently along a strike length of 3000 feet.

Economic Features

In 1966 and 1967, Broulan Reef Mines Limited assayed samples from drill holes and trenches. From the drill holes, the average value for uranium was 1.5 pounds U₃O₈ per ton over 3.7 feet. From the trenches, assays showed 1.2 pounds ThO₂ over 2.0 feet.

History of Development

1966-1967: A geological survey of the area was completed. Three trenches and 14 drill holes totalling 2806 feet were completed. An airborne electromagnetic and radiometric survey was carried out by Broulan Reef Mines Limited.

1970: Conrad Springer drilled three diamond drill holes.

1976: Magnetic and radiometric surveys were carried out by the Glencair Mining Company Limited.

1977: Prospecting was carried out by the Glencair Mining Company Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.65-66)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Baldwin Township, Drill Log Report
No. 34, Broulan Reef Mines Limited,
1966-1967.

Technical File No. 63.2211, Broulan
Reef Mines Limited, 1967-1968.

Technical File No. 2.2275, Glencair
Mining Company Limited, 1976.

Technical File No. 2.2377, 1977.

Technical File No. 63.3479, Glencair
Mining Company Limited.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto

File No. 00420

Regional Geologists Files, Sudbury
File S63-857.

BROULAN REEF LOOKOUT TOWER
OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°20'45"N, Longitude 81°42'45"W.
Baldwin Township.
Map Reference: ODM Map 1952-1.

Geology

Radioactivity is associated with quartz-pebble conglomerate lenses that occur in quartzite of the Matinenda Formation. Three radioactive zones were discovered. The average length is 106 feet and the average width approximately 2.0 feet.

Economic Features

Assays of samples taken by Dominion Gulf Company from lot 3, concession V were 0.12 percent U₃O₈ (radiometric equivalent) and 0.04 percent U₃O₈ (chemical) (Thomson 1960a, p. 26)

History of Development

1953-1954: Geological, radiometric and magnetic surveys were performed by Dominion Gulf Company.

1966-1967: Geological and geophysical airborne surveys were completed by Broulan Reef Mines Limited.

1969: One drill hole totalling 1004 feet was drilled by Broulan Reef Mines Limited.

References

Card et al. (1974)
Thomson (1953, p.33)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Baldwin Township, Drill Log Report
No. 29, Broulan Reef Mines Limited,
1969

Technical File No. 63.512, Dominion
Gulf Company 1954.
Technical File No. 63.2211, Broulan
Reef Mines Limited, 1968.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000379

CANADIAN JOHNS-MANVILLE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°20'08"N, Longitude 81°43'00"W.
Baldwin Township.
Map Reference: OGS Map 2361.

Geology
Radioactivity is associated with quartz-
pebble conglomerate embedded in quartzite.

Economic Features
Assays from a single drill hole graded 0.60
pounds U₃O₈ per ton over 1.1 feet.

History of Development
1966-1967: A single diamond drill hole
totalling 2378 feet, with a wedge at 548 feet
was completed by Canadian Johns-Manville
Company Limited.

References
Card and Lumbers (1977)
Robertson (1968a, p.66)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Baldwin Township, Drill Log Report
No. 22 Canadian Johns-Manville
Company Limited, 1966-1967.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000380.

JELICOE (ESPANOLA BAY) OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°20'35"N, Longitude 81°44'35"W.
Baldwin Township.
Map Reference: OGS Map 2361.

Geology
Radioactivity is associated with quartz-
pebble conglomerate containing streaks and
patches of chalcopyrite, and interbedded with
quartzite.

Economic Features
In 1954, Plum Uranium and Metal Mining
Company Limited drilled eight diamond drill
holes. One assay was 0.008 percent U₃O₈
over 2.6 feet.

History of Development
1954-1955: Five diamond drill holes were
drilled by Jellicoe Mines (1939) Limited
totalling 1889 feet.
1966: Consolidated Montclerg Mines Limited
drilled the showing; no radioactivity was
found.

References
Card and Lumbers (1977)
Robertson (1968a, p.67-68)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Baldwin Township, Drill Log Report
No. 32, Jellicoe Mines (1939) Limited,
1954
Baldwin Township, Drill Log Report
No. 19, Consolidated Montclerg Mines
Limited, 1966.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000384

JELICOE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°20'14"N, Longitude 81°45'28"W.
Baldwin Township.
Map Reference: OGS Map 2361.

Geology
Radioactivity is associated with quartz-
pebble conglomerate in bands that strike east
and dip approximately 45 to 70 degrees north.

Economic Features

Six drill holes were drilled on the SE 1/4, N 1/2, lot 7, concession V. Assays of samples taken were 0.03 percent U_3O_8 (chemical) over 3.5 feet, and 0.01 percent U_3O_8 (radiometric equivalent) over 3.0 feet.

History of Development

1954: Geological mapping and a magnetometer survey were completed by Jellicoe Mines (1939) Limited.

1967: Denison Mines Limited completed an airborne geophysical survey.

A geophysical survey was completed on the Burns and Maki option by Mattagami Mines Limited.

1975: Five diamond drill holes were drilled by Mattagami Lake Mines Limited.

References

Card and Lumbers (1977)

Robertson (1968a, p.67-68)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Baldwin Township, Drill Log Report
No. 11, Jellicoe Mines (1939) Limited,
1954

Baldwin Township, Drill Log Report
No. 36, Mattagami Lake Mines
Limited, 1975

Technical File No. 2.1774, Mattagami
Lake Mines Limited

Technical File No. 63.2170, Denison
Mines Limited, 1967.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000383.

PLUM OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude $46^{\circ}19'18''N$, Longitude $81^{\circ}43'52''W$.
Baldwin Township.

Map Reference: OGS Map 2361.

Geology

A zone of radioactive quartzite and conglomerate is contained in the quartzite formation and has a width of 1 to 9 feet over a strike length of 600 feet and to a slope depth of 185 feet.

Economic Features

The best assay recorded by Plum Uranium and Metal Mining Company Limited in 1954 was 0.069 percent U_3O_8 over 6.2 feet.

History of Development

1916: Prospecting was carried out by Nickol Chemical Company.

1952: An aeromagnetic and scintillometer survey was completed by the Ontario Department of Mines.

1954: Twelve drill holes totalling 1908 feet, and magnetometer and geological surveys were carried out by Plum Uranium and Metal Mines Limited.

1954-1955: Twenty-nine drill holes totalling 6445 feet were drilled by Jellicoe Mines (1939) Limited.

1960: An electromagnetic survey was completed by Evenlode Mines Limited.

1966: Six drill holes totalling 1655 feet were completed by Consolidated Montclerg Mines Limited.

1967: Diamond drilling was carried out by Denison Mines Limited.

1976: Geological mapping and a gamma ray spectrometer survey were carried out by D.S. Robertson and Associates Limited.

References

Card and Lumbers (1977)

Thomson (1953)

Robertson (1968a, p.67-68)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Baldwin Township, Drill Log Report
No. 10, Plum Uranium and Metal
Mining Limited, 1954

Baldwin Township, Drill Log Report
No. 21, Consolidated Montclerg Mines
Limited, 1967

Technical File No. 63.435, Plum
Uranium and Metal Mining Limited,
1954.

Technical File No. 2.2373, D.S.
Robertson and Associates Limited.

CREELMAN TOWNSHIP**LESLIE OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Uraninite.

Location

Latitude 46°55'33"N, Longitude 81°03'47"W.
Creelman Township.
Map Reference: ODM Map 2212.

Geology

Uraninite has been found in the bedded argillite of the Elliot Lake and Hough Lake Groups.

Economic Features

No. 1 Zone consists of a radioactive argillite bed, 18 inches thick, striking S80E and dipping 55 degrees north. Assays gave 0.48 percent U₃O₈ (chemical) and 0.05 percent ThO₂ (chemical) and 0.065 percent U₃O₈.

No. 2 Zone consists of interbedded argillite and conglomerate. Samples assayed 0.08 percent U₃O₈ (Thomson 1960a).

History of Development

1954: A geiger counter survey and 14 drill holes totalling 3635 feet were completed by MacLeod-Cockshutt Gold Mines Limited and Kenogamisis Gold Mines Limited.

1957: Geological mapping was completed by Rio Tinto Canadian Exploration Limited.

1967: Assembly Mines Limited drilled holes totalling 10,554 feet.

1973: An airborne geophysical survey was completed by Gulf Minerals Canada Limited.

1976: An airborne gamma ray spectrometer survey was completed by Ingamar Explorations Limited.

Erana Mines Limited completed a geophysical and radiometric survey.

1978: Issac Burns Exploration Company Limited completed an airborne magnetometer survey.

1979: TX Resources Limited performed a gamma ray spectrometer survey.

References

Meyn (1971, p.39-41)
Thomson (1960a)

**Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto**

Creelman Township, Drill Log Report
No. 10, MacLeod-Cockshutt Gold
Mines Limited, 1954

Drill Log Report No. 11, 12, Assembly
Mines Limited, 1967, 1968

Technical File No. 63A.338, Leslie
Uranium Option, 1957-1958

Technical File No. 2.1214, Gulf
Minerals Canada Limited, 1973

Technical File No. 2.2307, Ingamar
Explorations Limited, 1978.

Technical File No. 2.2994, TX

Resources Limited, 1979

Technical File No. 2.2306, Erana
Mines Limited, 1976

Technical File No. 2.2850, Issac Burns
Explorations Limited, 1978

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto

File No. 000388.

DRURY TOWNSHIP**ALANEN-MAKI OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°24'43"N, Longitude 81°29'42"W.
Drury Township.
Map Reference: OGS Map 2361.

Geology

The deposit consists of a large outcrop of greatly deformed quartzite, argillite, arkose, and pebble conglomerate. The beds of quartz-pebble conglomerate are radioactive (Thomson 1960a).

Economic Features

A grab sample taken by W. Alanen assayed 0.61 percent U₃O₈ (chemical) and 0.30 percent ThO₂ (chemical).

A sample taken by J.E. Thomson, Ontario Department of Mines, from a conglomerate bed 1.5 feet thick, assayed 0.38 percent U₃O₈ and 0.10 percent ThO₂ (chemical).

Acme Gas and Oil Company reported that two drill holes intersected a minimum of 6 beds across a width of 600 feet. Grades of samples from this drilling varied from 1.56 pounds U₃O₈ over 20 feet to 1.8 pounds over 4.5 feet.

History of Development

Pre-1955: Pitting by W. Alanen

1955: W. Alanen drilled a total of 322 feet

1967: Two holes totalling 223 feet were drilled by Acme Oil and Gas Company Limited.

1975: Espina Copper Development drilled some diamond drill holes.

References

Robertson (1968a, p.70-71)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Drury Township, Drill Log Report No. 13, Alanen and Maki, 1955.

Technical File No. 63.717, Garrison Harbour Mines Limited, 1956.

Technical File No. 63.2155, B.W. Lang, 1967.

Drury Township, Drill Log Report No. 20, Acme Oil and Gas Company Limited, 1967.

Technical File No. 2.378, Acme Oil and Gas Company Limited, 1971.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto

File No. 000392.

Regional Geologist's Files, Sudbury
Files S63.117, S63.122.

KERR ADDISON PROSPECT

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°24'36"N, Longitude 81°32'07"W.
Drury Township.
Map Reference: OGS Map 2361.

Geology
Two radioactive zones trend southeast, following the contact between Early Precambrian granite and Middle Precambrian Huronian rocks. The first radioactive zone lies within the basal metasedimentary formation which overlies the granitic and volcanic basement and underlies the main quartzite formation. Several radioactive quartz-pebble conglomerate beds occur in this zone. One is within 100 to 200 feet of the basement and extends along strike for 3000 feet.

A second zone of radioactivity lies within the main quartzite formation. It consists of 6 inch to 2 foot thick lenses of pebble-conglomerate traceable for 2500 feet.

Economic Features

Assays from samples were 0.24 percent U₃O₈ and 0.30 percent ThO₂ (chemical).

A grab sample taken from radioactive conglomerate assayed 0.11 percent U₃O₈ and 0.05 percent U₃O₈.

The average of assays from the second zone was 0.80 pounds U₃O₈ per ton.

History of Development

1956: A geological survey was completed by Sagamore Explorations Limited.

1957: Six drill holes totalling 905 feet were drilled by InSCO Mines Limited.

1958: A surficial geology survey, a radiometric survey and trenching were completed by Cody-Reco Mines Limited. Six drill holes were drilled by United MacFie Mines Limited.

1966: Three drill holes were completed by Kerr Addison Mines Limited.

1968-1969: Several trenches, pits and 14 drill holes totalling 9200 feet were completed by Acme Gas and Oil Company Limited.

1976: Kerr Addison Mines Limited completed geological, geophysical, and radiometric surveys.

1977: Kerr Addison Mines Limited drilled eight diamond drill holes.

References

Card and Lumbers (1977)

Robertson (1968a, p.70-71)

Thomson (1960a, p.19-22)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Drury Township, Drill Log Report No. 22, 23, Acme Gas and Oil Company Limited, 1968-69

Technical File No. 2.378, Acme Gas and Oil Company Limited, 1969-1971

Technical File No. 2.2229, Kerr Addison Mines Limited, 1976

Drury Township, Drill Log Report No. 30, Kerr Addison Mines Limited, 1977

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000391

ERMATINGER TOWNSHIP

BALBOA OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°36'36"N, Longitude 81°37'15"W.
Ermatinger Township.
Map Reference: OGS Map 2361.

Geology
At this showing, deformed Huronian

sedimentary rock lies unconformably on the granite basement complex. Immediately above the unconformity are outcrops of the Mississagi Formation. The Mississagi rocks consist of quartzite and arkosic beds with argillite partings and scattered thin interbeds of quartz-pebble conglomerate. The beds dip steeply to the northwest, and face in the same direction. The unconformity is exposed at a few places and is marked by a 3-foot transition zone between massive local arkose and massive pink granite.

The Mississagi Formation is locally radioactive near the granite contact.

Economic Features

The best assay was 0.042 percent U_3O_8 (chemical). Deeper drilling indicated 0.01 percent U_3O_8 and 0.03 percent ThO_2 (radiometric equivalent) over 3.0 feet (Thomson 1960a).

History of Development

1957-1959: Trenching, pitting and five drill holes were completed by Alcourt Mines Limited.

1968: Geological and geophysical exploration of the area was completed by Balboa Uranium Mines Limited.

1974-1975: An airborne geophysical survey was completed by Consolidated Morrison Explorations Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.71)
Thomson (1960a, p.20-21)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Ermatinger Township, Drill Log
Report, Alcourt Mines Limited and
Balboa Uranium Mines Limited, 1968.
Technical File No. 63.2307, Balboa
Uranium Mines Limited, 1968.
Technical File No. 2.1671,
Consolidated Morrison Explorations
Limited.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000394.

GRIGG TOWNSHIP

CANADIAN JOHNS-MANVILLE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 47°01'40"N, Longitude 80°52'54"W.
Grigg Township.
Map Reference: OGS Map 2361.

Geology

The metasediments are thought to be equivalent to those of the Elliot Lake and Hough Lake Groups. They are exposed in the central part of Grigg Township, along the Wanapitei River.

Economic Features

The first diamond drill hole graded 0.77, 0.56, and 0.47 pounds U_3O_8 per ton over 2.3, 1.0, and 4.0 feet respectively.

History of Development

1967: Airborne electromagnetic, magnetic and radiometric surveys and geological mapping were completed by Canadian Johns-Manville Company Limited.

1967-1971: Thirteen diamond drill holes totalling 4737 feet were drilled by Canadian Johns-Manville Company Limited. They also prepared a geological and airborne-geophysical report.

References

Card and Lumbers (1977)
Meyn (1972, p.32)
Robertson (1968a, p.72)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Grigg Township, Drill Log Report No.
10, Canadian Johns-Manville Company
Limited, 1967-1971.
Technical File No. 63A.548, Canadian
Johns-Manville Company Limited,
1969.
Technical File No. 63.2277A, Canadian
Johns-Manville Company Limited,
1967.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000396

HALLAM TOWNSHIP

AGGRESSIVE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°16'13"N, Longitude 81°52'58"W.
Hallam Township.
Map Reference: OGS Map 2361.

Geology

Huronian metasediments are exposed at the surface across 17 feet. They are thought to belong to the Elliot Lake Group. They comprise feldspathic quartzite, quartz-pebble conglomerate, arkose, protoquartzite, and greywacke. The radioactive zone is confined to the northwest corner of the claim and extends northward for 9000 feet.

Economic Features

Four thousand feet south of the main conglomerate-granite contact, assays from core samples yielded 0.3 percent U_3O_8 over 0.7 feet and 0.06 percent U_3O_8 (radiometric equivalent) over an unrecorded width.

History of Development

1955: Seven drill holes totalling 2260 feet were drilled by International Cobalt and Silver Mining Company Limited.
1955-1956: Five drill holes totalling 1400 feet and seven trenches were completed by Aggressive Mining Limited.
1957: Three drill holes totalling 1197 feet were drilled by Delcan Minerals Limited and Cleveland Copper Corporation.
1969: Two drill holes totalling 2504 feet were drilled by Aggressive Mining Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.85)
The Northern Miner (1969, Jan. 2, p.18, "Aggressive Mining Cuts Uranium Section")

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Hallam Township, Drill Log Report No. 11, Aggressive Mining Limited, 1955-1969.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000397.

HUTTON TOWNSHIP**ASSEMBLY OCCURRENCE****Commodity**

Uranium, thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°50'35"N, Longitude 80°59'35"W.
Hutton Township.
Map Reference: OGS Map 2361.

Geology

Uranium mineralization occurs in pyritic quartz-pebble conglomerate at or near the base of the Huronian metasedimentary sequence.

Economic Features

The best assays averaged 0.20 percent U_3O_8 and 0.08 percent ThO_2 .

History of Development

1955-1956: Geological and geophysical surveys were completed by Fano Uranium Mines Limited.
1955-1958: Several test pits and three drill holes totalling 2463 feet were completed by Assembly Mines Limited.
1967: Four drill holes totalling 1413 feet were drilled by Assembly Mines Limited.
1969: A geological survey was completed by International Mine Services Limited.
1973: An airborne geophysical survey was completed by Gulf Minerals Canada Limited.
1975: The Hanna Mining Company Limited completed a radiometric report.

References

Card and Lumbers (1977)
Meyn (1970, p.64-65)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Hutton Township, Drill Log Report No. 18, Assembly Mines Limited, 1967-1968.

Technical File No. 63A.533, International Mine Services Limited, 1969.

Technical File No. 2.1214, Gulf Minerals Limited, 1973.

Technical File No. 2.2252, Hanna Mining Company Limited, 1975

Regional Geologist's Files, Sudbury
File S63-207, Sudbury-Assembly Mines Limited

ASSEMBLY 1A OCCURRENCE**Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°52'23"N, Longitude 81°01'19"W.
Hutton Township.
Map Reference: OGS Map 2361.

Geology

Quartz-pebble conglomerate and quartzite lie on granitic rocks.

Economic Features

Three shallow holes were drilled in 1955 and all encountered interbedded conglomerate and argillite. The available assays show an average of 0.06 percent U₃O₈.

History of Development

1954-1955: Several small trenches and four drill holes totalling 21 feet were completed by Doyon, MacLeon, MacIntosh and Associates.

1966: Nine trenches and four drill holes were completed by Assembly Mines Limited.

1970: A geological survey was completed by Gui-Por Uranium Mines and Metals Limited.

References

Card and Lumbers (1977)
Meyn (1970, p.64)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto
Hutton Township, Drill Log Report No. 12 MacIntosh and Associates, 1970.
Technical File No. 63A.558, Gui-Por Uranium Mines and Metals Limited, 1970.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000400

HYMAN TOWNSHIP**NORANDA OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°23'08"N, Longitude 81°40'10"W.
Hyman Township.
Map Reference: OGS Map 2361

Geology

Uranium mineralization is associated with interbeds of quartz-pebble conglomerate in quartzite.

Economic Features

Eight showings of uranium mineralization occur on the property. Two uraniumiferous bodies are contained within the Ridge showing. The southwestern one averaged 0.28 percent U₃O₈ (chemical equivalent) across 2.0 feet. The northwestern body averaged 0.16 percent U₃O₈ across 2.1 feet for a length of 187 feet. Uranium occurs in quartzite and pebble conglomerate.

The Cabin showing, 600 feet northwest of the Ridge showing, assayed 0.17 percent U₃O₈ and 0.06 percent ThO₂ (radiometric equivalent). Samples from the Island showing assayed 0.152 percent U₃O₈ (radiometric equivalent) across 8 feet. Hilltop showing samples assayed 0.182 percent U₃O₈ (radiometric equivalent).

History of Development

1953-1954: Radiometric, geological, and geiger surveys were completed by Noranda Mines Limited.

1954: Eighteen diamond drill holes totalling 3093 feet were drilled by Noranda Mines Limited.

1976: A.E. Rose completed a radiometric survey.

1977: Two diamond drill holes totalling 1919 feet were completed by Kerr Addison Mines Limited.

Consolidated Morrison Explorations Limited drilled six diamond drill holes.

1978: One diamond drill hole totalling 3416 feet was drilled by Consolidated Morrison Explorations Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.76-77)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto

Hyman Township, Drill Log Report No. 22, 23, Consolidated Morrison Explorations Limited, 1977-1978.

Hyman Township, Drill Log Report No. 20, Kerr Addison Mines Limited, 1977.

Hyman Township, Drill Log Report No. 10, Noranda Mines Limited, 1954.

Technical File No. 63.513, Noranda Mines Limited, 1953-1954.

Technical File No. 2.2112, A.E. Rose, 1976.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000404

**AGNEW LAKE MINE
(AGNEW LAKE MINES LIMITED)**

Commodity

Uranium, thorium, and yttrium.

Radioactive Minerals

Uranothorite, monazite, and uraninite.

Location

Latitude 46°25'59"N, Longitude 81°37'30"W.
Hyman Township.
Map Reference: OGS Map 2361.

Geology

Radioactive minerals are found in quartz-pebble conglomerate, which is interbedded with sericitic quartzite and argillite. Recoverable amounts of yttrium oxide are indicated. The deposit consists of three zones, and No. 3 has been subdivided into 3A, 3B, and 3C.

Zone No. 1 occurs in quartzite. Zone No. 2 occurs in coarse-grained sericitic quartzite. Zone No. 3 occurs in parallel beds of conglomerate.

Economic Features

As of December 1, 1980, estimated mineral reserves in the proven and probable category were 5,803,000 tons at 1.0 pound U₃O₈ per ton (Kerr Addison Mines Limited, Company Annual Report).

History of Development

1954-1955: Diamond drilling totalling 36,100 feet was carried out by New Thurbois Mines Limited.

1956: Geological and ground geophysical surveys were carried out by Agnew Lakes Mines Limited.

1956-1966: Diamond drilling totalling 21,400 feet was carried out by Quebec Mattagami Minerals Limited.

1967-1968: No. 1 shaft was carried to 1400 feet. Sixty-four drill holes totalling 18,430 feet were drilled by Agnew Lake Mines Limited.

1970: A total of 8809 feet of drifting and crosscutting, 1802 feet of raising and 20,091 feet of drilling was completed by Agnew Lake Mines Limited. Work was suspended in late 1970 due to low uranium prices.

1974: The mine was dewatered to the 1750-foot level and a two year program of surface leaching began.

1976: Lateral development was up to 20,772 feet. Total diamond drilling from surface was 2120 feet, and from underground, 36,560 feet.

1977: Production resumed. At the end of 1976, 300,000 tons of ore were available for leaching.

1979: Mine development totalled 29,047 feet consisting of 25,047 feet of drives and 4000 feet of raises. The mine produced 448,000 pounds of U₃O₈.

1980: Production increased despite plans to phase out operations. The mine produced 507,000 pounds of U₃O₈.

1981-1983: Leaching operations have been continued as long as they are economic. Final termination is anticipated for early 1983.

References

Kerr Addison Mines Limited, Annual Report.

Thomson (1960a, p.22-24)

Robertson (1968a, p.74)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Hyman Township, Drill Log Report No. 13, New Thurbois Mines Limited, 1954-1955.

Hyman Township, Drill Log Report No. 18, Agnew Lake Mines Limited, 1965-1966.

Technical File No. 63A.276, Agnew Lake Mines Limited, 1956.

Regional Geologist's Files, Sudbury
File No. S63.217, New Thurbois Mines Limited.
File No. S63.751, Kerr Addison Mines Limited

EAST BAY OCCURRENCE

Commodity

Uranium and thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°25'32"N, Longitude 81°40'38"W.
Hyman Township.
Map Reference: OGS Map 2361.

Geology

Several major faults cut the area. A number of radioactive beds have been found in polymictic conglomeratic where the Sealine and Cygnet faults cut each other.

Economic Features

Two of the conglomerate beds were trenched north of John Creek. A sample assayed 3.62 pounds U_3O_8 per ton and 2.84 pounds ThO_2 per ton over 1.5 feet. Other assays from these beds were 1.08 pounds U_3O_8 per ton and 1.34 pounds ThO_2 per ton over 5.5 feet.

History of Development

1967: A radiometric and geological survey and five drill holes totalling 1084 feet were completed by East Bay Gold Limited.
 1968-1969: Four drill holes totalling 5219 feet and some trenching were completed by Monteagle Minerals Limited.
 1969-1970: Seven drill holes totalling 8000 feet were drilled by Imperial Oil Enterprises.
 1976: E.A. Rose completed a radiometric survey.
 1977: Two diamond drill holes totalling 1919 feet were drilled by Kerr Addison Mines Limited.
 Consolidated Morrison Explorations Limited drilled six diamond drill holes.
 1978: One diamond drill hole totalling 3146 feet was drilled by Consolidated Morrison Explorations Limited.

References

Card and Lumbers (1977)
 Robertson (1968a, p.76)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto

Hyman Township, Drill Log Report No. 14, East Bay Gold Limited, 1967-1969.
 Technical File No. 63.2176, East Bay Gold Limited, 1967-1968.
 Technical File No. 2.2112, E.A. Rose, 1976
 Technical File No. 2.1800, E.A. Rose, 1975

Source Mineral Deposits Record, Ontario
 Geological Survey, Toronto
 File No. 000403

MACLENNAN TOWNSHIP**PICTON OCCURRENCE (LECLERC OCCURRENCE)**

Commodity
 Uranium.

Radioactive Minerals
 Unknown.

Location

Latitude 46°40'55"N, Longitude 80°46'22"W.
 MacLennan Township.
 Map Reference: ODM Map 2009.

Geology

Radioactive quartz-pebble conglomerate is exposed at four places along the west shore of Massey Bay, Wanapitei Lake. It contains quartz and chert pebbles in an arkose matrix with sparse to profuse pyrite mineralization.

Economic Features

Representative samples taken assayed 0.01 percent U_3O_8 (radiometric equivalent). Samples taken by Picton Uranium Mines Limited gave 0.06 percent U_3O_8 (radiometric equivalent). The radioactive conglomerate was traced along the lake shore for a distance of approximately 600 feet.

History of Development

1957: A ground geophysical survey was carried out by El Pen-Rey Oil and Mines Limited.
 1959: Three drill holes totalling 532 feet were drilled by Picton Uranium Mines Limited.
 1970: An airborne geophysical survey was carried out by Tomrose Mines Limited and Kennco Explorations (Canada) Limited.
 1975: One short drill hole was drilled by M. Burton.
 1976: Trenching, geological and geophysical surveys and diamond drilling were carried out by Hollinger Mines Limited.

References

Robertson (1968a, p.78)
 Thomson (1960b, p.29-30)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto

Technical File No. 63.825, El Pen-Rey Oil and Mines Limited, 1957
 Drill Log Report No. 15, Picton Uranium Mines Limited, 1959
 Technical File No. 63.210, Tomrose Mines Limited, 1970

Source Mineral Deposits Record, Ontario
 Geological Survey, Toronto
 File No. 000409

MAY TOWNSHIP**DOMINION GULF OCCURRENCE**

Commodity
 Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°16'11"N, Longitude 81°57'23"W.
Map Reference: OGS Map 2419
May Township.

Geology

Sheared and deformed quartzite and greywacke are the host rocks for radioactive quartz-pebble conglomerate.

Economic Features

A 6-inch rusty band of conglomerate in quartzite was drilled. Assays averaged 0.07 percent U₃O₈ (radiometric equivalent).

In 1967, a single drill hole intersected radioactive quartzite and argillaceous material at 447.0 feet with 0.02 percent U₃O₈ (radiometric equivalent).

History of Development

1950: Some geological exploration and sampling was completed by A. Alexander.

1954: Surficial geological exploration and three drill holes totalling 434.1 feet were completed by Dominion Gulf Company Limited.

1967: Surficial geological exploration and a single drill hole totalling 531.0 feet were completed by McIntyre Porcupine Mines Limited.

References

Giblin et al. (1979)
Robertson (1968a, p. 79)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

May Township, Drill Log Report No. 12, McIntyre Porcupine Mines Limited, 1967.

May Township, Drill Log Report No. 13, Dominion Gulf Company Limited, 1954.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000411

PARKIN TOWNSHIP**ASSEMBLY 17A OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 46°52'37"N, Longitude 80°55'59"W.
Parkin Township.
Map Reference: OGS Map 2361

Geology

Radioactivity was observed in the main conglomerate bed, and in a weakly sheared bed of greywacke northeast of the nose of an anticlinal structure which is found in the area of the deposit.

Economic Features

Two grab samples taken by H. Meyn, Ontario Department of Mines, in 1966 and assayed radiometrically showed 0.003 percent U₃O₈ (Meyn 1970).

History of Development

1953-1954: Twelve drill holes were drilled totalling 2000 feet by Rhodes Exploration and Finance of Canada Limited.

1957: Rhodes Exploration drilled another 249 feet.

1973: An airborne geophysical survey was completed by Gulf Minerals Canada Limited.

1974: Four drill holes totalling 1965 feet were drilled by Gulf Minerals of Canada Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.80)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Parkin Township, Drill Log Report No. 15, 16, Rhodes Exploration, 1953-1954, and 1957.

Technical File No. 2.1214, Gulf Minerals of Canada Limited, 1973.
Drill Log Report No. 29, Gulf Minerals of Canada Limited 1974.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000413.

PORTER TOWNSHIP**BREWIS-WHITE OCCURRENCE**

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 46°24'58"N, Longitude 81°44'19"W.
Porter Township.
Map Reference: OGS Map 2361.

Geology

Radioactivity is associated with conglomerate and quartzite of the Huronian Supergroup.

Economic Features

Grab samples of the conglomerates assayed 0.006 percent and 0.20 percent U₃O₈ (radiometric equivalent).

History of Development

1954: A geological and radiometric survey was completed by Brewis and White Limited.
1955: A geological and ground radiometric survey was carried out by Agnew Lake Mines Limited.
1969: Ground and airborne geophysical surveys were carried out by Canadian Johns-Manville Company Limited.
1974: An airborne geophysical survey was completed by Consolidated Morrison Explorations Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A.276, Agnew Lake Uranium Mines Limited, 1955-1956
Technical File No. 63A.276, New Thurbois Mines Limited, 1955-1956.
Technical File No. 63.2517, Canadian Johns-Manville Company Limited, 1969.
Technical File No. 2.1671, Consolidated Morrison Explorations Limited, 1974.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000414.

CHEMICAL RESEARCH OCCURRENCE**Commodity**

Uranium and thorium.

Radioactive Minerals

Unknown.

Location

Latitude 46°22'04"N, Longitude 81°42'12"W.
Porter Township.
Map Reference: OGS Map 2361.

Geology

Radioactive quartz-pebble conglomerate occurs adjacent to a volcanic complex and a ridge of quartzite and quartz-pebble conglomerate, that is much contorted and faulted.

Economic Features

The radioactive conglomerate was sampled from 11 trenches at intervals of 240 feet in a direction N55E over a width of 12 feet. The average assay was 0.17 percent U₃O₈.

History of Development

1954: Geological mapping, a scintillometer survey and trenching were carried out by Chemical Research Corporation (Canada) Limited.
1968-1969: Drilling of eight holes totalling 1674 feet was completed by Reactor Uranium Mines Limited.
1975: An airborne geophysical survey was completed by Consolidated Morrison Explorations Limited.
1977: Five diamond drill holes were drilled by Consolidated Morrison Explorations Limited.
1979: The company's name was changed to Consolidated Reactor Uranium Mines Limited.

References

Card and Lumbers (1977)
Thomson (1960a, p.24-25)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63A 200, Chemical Research Corporation (Canada) Limited, 1954.
Porter Township, Drill Log Report No. 14, Reactor Uranium Mines Limited, 1968-1969.
Technical File No. 63.2423, Chemical Research Corporation (Canada) Limited, 1968.
Technical File No. 2.2613, D.S. Robertson and Associates Limited.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000402.

HUNTER LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°22'50"N, Longitude 81°41'55"W.
Porter Township.
Map Reference: OGS Map 2361.

Geology
Mineralization is associated with the quartz-pebble conglomerate. The bed has a maximum thickness of 7 feet. Mineralization cannot be traced continuously for more than 200 feet over a total length of 3100 feet.

Economic Features
A grab sample assayed 0.50 percent and 0.20 percent U₃O₈.

History of Development
1954: A geological and geophysical survey was completed by the Chemical Research Corporation (Canada) Limited.
1961: Surface sampling was done by Ontario Department of Mines geologist R.M. Ginn.
1966-1967: A ground geophysical survey was completed by Reactor Uranium Mines Limited.
1968-1969: A geological survey was completed by Watts, Griffis, and McQuat Limited.
1974: An airborne geophysical survey was carried out by Consolidated Morrison Explorations Limited.
1975: An airborne geophysical survey was completed by Amax Exploration Incorporated.

References
Card and Lumbers (1977)
Robertson (1968a, p.90-91)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto
Technical File No. 63A 200, Chemical Research Corporation (Canada) Limited, 1954.
Technical File No. 63.2211 Reactor Uranium Mines Limited 1966-1967.
Technical File No. 63.2211A Broulan Reef Mines Limited 1967.
Technical File No. 63.2423, Watts, Griffis, and McQuat Limited, 1968-1969.

Technical File No. 2.1671
Consolidated Morrison Explorations Limited, 1974.
Technical File No. 2.1618, Amax Exploration Incorporated, 1975.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000415.

NEW MYLAMAQUE (MYMAR) OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°25'10"N, Longitude 81°42'50"W.
Porter Township.
Map Reference: OGS Map 2361.

Geology
Radioactivity was observed along an east-trending radioactive zone of quartz-pebble conglomerate.

Economic Features
Grab samples from this zone assayed 0.18 and 0.20 percent U₃O₈.

History of Development
1954: A geological and ground geophysical survey was carried out by New Mylamaque Explorations Limited
1969: An airborne geophysical survey was completed by Canadian Johns-Manville Company Limited.
Four drill holes totalling 2634 feet were drilled for Projex Limited.
1974: Consolidated Morrison Explorations Limited carried out an airborne geophysical survey.

References
Card and Lumbers (1977)
Robertson (1968a, p.86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario Geological Survey, Toronto
Technical File No. 63.501, New Mylamaque Explorations Limited, 1954.
Technical File No. 63.2517, Canadian Johns-Manville Company Limited, 1969.

Technical File No. 2.1671,
Consolidated Morrison Explorations
Limited, 1974.
Drill Log Report No. 19, Projex
Limited, G.L. Phelan, 1969.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000417.

ROBERTS TOWNSHIP

NORDIC RESOURCES PROSPECT

Commodity
Uranium and thorium.

Radioactive Minerals
Uraninite.

Location
Latitude 46°54'57"N, Longitude 81°06'04"W.
Roberts Township.
Map Reference: ODM Map 2212.

Geology
Mineralization occurs in Proterozoic
argillite containing quartz-pebble
conglomerate. The main showing is
approximately 90 by 460 m.

Economic Features
According to assays by Amax Exploration
Incorporated, the area of the main showing
has a grade potential of 0.025 percent U₃O₈.
The best drill intersection by Amax assayed
0.06 percent U₃O₈ and 0.02 percent ThO₂
over 1.4 m of core.

History of Development
1975: Geological mapping and a radiometric
survey were carried out by E.A. Rose.
1976: Erana Mines Limited carried out
geological and radiometric surveys.
TX Resources Limited completed a
geological survey.

References
Meyn (1971)
Thomson (1960a)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 2.1605, Amax
Exploration Incorporated, 1974
Mineral Deposits Inventory Record, Ontario
Geological Survey, Toronto
File No. S0087.

ROBERTS LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°57'55"N, Longitude 81°06'12"W.
Roberts Township.
Map Reference: ODM Map 2212.

Geology
Sedimentary formations of quartzite,
greywacke, arkose and pebble to boulder
conglomerate are present. Scattered
conglomerate layers are found within the
quartzites. These layers consist of granitic
pebbles in a quartzite matrix.

Economic Features
Several drill holes were drilled in the
Huronian metasediments. Radioactivity
readings were 0.010 mR/h. Other assays were
0.001 percent U₃O₈ (radiometric equivalent)
and 0.006 percent U₃O₈ (radiometric
equivalent).

History of Development
1954: A geological survey was completed by
Dyno Mines Limited.
1954-1955: A geological survey and surface
sampling were carried out by Elmridge Mines
Limited.
1966-1973: H.C. Humphries and Assembly
Mines Limited drilled 42 drill holes totalling
10,584 feet.
1973: An airborne geophysical survey was
carried out by Gulf Minerals Canada Limited.

References
Meyn (1971)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 63A229, Elmridge
Mines Limited, 1954
Technical File No. 63A204, Dyno
Mines Limited, 1954
Roberts Township, Drill Log Report
No. 14 and 16, Assembly Mines
Limited, 1966-68.
Roberts Township, Drill Log Report
No. 15 and 20, H.C. Humphries, 1967-
73.
Technical File No. 2.1214, Gulf
Minerals Canada Limited, 1973.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001104.

SHAKESPEARE TOWNSHIP

ALEXANDER OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°17'02"N, Longitude 81°56'31"W.
Shakespeare Township.
Map Reference: ODM Map 2313.

Geology
The Early Precambrian granitic basement is in contact with Huronian metasediments. Adjacent to the contact zone, a 15-foot width of radioactive quartz-pebble conglomerate is exposed.

Economic Features
Samples from the main conglomerate bed assayed 0.13 percent U₃O₈.

History of Development
1968-1969: A geological and geophysical ground survey and several drill holes were completed by Aggressive Mining Limited.
1969: A geological and ground geophysical survey, and two drill holes totalling 1337 feet were completed by Moncrieff Uranium Mines Limited.
1976: Kerr Addison Mines Limited performed geophysical and geological surveys.

References
Card and Palonen (1976, p.43)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Shakespeare Township, Drill Log
Report No. 14, Moncrieff Uranium
Mines Limited, 1969.
Technical File No. 2.2294, Kerr
Addison Mines Limited, 1976.
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001103.

BROULAN REEF OCCURRENCE (SUTHERLAND CREEK OCCURRENCE)

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 46°22'27"N, Longitude 81°48'58"W.
Shakespeare Township.
Map Reference: ODM Map 2313.

Geology
Radioactivity was found in oligomictic quartz-pebble lenses in the Matinenda Formation. The lenses range in thickness from a few inches to 3 feet.

Economic Features
The average assay was 0.02 percent U₃O₈.

History of Development
1947: Two drill holes totalling 3416 feet were drilled by Falconbridge Nickel Mines Limited.
1953-54: A geological survey was completed by Dominion Gulf Company Limited.
1954: A geological and ground geophysical survey was completed by Dominion Gulf Company Limited.
1968: A geophysical survey was completed by Broulan Reef Mines Limited.
1968-69: An airborne geophysical survey was completed by Broulan Reef Mines Limited. Eighteen diamond drill holes totalling 13,086 feet were drilled by Broulan Reef Mines Limited.
Two drill holes totalling 1294 feet were drilled by Broulan Reef Mines Limited.
1969-1970: A single drill hole to 2070 feet was drilled by Broulan Reef Mines Limited.

References
Card and Palonen (1976, p.45)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Shakespeare Township, Drill Log
Report No. 17, Falconbridge Nickel
Mines Limited, 1947.
Shakespeare Township, Drill Log
Report No. 10, 11, 16, Broulan Reef
Mines Limited 1968-1970.
Technical File No. 63A203, Dominion
Gulf Company Limited, 1954.
Technical File No. 63.502, Dominion
Gulf Company Limited, 1954.

Technical File No. 63.2211, Broulan Reef Mines Limited, 1968.
Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 001102.

DOMINION GULF OCCURRENCE

Commodity

Uranium and thorium.

Radioactive Minerals

Pitchblende, brannerite, and thucolite.

Location

Latitude 45°20'48"N, Longitude 81°50'45"W.
Shakespeare Township.
Map Reference: ODM Map 2313.

Geology

The sedimentary rocks here consist of quartzite with local interbeds of quartz-pebble conglomerate and are intruded by large and small bodies of diabase, diorite, and gabbro. Five zones of radioactivity were located in the metasediments, four of these in quartz-pebble conglomerate beds associated with quartzite.

Economic Features

The two most important zones occur on the south shore of Agnew Lake. Both are exposed in a few places over strike lengths of 150 to 200 feet. Assays recorded were up to 0.054 percent U_3O_8 and 0.9 percent ThO_2 over widths of 8 inches to 2 feet. In the other areas, radioactivity was up to 0.03 percent U_3O_8 and 0.05 percent ThO_2 (Resident Geologists Files, Ontario Ministry of Natural Resources, Sudbury).

History of Development

1954: Geological and ground surveys were completed by Dominion Gulf Company Limited.

1961: Two drill holes totalling 720 feet were drilled by Falconbridge Nickel Mines Limited.

1968: A ground geophysical survey was completed by Broulan Reef Mines Limited.

1974: An airborne geophysical survey was completed by Consolidated Morrison Explorations Limited.

References

Card and Palonen (1976, p.44)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63.203, Dominion

Gulf Company Limited, 1954.

Technical File No. 63.502, Dominion Gulf Company Limited, 1954-1955.

Technical File No. 63.2211, Broulan Reef Mines Limited, 1968.

Technical File No. 63.2339, Pick Mines Limited, 1969.

Technical File No. 2.1671, Consolidated Morrison Explorations Limited, 1974

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000423

MCDONALD BAY OCCURRENCE

Commodity

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 46°20'48"N, Longitude 81°50'06"W.
Shakespeare Township.
Map Reference: ODM Map 2313.

Geology

The most important discovery was found on N1/2, lot 2, concession IV. It consists of several parallel conglomerate beds that strike northeast and dip 70°NW. Diamond drilling revealed quartz-pebble conglomerate lenses within the sections of interbedded arkose, pelite, volcanic rocks, and conglomerate.

Economic Features

Assays ranged from traces to 0.14 percent U_3O_8 (radiometric equivalent) over widths of 1 to 3 feet. Other assays averaged 0.10 percent U_3O_8 across 2.0 feet.

History of Development

1953-1954: Geological and geophysical surveys were completed by Shakespeare Uranium Mines Limited.

1954: Thirty-one drill holes totalling 7294 feet were drilled by Shakespeare Uranium Mines Limited.

References

Card and Palonen (1976, p.45-46)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63A 518,

Shakespeare Uranium Mines, 1954.

Shakespeare Township, Drill Log
Report No. 18, Shakespeare Uranium
Mines Limited, 1954-1955.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000425.

STETHAM TOWNSHIP

JONSMITH OCCURRENCE (MCKINNON OCCURRENCE)

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 47°45'54"N, Longitude 81°39'07"W.
Stetham Township.
Map Reference: OGS Map 2361.

Geology
Granitic rocks with partially assimilated
inclusions of mafic metavolcanics are cut by
northwest- and northeast-trending diabase
dikes and late northwest-trending faults.
Radioactivity was found in pegmatitic and
mafic rocks on the surface.

Drilling was concentrated on the southern
shore of the west arm of Kenetogami Lake. It
was determined that radioactivity is
associated with widespread carbonization and
the introduction of brown hematite into the
mafic rocks.

Economic Features

Two diamond drill hole intersections
returned assays of 0.6 pounds U_3O_8 per ton
over 30 feet and 1.6 pounds U_3O_8 per ton
over 16.8 feet (The Northern Miner, 1976,
Sept. 9, p.22).

In 1977, Beach Gold Mines Limited drilled
five holes and outlined the A Zone, which
extends for 1600 feet. The average assay is
1.33 pounds U_3O_8 per ton.

Ground radiometric surveys have
indicated two new zones. The B Zone, 1100
feet west of the A Zone, has an indicated
length of 1000 feet and is open to the north.
The C Zone, about 1/2 mile west of the B
Zone is approximately 1400 feet long and
open to the north (The Northern Miner, 1977,
May 15, p.5).

History of Development

1967: Ground geophysical and geological
surveys, prospecting, pitting and grab
sampling were carried out by Jonsmith
Uranium Mines Limited.

1968: Pitting, trenching and 14 diamond d
holes totalling 5303 feet were completed
Jonsmith Uranium Mines Limited.

1970: Pitting and diamond drilling were
carried out by Jonsmith Uranium Mines
Limited.

1976: An airborne radiometric survey was
carried out by Beach Gold Mines Limited.

1977: Five diamond drill holes totalling 19
feet were drilled by Beach Gold Mines
Limited.

References

Card and Lumbers (1977)
Robertson (1968a, p.83-84)
The Northern Miner (1977, May 12, p.5,
"Beach Gold Tests Zone A on "U" Prospe

Ontario Ministry of Natural Resources Fi
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.3283, Jonsmi
Uranium Mines Limited, 1967.

Technical File No. 63.3279, Jonsmi
Uranium Mines Limited, 1968.

Stetham Township, Drill Log Report
No. 11, 12, 13 Jonsmith Uranium
Mines Limited, 1968-1970.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000426.

STOBIE TOWNSHIP

STOBIE TOWNSHIP OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 47°01'48"N, Longitude 80°46'33"W
Stobie Township.
Map Reference: ODM Map 2238.

Geology

Uranium mineralization is present in the
Matinenda sedimentary rocks interbedded
with sheared, rusty, argillaceous schist.
Mineralization is also found in an angular
quartz conglomerate with a quartz
greywacke matrix.

Economic Features

An assay of sheared argillite returned
0.08 percent U_3O_8 . Analysis of the
conglomerate matrix showed 0.021 percent
 U_3O_8 was present.

History of Development

1967: Geophysical and geological surveys were carried out by Canadian Johns-Manville Company Limited.

1976: Stripping, trenching and assaying were completed by J.A. Pollock.

1977: Two drill holes totalling 628 feet were completed by the Grandad-Warren-Consolidated Professor joint option. Further trenching was carried out by J.A. Pollock.

References

Meyn (1972, p.32)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 63A.548, Canadian Johns-Manville Company Limited, 1969.

Technical File No. 63.2277, Canadian Johns-Manville Company Limited, 1967.

Stobie Township, Drill Log Report No. 10, Grandad-Warren Consolidated Professor, 1977.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000174.

Economic Features

Assays from the conglomerate ranged from 0.001 percent to 0.27 percent U_3O_8 .

History of Development

1953-1954: Discovered by T. Saville and staked by Normingo Mines Limited.

1954: Surface clearing, a geophysical survey, and three drill holes were completed by Harrison Minerals Limited.

1954-1955: A geological and radioactivity survey and nine diamond drill holes totalling 1579 feet were drilled by H. Hibbert Mines Limited.

1967: Three drill holes totalling 3043 feet were drilled by Canadian Johns-Manville Company Limited.

1968-1969: An airborne geophysical survey was carried out by MacRae Mining Corporation Limited for Henry Last and Associates.

1971: Nine drill holes totalling 1540 feet were drilled by Canadian Johns-Manville Company Limited.

1976: Six diamond drill holes were drilled by Aggressive Mining Limited.

1978: Aggressive Mining Limited completed a scintillometer survey.

References

Thomson (1960a, p.29)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario Geological Survey, Toronto

Turner Township, Drill Log Report No. 10, H. Hibbert Mines Limited, 1954.

Drill Log Reports No. 12, 13, 15, Canadian Johns-Manville Company Limited, 1967, 1971.

Technical File No. 63.2275, Canadian Johns-Manville Company Limited, 1968-1969.

Technical File No. 63.2542, MacRae Mining Corporation Limited, 1969.

Drill Log Report No. 17, MacRae Mining Corporation Limited, 1969.

Technical File No. 2.2715, Aggressive Mining Limited, 1978.

Technical File No. 2.2242, Aggressive Mining Limited, 1976.

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000427.

TURNER TOWNSHIP**HARRISON OCCURRENCE****Commodity**

Uranium, thorium, and zirconium.

Radioactive Minerals

Unknown.

Location

Latitude 47°05'13"N, Longitude 80°34'41"W.
Turner Township.

Map Reference: ODM Map 2260.

Geology

Quartz-pebble conglomerate lies on basement rocks and is interbedded with argillite, pyritic quartz-pebble conglomerate, green quartzite, and greywacke. A radioactive bed of pyritic "microconglomerate" is approximately 30 feet thick and is exposed at intervals over a length of 2 miles. The rocks are folded and dips range from 25 to 70 degrees north, east and south.

MINOR URANIUM AND THORIUM OCCURRENCES OF SUDBURY DISTRICT

BALDWIN TOWNSHIP

DOMINION GULF OCCURRENCE

Location

Latitude 46°21'21"N, Longitude 81°42'50"W.
Baldwin Township.

References

Giblin et al. (1979)
Thomson (1953)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 63A.254, Dominion
Gulf Company Limited, 1955.
Technical File No. 63.1049, Evenlode
Gold Limited, 1960.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000382.

Regional Geologist Files, Sudbury
File No. S63.26.

UNITED COBALT OCCURRENCE

Location

Latitude 46°20'06"N, Longitude 81°47'56"N.
Baldwin Township.

Remarks

Two parallel beds of radioactive quartz-
pebble conglomerate approximately 175 feet
apart have been traced intermittently in a
northeasterly direction for 1.5 miles.

Extensive drilling by United Cobalt Mines
Limited established the presence of low
radioactivity. No assays were available.

References

Canadian Mines Handbook (1972-1973, p.379)
Robertson (1968a, p.85)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Baldwin Township, Drill Log Report
No. 35, United Cobalt Mines Limited,
1956.

Baldwin Township, Drill Log Report
No. 27, Rowan Consolidated Mines
Limited, 1967.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000387.

BIGWOOD TOWNSHIP

FRENCH RIVER OCCURRENCE (SOUTH
SIDE OF MAIN CHANNEL)**Location**

Latitude 46°10'12"N, Longitude 80°13'01"W.
Bigwood Township.

Remarks

Crystals and seams of allanite occur with
abundant red alkalic feldspar in an intensely
hematized pegmatite dike.

References

Lumbers (1975)

CASSIDY TOWNSHIP

AUBINADONG RIVER OCCURRENCE

Location

Latitude 47°15'51"N, Longitude 83°18'20"W.
Cassidy Township.

Remarks

The occurrence is situated in a cliff face
on the east side of the Aubinadong River. A.
Zeemel of Gunnar Mines Limited, described
the occurrence as pitchblende filling
fractures in pink medium-grained granite.
The main fractures are 3/8 to 1/2 inch wide
and strike northeast. Tiny subsidiary
fractures are coated in places with yellow
secondary uranium oxides.

References

Thurston et al. (1977)

CLARY TOWNSHIP

NORANDA OCCURRENCE (MARCIA LAKE
OCCURRENCE)**Location**

Latitude 47°02'15"N, Longitude 80°29'46"W.
Clary Township.

Remarks

The host rocks for radioactive pebble-
conglomerate are argillite and greywacke.

References

Card et al. (1973, p.115)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Clary Township, Drill Log Report No.
10, Noranda Exploration Company
Limited, 1967.

Technical File No. 63.2524, MacRae
Mining Corporation Limited, 1969.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000393.

radioactive zone. A grab sample taken by
H.D. Meyn, Ontario Department of Mines,
assayed 0.003 percent U_3O_8 .

References

Meyn (1971, p.39)

Ontario Ministry of Natural Resources Files
Mineral Deposits Inventory Record, Ontario
Geological Survey, Toronto
File No. S0086.

COLLINS AND CHEWETT TOWNSHIPS

NEMEGOSENDA LAKE OCCURRENCE

Location

Latitude $48^{\circ}00'20''N$, Longitude $83^{\circ}06'30''W$.
Collins and Chewett Township.

Remarks

The carbonatite complex is composed of
incomplete rings of various alkalic syenites
which have intruded a sequence of tonalitic
and monzonitic rocks of the Shawmere
anorthosite complex. Parsons (1961) indicated
that three fenite zones occur from the outer
perimeter inward: ijolite fenite, red alkalic
fenite, and pyroxenitic fenite. The red
alkalic fenite is consistently radioactive.
Uranium assays were in the range of 0.02 to
0.03 percent U_3O_8 .

References

Parsons (1961)
Thurston et al. (1977)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Chewett Township, Drill Log Report
No. 10, Dominion Gulf Company
Limited, 1955-1956.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001108.

CREELMAN TOWNSHIP

NORTH CREELMAN OCCURRENCE

Location

Latitude $46^{\circ}57'21''N$, Longitude $81^{\circ}04'13''W$.
Creelman Township.

Remarks

Uranium mineralization occurs in
greywacke and pebble-conglomerate of the
Mississagi Formation. Drilling intersected a

CURTIN TOWNSHIP

BRIDGES OCCURRENCE

Location

Latitude $46^{\circ}09'00''N$, Longitude $81^{\circ}34'45''W$.
Curtin Township.

Remarks

A radioactive occurrence was discovered
at Howrey Creek, 12 miles southeast of
Espanola, in polymictic conglomerate of the
Gowganda Formation.

References

Card (1975)
Robertson (1968a, p.85)

DEMOREST TOWNSHIP

NORANDA OCCURRENCE (YORSTON LAKE OCCURRENCE)

Location

Latitude $47^{\circ}02'27''N$, Longitude $80^{\circ}35'15''W$.
DeMorest Township.

Remarks

Radioactivity is associated with quartz-
pebble conglomerate interbedded with
quartzite.

References

Card and Lumbers (1977)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Demorest Township, Drill Log Report
No. 10, Noranda Exploration Company
Limited, 1967.

Technical File No. 63.2277, Canadian-
Johns Manville Company Limited,
1967-1968.

Technical File No. 63.2542, MacRae
Mining Corporation Limited, 1969.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000313

FRALECK TOWNSHIP

INGAMAR OCCURRENCE

Location

Latitude 46°57'32"N, Longitude 80°51'36"W.
Fraleck Township.

Remarks

Locally the radioactive zones occur in quartzite and quartz-pebble conglomerate of the Mississagi Formation of the Hough Lake Group.

References

Meyn (1971, p.8-13)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2277, Canadian
Johns-Manville Company Limited,
1967.

Technical File No. 2.2307, M. LeFort,
1976.

Fraleck Township, Drill Log Report
No. 10, Canadian Johns-Manville
Company Limited.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001142.

GALLAGHER TOWNSHIP

BORDEN LAKE OCCURRENCE

Location

Latitude 47°48'55"N, Longitude 83°18'55"W.
Gallagher Township.

Remarks

Radioactivity, associated with gneissic to massive granitic rocks, is four to five times the background. The rocks strike S33E and dip 10°N. The radioactivity seems to be due to thorium.

References

Thurston et al. (1977, p.261)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000395.

HARROW TOWNSHIP

STENCIL OCCURRENCE

Location

Latitude 46°08'21"N, Longitude 82°01'20"W.
Harrow Township.

Remarks

Radioactivity is associated with a dike in quartzite.

The following assays were obtained from three holes: 0.04 percent U₃O₈ over 5.2 feet, 0.06 percent U₃O₈ over 5.5 feet and 0.02 percent U₃O₈ over 3.0 feet (Resident Geologist's Files, Ontario Ministry of Natural Resources, Sudbury)

References

Card and Lumbers (1977)

Robertson et al. (1972)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Harrow Township, Drill Log Report
No. 11, V. Stencil, 1954.

HYMAN TOWNSHIP

RICHORE OCCURRENCE

Location

Latitude 46°25'02"N, Longitude 81°39'10"W.
Hyman Township.

Remarks

The occurrence is underlain by the Birch Lake granite batholith and by metasediments. There is a series of northeasterly trending major faults in the area. Three zones of radioactivity were detected in the underlying intrusive formations and in the vertical fracture zones in the granite batholith.

Shallow diamond drilling revealed 10 radioactive zones. The best assays were 0.08 and 0.12 pounds U₃O₈ per ton.

References

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2176, East Bay
Gold Mines Limited, 1967.

Hyman Township, Drill Log Report No.
16, Richore Gold Mines Limited, 1969.

Technical File No. 2.50, Richore Gold
Mines Limited.

Technical File No. 2.2112, E.A. Rose, 1976
 Technical File No. 2.1996, E.A. Rose, 1975
 Technical File No. 2.1800, E.A. Rose, 1975.
 Technical File No. 2.1671,
 Consolidated Morrison Explorations Limited.
 Technical File No. 63A.200, Chemical Research Corporation (Canada) Limited.

Energy, Mines and Resources Canada Files
 Mineral Development Sector, Department of Energy, Mines and Resources, Ottawa
 National Mineral Inventory, File U 14, 41/1/5, Richore, June 1971.

LANGLOIS TOWNSHIP

KRAM-GLOWASKI OCCURRENCE

Location

Latitude 47°25'05"N, Longitude 83°12'11"W.
 Langlois Township.

Remarks

A band of highly folded Early Precambrian greywacke strikes northeast. The greywacke is cut by quartz veins which carry pyrite, pyrrhotite, and sphalerite.

Seven samples gave assays up to 0.09 percent U_3O_8 (radiometric equivalent).

References

Robertson (1968a, p.84-85)
 Thurston et al. (1977)

Ontario Ministry of Natural Resources Files
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 000429.

MACLENNAN TOWNSHIP

CHEVRETTE OCCURRENCE

Location

Latitude 46°39'13"N, Longitude 80°44'45"W.
 MacleNNan Township.

Remarks

Radioactivity occurs in granite. Grab samples collected by M.E. Chevrette averaged 0.08 percent U_3O_8 .

In 1970 an airborne geophysical survey was completed by Tomrose Mines Limited and Kennco Explorations (Canada) Limited.

References

Card and Lumbers (1977)
 Robertson (1968a, p.85)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario Geological Survey, Toronto
 Technical File No. 63.210, Tomrose Mines Limited, 1970.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 000407.

MASSEY BAY OCCURRENCE

Location

Latitude 46°39'25"N, Longitude 80°46'09"W.
 MacleNNan Township.

Remarks

Radioactivity was encountered in quartz-pebble conglomerate along the west shore of Massey Bay. The conglomerate lies with great angular unconformity on pre-Huronian greywacke. Quartz-pebble conglomerate and quartzite are pyritic and weakly radioactive. The only assay showed 0.009 percent U_3O_8 (radiometric equivalent).

References

Robertson (1968a, p.86)
 Thomson (1960b, p.28)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario Geological Survey, Toronto
 MacleNNan Township, Drill Log Report No. 16, Nalgar Nickel Mines Limited, 1957.
 MacleNNan Township, Drill Log Report No. 18, Pan Canadian Development Company Limited, 1961.
 MacleNNan Township, Drill Log Report No. 19, International Nickel Company of Canada Limited, 1967-1968.
 Technical File No. 63.210, Tomrose Mines Limited, 1970.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 000408.

MARCONI TOWNSHIP

CANADIAN JOHNS-MANVILLE OCCURRENCE

Location

Latitude 47°05'43"N, Longitude 80°43'57"W.
 Marconi Township.

Remarks

The contact between the Superior and Southern structural provinces occurs in Marconi Township. Diamond drilling showed that there are two separate radioactive occurrences in two different settings. The southwestern occurrence is found in pegmatite dikes. The northeastern occurrence is found in quartz-pebble conglomerate.

References

Card and Lumbers (1977)
Card et al. (1973, p.110-118)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2277, Canadian
Johns-Manville Company Limited,
1967-1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000386.

CAN-FER OCCURRENCE**Location**

Latitude 47°06'07"N, Longitude 80°46'26"W.
Marconi Township.

Remarks

Uranium mineralization was found in pegmatite dikes along the eastern shore of an unnamed lake near the centre of the Marconi Township. Assays of samples were 0.8 pounds U_3O_8 per ton.

References

Card et al. (1973, p.114)
Robertson (1968a, p.78)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2277, Canadian
Johns-Manville Company Limited,
1969.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000410.

MCNISH TOWNSHIP**SAVILLE OCCURRENCE****Location**

Latitude 46°46'50"N, Longitude 80°18'40"W.
McNish Township.

Remarks

Uranium occurs in pyritiferous quartz-pebble conglomerate of the Mississagi Formation.

References

Dressler (1979, p.81-82)

MONCRIEFF TOWNSHIP**HOLLINGER RADIOACTIVITY OCCURRENCE****Location**

Latitude 46°46'14"N, Longitude 81°36'12"W.
Moncrieff Township.

Remarks

Most of the radioactivity is confined to the Lorrain Formation.

References

Card and Lumbers (1977)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Moncrieff Township, Drill Log Report
No. 11, Nickel Rim Mines Limited.
Moncrieff Township, Drill Log Report
No. 15, Hollinger Mines Limited,
1976.

Technical File No. 2.2272, Hollinger
Mines Limited, 1977.
Technical File No. 2.2484, Hollinger
Mines Limited, 1978.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001139.

NAIRN TOWNSHIP**DEAUVILLE OCCURRENCE****Location**

Latitude 46°16'52"N, Longitude 81°37'23"W.
Nairn Township.

Remarks

Mineralization occurs in quartzite. The radioactive zone is 45 m wide. The best drill hole intersection was 0.05 percent U_3O_8 and 0.03 percent ThO_2 over 1.5 m.

References

Card and Lumbers (1977)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Nairn Township, Drill Log Report No.
14, Deauville Explorations Limited,
1968.

Technical File No. 63.2200, Deauville
Explorations Limited, 1967.

Mineral Deposits Inventory Record, Ontario
Geological Survey, Toronto
File No. S0128.

PORTER TOWNSHIP

MACFIE OCCURRENCE

Location

Latitude 46°22'06"N, Longitude 81°49'02"W.
Porter Township.

Remarks

A ground geological survey was completed
on a zone following a faulted contact
between quartz-pebble conglomerate and
feldspathic quartzite. The uraniferous zone is
about 3000 feet long and 200 feet wide.

In 1954, geological and geophysical
surveys were completed by MacFie
Explorations Limited.

In 1955, four drill holes totalling 1104
feet were drilled by Mid-North Engineering
Services Limited.

References

Robertson (1968a, p.86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Porter Township, Drill Log Report No.
10, Mid-North Engineering Services
Limited, 1955.

Technical File No. 63.499, Gardiner,
Low, and Morrow, 1954.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000416.

PENNBEC OCCURRENCE

Location

Latitude 46°26'07"N, Longitude 81°41'59"W.
Porter Township.

Remarks

A radioactive zone was located in
conglomerate interbedded with quartzite.
The conglomerate beds, as a result of
faulting, occur in lenses and patches and
cannot be traced for more than 100 feet.

References

Card and Lumbers (1977)

Robertson (1968a, p.86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Porter Township, Technical File No.
63A.276, Agnew Lake Mines Limited
and New Thurbois Mines Limited,
1955.

Porter Township, Drill Log Report No.
13, Pennebec Mining Company
Limited, 1966.

Technical File No. 2.1671,
Consolidated Morrison Explorations
Limited, 1974.

WADGE OCCURRENCE

Location

Latitude 46°22'00"N, Longitude 81°41'32"W.
Porter Township.

Remarks

A small area of anomalous radioactivity
occurs in rusty, weathered conglomerate
close to its contact with Mississagi quartzite
and a large fault which crosses the property.

References

Card and Lumbers (1977)

Robertson (1968a, p.86)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.500, Wadge
Mines Limited, 1954.

Porter Township, Drill Log Report
Nos. 15 and 16, Hanover Explorations
Limited.

SEAGRAM TOWNSHIP

DENISON OCCURRENCE

Location

Latitude 47°04'34"N, Longitude 80°33'03"W.
Seagram Township.

Remarks

In 1969, Denison Mines Limited drilled
three diamond-drill holes totalling 5170 feet.
The first hole started in the transition zone
between the Bruce and Mississagi
Formations. In the last 200 feet some
radioactive rock units were encountered.

The second hole started in the Gowganda Formation, cut a 6-foot minette dike at 180 feet and entered the Mississagi Formation at 331 feet. It continued in quartzite, greywacke, and quartz-pebble conglomerate of the Mississagi Formation to a depth of 1361 feet. Radioactive zones were encountered in the last 200 feet (Card et al. 1973).

References

Card et al. (1973, p.114)

SHAKESPEARE TOWNSHIP

BLUE OCCURRENCE

Location

Latitude 46°17'30"N, Longitude 81°51'45"W.
Shakespeare Township.

Remarks

According to Map 2361 (Card and Lumbers 1977), this occurrence is situated near the contact of the Salmay Lake Formation and the Matinenda Formation.

Radioactivity was found to average 0.003 percent U_3O_8 (radiometric equivalent).

References

Card and Palonen (1976)

Ontario Ministry of Natural Resources Files Assessment Files Research Office, Ontario Geological Survey, Toronto

Technical File No. 23718, P.G. Blue, 1981.

DELCAN OCCURRENCE

Location

Latitude 46°16'59"N, Longitude 81°52'42"W.
Shakespeare Township.

Remarks

Interbedded sandstone and pelite of the Matinenda Formation are exposed in lots 8 and 9, concession 1, Shakespeare Township.

In 1956 and 1957, Delcan Minerals Limited diamond drilled four holes, three of which totalled 1297 feet in lots 8 and 9 Shakespeare Township. In one diamond-drill hole, an assay of 0.63 percent U_3O_8 over a length of 1.5 feet of pyritic sandstone was obtained.

References

Card and Palonen (1976, p.42)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000422

PICK OCCURRENCE

Location

Latitude 46°21'29"N, Longitude 81°52'44"W.
Shakespeare Township.

Remarks

Four drill holes, drilled from the ice on Agnew Lake, intersected quartz-pebble conglomerate and pelite of the Matinenda Formation at the 210-foot level. Radioactive lenses were pyritic and geiger counter readings were up to three times the background.

References

Card and Palonen (1976, p.44)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Shakespeare Township, Drill Log
Report No. 12 and 13, Falconbridge
Nickel Mines Limited, 1968.

Technical File No. 63.2339, Pick Mines
Limited, 1968.

Technical File No. 63.2211, Broulan
Reef Mines Limited, 1968-1969.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No 001101.

SATELLITE OCCURRENCE

Location

Latitude 46°21'47"N, Longitude 81°49'23"W.
Shakespeare Township.

Remarks

A scintillometer survey and drilling in northern Shakespeare Township located one radioactive zone 30 feet wide in Huronian argillaceous quartzite.

References

Card and Palonen (1976, p.45)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2211, Broulan
Reef Mines Limited, 1968

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000424

SCHLESINGER OCCURRENCE

Location

Latitude 46°17'02"N, Longitude 81°50'31"W.
Shakespeare Township.

Remarks

A slightly radioactive zone was encountered during geological and scintillometer surveys. The best assay from pebble conglomerate was 0.015 percent U₃O₈.

References

Card and Palonen (1976, p.45)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2246, Birch Point Mines Limited, Great Yukon Mines Limited, and Sheeley Mining Corporation Limited, 1967.

Technical File No. 2.2293, Kerr Addison Mines Limited, 1976.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001100.

STETHAM TOWNSHIP

ASTRABRUN RADIOACTIVITY OCCURRENCE

Location

Latitude 47°48'29"N, Longitude 81°38'56"W.
Stetham Township.

Remarks

Early Precambrian granitic rocks are exposed along the southern shore of Kenetogami Lake. At one location along the shore, mafic rocks are intruded by radioactive pegmatite. Shearing and faulting occur at the contact. This general structure would project along strike to the occurrence.

References

Pyke et al. (1973)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario
Geological Survey, Toronto

Stetham Township, Drill Log Report
No. 10, Astrabrun Mines Limited

Technical File No. 63.3243, Astrabrun Mines Limited
Stetham Township, Drill Log Report
No. 13 and 14, Jonsmith Mines Limited
Technical File No. 2.2131, Beach Gold Mines Limited

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000428.

TURNER TOWNSHIP

DENISON (BULL LAKE) OCCURRENCE

Location

Latitude 47°03'50"N, Longitude 80°33'39"W.
Turner Township.

Remarks

In 1968, Denison Mines Limited drilled to a depth of 2159 feet. Minor radioactivity was detected in quartzite at 1794 feet. Poorly sorted, dark grey, pyritiferous quartzite grades into greywacke. Several pebble bands were located in the quartzite. Radioactivity was approximately three times the background.

References

Card et al. (1973, p.110-118)

Ontario Ministry of Natural Resources Files

Assessment Files Research Office, Ontario
Geological Survey, Toronto

Turner Township, Drill Log Report No. 16, Denison Mines Limited, 1968.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000406.

NORANDA (PILGRIM CREEK) OCCURRENCE

Location

Latitude 47°03'55"N, Longitude 80°37'17"W.
Turner Township.

Remarks

The Huronian metasediments consist of quartzite, argillite, greywacke, and quartz-pebble conglomerate. The quartzite is slightly radioactive.

References

Card et al. (1973, p.115)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto
 Technical File No. 63.2542, Canadian
 Johns-Manville Company Limited,
 1969.
 Turner Township, Drill Log Report No.
 11 and 14, Noranda Exploration
 Company Limited, 1967 and 1969.
 Source Mineral Deposits Record, Ontario
 Geological Survey, Toronto
 File No. 001099.

References

Robertson (1968a, p.51, 1976b, p.105)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto
 Technical File No. 63A.235, J.C.
 Humpage and G. Towle, 1955-1962.
 Source Mineral Deposits Record, Ontario
 Geological Survey, Toronto
 File No. 001111.
 Regional Geologist's Files, Sudbury

VICTORIA TOWNSHIP

SUGAR LAKE OCCURRENCE

Location

Latitude 46°14'11"N, Longitude 82°13'48"W.
 Victoria Township.

Remarks

The Espanola Formation here is locally radioactive. Assays indicated uranium, thorium, and niobium in sheared, impure quartzite.

During 1954 and 1956, an aeroradiometric survey, and geological and geophysical mapping were carried out by Blind River Uranium Mines Limited.

References

Robertson (1976b, p.50,76)
 Financial Post Survey of Mines (1957, p.203)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario
 Geological Survey, Toronto
 Technical File No. 63A.234, Blind
 River Uranium Mines Limited
 Technical File No. 2.2294
 Technical File No. 2.2452

TOWLE-HUMPAGE OCCURRENCE

Location

Latitude 46°13'00"N, Longitude 82°15'45"W.
 Victoria Township.

Remarks

Principal sedimentary formations outcropping in the area are quartzite and argillaceous metasediments. In pebbly quartzite, radioactivity is up to four times the background. Greater radioactivity is found in the shear zones in the argillaceous metasediments.

URANIUM AND THORIUM DEPOSITS OF THUNDER BAY DISTRICT

OCCURRENCES ON UNSURVEYED LAND

GREENWICH LAKE OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Uraninite.

Location
Latitude 48°46'57"N, Longitude 88°51'20"W.
District of Thunder Bay
Map Reference: ODM Map 2232.

Geology

The uranium mineralization is found where Early Precambrian (Kenoran) granite and pegmatite intrude granitic, biotite gneiss at a strike of N75E. The radioactivity occurs mainly in tension and shear fractures and joints dipping 80°E and striking N30W. To the west of the fault is a white-to-grey, medium-to coarse-grained biotite granite with scattered pegmatite segregations. Some of the fractures contain 1/4 inch stringers of pitchblende assaying up to 27 percent U₃O₈ (radiometric equivalent), while the intervening granite assayed 0.08 percent U₃O₈ (radiometric equivalent).

Economic Features

Two mineralized lenses outlined are 5.9 feet by 150 feet averaging 0.29 percent U₃O₈ (radiometric equivalent).

In 1975, Copper Lake Explorations Limited obtained assays of up to 1.6 pounds per ton U₃O₈. The average grade across 37 feet was 0.4 to 0.6 pounds U₃O₈ per ton.

In 1977, Rio Tinto Canadian Exploration Limited drilled three diamond drill holes. Assays ranged from 0.1 pound U₃O₈ per ton over 3 feet to 0.5 pounds per ton over two feet.

In 1978, Greenwich Lake Explorations Limited performed drilling, trenching, and sampling. The assays ranged from 0.001 to 0.073 percent U₃O₈.

History of Development

1949: Tom Christianson discovered radioactivity 500 feet west of the shore of Greenwich Lake.

Pre-1954: Trenching, line cutting, mapping, diamond drilling, and prospecting were completed by Great Lake Uranium Mines Limited.

1954: Sixteen diamond drill holes totalling 1319 feet were drilled by Pan Canadian Development Company Limited.

1955: Trenching and drilling were completed by The Associates.

1956: Trenching and drilling were carried out by The Associates and Climax Molybdenum Corporation of British Columbia Limited.

1958: A geological survey was carried out F.R. Joubin and Associates Limited.

1969: An airborne radiometric survey was completed by Univex Exploration and Development Corporation Limited.

1970: Fifty-one diamond drill holes totalling 6580 feet were drilled by Univex Exploration and Development Corporation.

1974: Consolidated Shunsby Mines Limited acquired these claims.

1975: Copper Lake Explorations Limited performed radiometric and geological surveys.

Six diamond drill holes were drilled by Consolidated Monarch Metal Mines Limited.

1976: Geological and geophysical surveys were completed by Rio Tinto Canadian Exploration Limited.

1977: Rio Tinto Canadian Exploration Limited carried out scintillometer, spectrometer, and geological surveys as well as diamond drilling and trenching.

Late in 1977 Consolidated Shunsby Mines Limited changed their name to Greenwich Lake Explorations Limited.

1978: Greenwich Lake Explorations Limited drilled 21 diamond drill holes for a total of 4024 feet.

1979: Turam and induced polarization surveys were completed by Greenwich Lake Explorations Limited.

References

- Canadian Mines Handbook (1975-1976, "MW Resources Limited")
Carter (1977)
Carter et al. (1973)
Robertson (1968a, p.88-89)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

- Greenwich Lake Area, Drill Log
Report No. 10, Pan Canadian
Development Company Limited, 1954.
Technical File No. 63A.358,
Greenwich Lake Area, 1958.
Technical File No. 63.2484, Univex
Exploration and Development
Company, 1954.

Greenwich Lake Area, Drill Log Report No. 11 and 13, Univex Exploration and Development Corporation Limited, 1970.
 Technical File No. 2.2072, Copper Lake Explorations Limited, 1975.
 Technical File No. 2.2295, Rio Tinto Canadian Exploration Limited, 1977.
 Technical File No. 2.3128, Greenwich Lake Explorations Limited, 1979.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 000277.

NEW INSCO PROSPECT (PRAIRIE LAKE COMPLEX)

Commodity
 Uranium, niobium

Radioactive Minerals
 Uraniferous pyrochlore, betafite, and apatite.

Location
 Latitude 49°01'48"N, Longitude 86°42'54"W.
 District of Thunder Bay
 Map Reference: ODM Map 2232.

Geology
 The Prairie Lake carbonatite-alkalic complex is a circular intrusion approximately 3.4 square miles in surface area. The entire complex is radioactive with the highest readings occurring at the contact between the outer carbonatite ring and the core of nepheline syenite. This contact can be traced for 2 miles between "Centre" and "Anomaly" Lakes. New InSCO Mines Limited have exposed the contact in three places (showings A, B, and C). In each location the contact is marked by betafite-bearing pyroxenite. This betafite is confined to the mixed rock - pyroxenite phase developed at the main carbonatite-syenite contact.

Economic Features
 A trench on showing A assayed 0.045 percent U_3O_8 over 10 feet. Four drill holes on the B showing (Jim's showing) over a strike length of 320 feet averaged 0.1 percent U_3O_8 and 0.27 percent Nb_2O_5 across an average of 23 feet. Four trenches on the C showing (North Highgrade showing) over a distance of 130 feet averaged 0.045 percent U_3O_8 across an average width of 14 feet.
 Work on the remainder of the complex has located significant concentrations of uraniumiferous pyrochlore and a fourth betafite occurrence at showing D (Discovery Zone) which averages 0.085 percent U_3O_8 .

Estimated mineralized reserves are 109,024 tons grading 0.12 percent U_3O_8 .

History of Development
 1968-1970: Ground radiometric, magnetic and geochemical surveys, diamond drilling totalling 1742 feet and trenching totalling 1375 feet were completed by Newmont Mining Corporation of Canada Limited.
 1976: A geochemical survey was completed by International Minerals and Chemical Corporation (Canada) Limited.
 Ground radiometric and geological surveys, pitting and trenching, and 150 overburden drill holes were all completed by New InSCO Mines Limited.
 1977: Fifteen diamond drill holes totalling 5053 feet were drilled by New InSCO Mines Limited.

References
 Carter et al. (1973)
 Robertson (1981)
 Sage (1975)
 The Northern Miner (1976, Nov. 11, p.17)

Ontario Ministry of Natural Resources Files
 Assessment Files Research Office, Ontario Geological Survey, Toronto
 Killala Lake Area, Drill Log Reports No. 10 and 11, 1969 and 1976.
 Technical Report No. 2.2099, International Minerals and Chemical Corporation, 1976.
 Technical Report No. 2.2372, New InSCO Mines Limited, 1976.
 Technical Report No. 2.2555, New InSCO Mines Limited, 1977.
 Source Mineral Deposits Record, Ontario Geological Survey, Toronto
 File No. 001127.

NEW SANTIAGO (STURGEON NARROWS) OCCURRENCE

Commodity
 Uranium, thorium, and niobium.

Radioactive Minerals
 Unknown.

Location
 Latitude 49°56'10"N, Longitude 90°51'00"W.
 District of Thunder Bay
 Map Reference: ODM Map 2169.

Geology
 This occurrence is situated in the Sturgeon Lake alkaline syenite complex

which cuts Early Precambrian volcanic rocks. Several radioactive zones up to 25 feet wide are associated with the contact and alteration zones and with the late pegmatites.

The Coveney Island contact zone has veinlets of red feldspar rock containing purple fluorite, sulphides, magnetite, and secondary carbonate.

Economic Features

Four radioactive dikes were located on Coveney Island. The best assays were 0.15 percent U_3O_8 and 0.48 percent ThO_2 (radiometric equivalent) and 0.015–0.020 percent Nb_2O_5 (radiometric equivalent).

The Anderson Island zone, located 1500 feet south of Anderson Lake, is geologically similar to the Coveney Island zone and assayed 0.006 percent U_3O_8 . A sample from the Sturgeon Narrows zone assayed 0.01 percent U_3O_8 (radiometric equivalent) and 0.01 percent Nb_2O_5 . A sample from the Seaton Island zone approximately 1.5 miles southwest of the above location assayed 0.04 percent U_3O_8 (radiometric equivalent).

History of Development

1955: An Ontario Department of Mines field party dug pits 6 feet deep and 4–5 feet wide, and also did geological mapping and spot radioactivity measurements.

1969: A geological survey was completed by W.G. Wahl Limited.

1970: An airborne electromagnetic and magnetic survey was completed by Consolidated Red Poplar Mines Limited and Green Point Mines Limited.

Electromagnetic and magnetometer surveys of the Six Mile Lake area were completed by Texmont Mines Limited.

1971: Magnetometer and electromagnetic surveys were completed by Green Point Mines Limited.

An airborne electromagnetic and magnetic survey was completed by Texmont Mines Limited and Sturdy Mines Limited.

References

Davies et al. (1970)
Robertson (1968a, p.89)
Rodgers (1964, p.42–44)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.2437, W.G.
Wahl Limited, 1969.
Technical File No. 63.2729,
Consolidated Red Poplar Mines
Limited and Green Point Mines
Limited, 1970.

Technical File No. 2.180, Texmont
Mines Limited, 1970.

Technical File No. 2.416, Green Point
Mines Limited, 1971.

Technical File No. 2.382, Texmont
Mines Limited, 1971.

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File 000314.

SANDY STONE (CHARRON LAKE) OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location

Latitude 49°38'46"N, Longitude 85°58'54"W.
District of Thunder Bay
Map Reference: ODM Map 2141.

Geology

The host rock consists of metasediments underlain by granitic rocks of Early Precambrian age. Pegmatites have intruded the mica schists.

Economic Features

The best assays averaged 0.02 percent U_3O_8 (radiometric equivalent).

History of Development

1955–1957: 1081 feet of diamond drilling was completed by Sandy Stone Exploration and Development Company.

References

Coates (1967a)
Hewitt (1967, p.55)
Lang et al. (1962)
Robertson (1968a, p.90)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000278.
Resident Geologist's Files, Kenora.

SANDY STONE (KASSAGIMINI LAKE) OCCURRENCE

Commodity
Uranium.

Radioactive Minerals

Uraninite.

Location

Latitude 49°43'20"N, Longitude 85°51'20"W.
District of Thunder Bay
Map Reference: ODM Map 2141.

Geology

Uranium mineralization is associated with pegmatite intrusions in mica schist.

Economic Features

The best assay obtained was 0.08 percent U_3O_8 (radiometric equivalent).

History of Development

1955-1957: Five diamond drill holes totalling 1084 feet were completed by the Sandy Stone Exploration and Development Company.

References

Coates (1967a)
Robertson (1968a, p.87)

Ontario Ministry of Natural Resources Files

Source Mineral Deposits Record, Ontario Geological Survey, Toronto
File No. 000279
Resident Geologist's Files, Kenora

DALEY TOWNSHIP**LONGLAC OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 49°45'00"N, Longitude 86°31'00"W.
Daley Township.
Map Reference: ODM Map 2101.

Geology

Radioactivity, associated with thorium, was encountered in a zone 2.0 feet wide and 40.0 feet long, in a quarry 1 mile east of Longlac (Lang 1952).

Economic Features

Core samples averaged 0.006 percent U_3O_8 (radiometric equivalent). The best core sample was 0.04 percent U_3O_8 (radiometric equivalent) over 1.75 feet and 0.04 percent U_3O_8 (chemical) (Assessment Files Research Office, Ontario Geological Survey, Toronto, Drill Log Reports No. 10 and 11, Floranda Mines Limited).

References

Lang (1952, p.119)
Pye et al. (1966)
Robertson (1968a, p.90)

DORION TOWNSHIP**INNES LAKE OCCURRENCE****Commodity**

Uranium.

Radioactive Minerals

Unknown.

Location

Latitude 48°50'50"N, Longitude 88°45'18"W.
Dorion Township.
Map Reference: ODM Map 2232.

Geology

This claim straddles the Good Morning Lake Fault. The fault strikes N30W and predates Late Precambrian fractures which strike N60E. Radioactive concentrations have been detected in hematite-quartz breccia. Assays of grab samples by the Geological Survey of Canada indicated 0.13 to 1.27 pounds per ton U_3O_8 (Franklin 1978).

References

Carter et al. (1973)
Franklin (1978)
Robertson (1981)

Ontario Ministry of Natural Resources Files

Resident Geologist' Files, Kenora.

MINOR URANIUM AND THORIUM OCCURRENCES OF THUNDER BAY DISTRICT

OCCURRENCES ON UNSURVEYED LAND

CELOTTI OCCURRENCE

Location

Latitude 48°47'50"N, Longitude 88°52'30"W.
District of Thunder Bay

Remarks

A zone of uranium mineralization extends for approximately 1300 feet across the southern portion of claim TB430716. The most radioactive section occurs near the east end of this zone and is 37 feet in width. Uranium mineralization occurs in two albitite layers and within para-gneissic metasediments adjacent to the albitite bands. The albitite consists of coarse-grained white plagioclase feldspar, quartz as individual grains and segregations, subordinate biotite, and minor muscovite.

Average radiometric assays across 37 feet were 0.4 to 0.6 pounds per ton U₃O₈.

References

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto
Technical File No. 2.2072, Greenwich
Lake Area, 1975.
Resident Geologist's Files, Kenora

PIC BAMOOS (PORT COLDWELL
COMPLEX) OCCURRENCE**Location**

Latitude 48°42'31"N, Longitude 86°20'24"W.
District of Thunder Bay

Remarks

The area is dominated by the Port Coldwell Precambrian alkalic complex. Radioactive mineralization occurs in deposits of fractured and sheared pegmatites. Samples submitted in 1949 were from a zone of pegmatite 40 feet thick, and assayed 0.01 percent and 0.45 percent U₃O₈ (radiometric equivalent).

References

Milne et al. (1972)
Puskas (1967, p.86-87)
Robertson (1968a, p.87)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000434.

SQUAW LAKE OCCURRENCE

Location

Latitude 50°03'45"N, Longitude 90°43'00"W.
District of Thunder Bay

Remarks

Trowell (1976) divided these rocks into two broad groupings. These are (1) coarse-grained to pegmatitic alkalic phases of syenite and monzonite in the north half of the Squaw Lake complex, and (2) medium-grained, equigranular and quartz-bearing syenite, syenomonzonite and monzonite in the south half.

References

Davies et al. (1970)
Sage (1976)

UNIVEX OCCURRENCE

Location

Latitude 48°50'30"N, Longitude 88°53'00"W.
District of Thunder Bay

Remarks

Airborne radiometric surveys were performed over migmatite areas. Diamond drilling that followed detected low amounts of uranium and thorium.

References

Carter (1977)

YZERDRAAT OCCURRENCE

Location

Latitude 50°09'41"N, Longitude 87°38'40"W.
District of Thunder Bay

Remarks

A large northwest-trending metavolcanic-metasedimentary belt is bordered by granite and other felsic intrusives to the west and the south. These rocks are intruded by diabase and porphyritic diabase.

A radiometric survey gave spot highs of 30 counts per second over a background of 3

to 5 counts per second. No radioactive mineral source was identified however.

References

Amukun (1977, p.80)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 2.2114, Walter
Yzerdraat, 1967.

Technical File No. 2.2283, Walter
Yzerdraat, 1967.

Technical File No. 2.2572,
Radiometric Survey of the Metcalfe
Lake Area, W. Yzerdraat, 1977.

Technical File No. 2.2874, Geological
Survey of the Metcalfe Lake Area, W.
Yzerdraat, 1979.

Technical File No. 2.3043,
Radiometric Survey of the Metcalfe
Lake Area, 1979

Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 001140.

BOMBY TOWNSHIP

HEMLO OCCURRENCE

Location

Latitude 48°41'50"N, Longitude 85°55'00"W.
Bomby Township.

Remarks

Radioactivity was discovered at a gold prospect owned by Lake Superior Mining Corporation Limited near Hemlo. Five parallel radioactive zones have been found along a contact between biotite granite and metavolcanics. Anomalies up to ten times the background have been obtained. Two samples sent to the Geological Survey of Canada showed 0.06 and 0.09 percent U_3O_8 (radiometric equivalent) (Lang 1952).

References

Lang (1952, p.118)

Robertson (1968a, p.90)

CONMEE TOWNSHIP

KAKABEKA FALLS OCCURRENCE

Location

Latitude 48°24'00"N, Longitude 89°35'15"W.
Conmee Township.

Remarks

Two grab samples of mud-ball tuff of the upper tuffaceous shale facies of the Gunflint Formation, at the Ontario Hydro diversion plant at Kakabeka Falls, each assayed 0.004 percent U_3O_8 .

References

Fenwick and Scott (1977)

Pye and Fenwick (1963)

Robertson (1981)

DORION TOWNSHIP

DORION AMETHYST MINE

Location

Latitude 48°49'00"N, Longitude 88°35'40"W.
Dorion Township.

Remarks

The Dorion vein has dolomite of the Rossport Formation on its southeast wall and Early Precambrian monzonite on its northwest wall. The vein extends along strike for approximately 5000 feet and consists of five separate veins. The veins contain 20 percent and locally up to 80 percent breccia fragments. They include unaltered quartz monzonite and highly silicified dolomite. Radioactivity and sulphide mineralization are associated in the matrix of the breccia.

References

Franklin and Mitchell (1977)

McIlwaine and Tihor (1975)

Robertson (1981)

GOLDIE TOWNSHIP

NELSON OCCURRENCE

Location

Latitude 48°39'48"N, Longitude 89°53'25"W.
Goldie Township.

Remarks

Uraninite occurs in a feldspar-quartz-biotite veinlet, 1 inch wide. Radioactivity is two times the background.

References

Pye and Fenwick (1963)

Robertson (1968a, p.87)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000432.

HELE TOWNSHIP

TESSIER-WILLIAMSON OCCURRENCE

Location

Latitude 49°00'10"N, Longitude 88°29'00"W.
Hele Township.

Remarks

Uranium mineralization occurs in a pegmatite dike 2 to 8 feet wide. This anomalous zone corresponds with a narrow steep-walled valley in biotite-quartz-feldspar gneiss that was intruded lit-par-lit by pegmatite.

References

Coates (1972, p.33)
Robertson (1968a, p.86)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000043.

LABERGE TOWNSHIP

MOLSON LAKE OCCURRENCE

Location

Latitude 48°41'40"N, Longitude 85°54'10"W.
Laberge Township.

Remarks

Radioactivity was found in five zones along the contact between granite and metavolcanics.

References

Milne et al. (1972)
Prasad (1981, p.31)

PUNKARI OCCURRENCE

Location

Latitude 48°41'00"N, Longitude 85°37'20"W.
Laberge Township.

Remarks

This occurrence is in an area underlain by felsic igneous rocks of Early Precambrian age, located in the Superior Structural Province.

No other data is available.

References

Milne et al. (1972)
Robertson (1968a, p.90)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000262.

MCCOY TOWNSHIP

PORT MONROE (POTVINE) OCCURRENCE

Location

Latitude 48°46'42"N, Longitude 86°26'27"W.
McCoy Township.

Remarks

Radioactive fractures occur in red syenite dikes cutting amygdaloidal lavas. The fracture system averages approximately 3 inches wide and 200 hundred feet long. Samples assayed up to 0.19 percent U_3O_8 (chemical) and 0.07 percent U_3O_8 (radiometric equivalent).

References

Milne et al. (1972)
Robertson (1968a, p.90)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000266.

MCINTYRE TOWNSHIP

PORT ARTHUR OCCURRENCE

Location

Latitude 48°24'53"N, Longitude 89°15'48"W.
McIntyre Township.

Remarks

Uranium mineralization occurs in narrow veins, seldom more than 6 inches wide, of anthraxolite which fills fractures in Animikie sedimentary rocks and diabase. Also present in minor amounts are silver, argentite, galena, chalcopryrite, sphalerite, pyrite, quartz, calcite, fluorite, and barite. A grab sample assayed 0.0034 percent U_3O_8 (chemical).

References

Pye and Fenwick (1963)

Robertson (1968a, p.87)
Tanton (1931, p.201)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000274

MCTAVISH TOWNSHIP

ENTERPRISE MINE

Location

Latitude 48°30'46"N, Longitude 88°37'12"W.
McTavish Township.

Remarks

The Enterprise mine's main amethyst vein is the only uranium-bearing vein of a group of lead-zinc-barite veins found throughout the southern part of the Sibley Basin. These veins occur in Rosspoint dolomite immediately above the zone where the Pass Lake sandstone thins against the Early Precambrian paleo-positive area that is composed of quartz monzonite (Franklin 1978).

According to Ruzicka (1977), uranium mineralization in the Sibley Formation, approximately 10 m above the pre-Sibley unconformity was identified by H.R. Steacy and A.G. Plant as coffinite. The coffinite rims and replaces pyrite grains and also occurs as irregular masses.

References

Franklin (1978)
Robertson (1981)
Ruzicka (1977)

PEARL LAKE OCCURRENCE

Location

Latitude 48°39'40"N, Longitude 88°39'35"W.
McTavish Township.

Remarks

The regional geology here includes the sedimentary rocks of the Sibley Group, and the contact between Early Precambrian biotite quartz monzonite to granodiorite and sedimentary rocks of the Sibley Group.

References

McIlwaine (1971)
Robertson (1971)

O'NEILL TOWNSHIP

CRAMETTE OCCURRENCE

Location

Latitude 48°46'00"N, Longitude 86°25'00"W.
O'Neill Township.

Remarks

Two samples from a showing discovered by C. Cramette assayed 0.06 and 0.07 percent U₃O₈ (radiometric equivalent).

References

Lang et al. (1962, p.257)

PIC TOWNSHIP

SMITH OCCURRENCE

Location

Latitude 48°35'28"N, Longitude 86°09'20"W.
Pic Township.

Remarks

Mineralization is associated with a shear zone located in syenite-pegmatite areas up to 3 feet by 3 feet. Radiometric readings are up to 11 times the background. Tests indicated radioactivity is due entirely to thorium.

References

Milne et al. (1972)
Robertson (1968a, p.90)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto
File No. 000267.

WALSH TOWNSHIP

DEADHORSE CREEK (DIATREME) OCCURRENCE

Location

Latitude 48°53'00"N, Longitude 86°39'20"W.
Walsh Township.

Remarks

This occurrence is associated with a diatreme located in the Early Precambrian metavolcanic-metasedimentary belt west of the Port Coldwell alkaline complex. The diatreme crosscuts the metavolcanics and pyroclastics.

The matrix is green and consists of carbonate and a greenish amphibole.

Embedded in this matrix are clasts of varying size and angularity.

A scintillometer survey traced the diatreme for 500 m in a north-south direction, 275 m west of, and 400 m northeast of Deadhorse Creek. Assays of grab samples showed values of 0.003 and 0.004 percent U_3O_8 .

References

Fenwick and Scott (1978)
Sage (1978)
Walker (1967)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File 2.2844, Gulf Minerals
Canada Limited, 1978
Walsh Township, Drill Log Reports No.
20 and 21, Gulf Minerals Canada
Limited, 1978.

MCKELLAR CREEK (DIATREME) OCCURRENCE

Location

Latitude 48°51'40"N, Longitude 86°41'00"W.
Walsh Township.

Remarks

This complex consists of two outcrops and numerous patches of radioactive rubble. The diatreme has a north-trending axis, and is approximately 240 m long by 60 m wide. Radioactive breccia occurs within outcrops of Early Precambrian fine-grained schistose, argillite and siltstone.

References

Robertson (1981)
Sage (1978)
Walker (1967)

URANIUM AND THORIUM DEPOSITS OF TIMISKAMING DISTRICT

LEBEL TOWNSHIP

KAPLAN OCCURRENCE

Commodity
Uranium.

Radioactive Minerals
Unknown.

Location
Latitude 48°09'48"N, Longitude 79°53'10"W.
Lebel Township.
Map Reference: ODM Map 2205.

Geology
This occurrence is associated with greenish-black, tuffaceous greywacke with areas of quartzite. To the west, the radioactive beds branch out for a total length of 1600 feet with widths up to 8 feet. This unit has an average dip of 80° to the north.

Economic Features

In 1957, a radioactive boulder assayed 2.16 pounds U₃O₈ per ton. A 5-foot chip sample yielded the highest assay at 2.0 pounds U₃O₈ per ton. Diamond drilling revealed a section 3.5 feet wide, containing 1.5 pounds U₃O₈ per ton.

History of Development

1928: Initially explored for gold as part of the Crystal Kirkland property.
1957: Rio Algom Mines Limited dug trenches.
1966: Rio Algom Mines Limited drilled seven diamond drill holes.

References

Pyke et al. (1973)
Robertson (1968a, p.92; 1981)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kirkland Lake.

MINOR URANIUM AND THORIUM OCCURRENCES OF TIMISKAMING DISTRICT

BOSTON TOWNSHIP

EMPIRE OCCURRENCE

Location

Latitude 48°03'58"N, Longitude 79°59'09"W.
Boston Township.

Remarks

The Empire occurrence is located within a siliceous band of very "lean" iron formation that is cut by stringers of fine-grained, mafic intrusive rocks. Radioactivity is confined to the face of one of the several joints that strike S70E and dip steeply to the north.

References

Pyke et al. (1973)
Robertson (1968a, p.93)

LUNGE OCCURRENCE

Location

Latitude 48°04'51"N, Longitude 79°59'10"W.
Boston Township.

Remarks

Radioactivity is found within irregular dikes of pink syenite cutting hornblendite. The dikes are situated close to the southwest contact of the Lebel Syenite Stock and the metavolcanic-metasedimentary belt to the south. In 1953, the occurrence was examined with a geiger counter by W.S. Savage, Resident Geologist, Ontario Department of Mines.

In 1972, Marshall Boston Iron Mines Limited conducted magnetometer, electromagnetic, geological and geochemical surveys.

References

Robertson (1968a, p.93)

Ontario Ministry of Natural Resources Files
Assessment Files Research Office, Ontario
Geological Survey, Toronto

Technical File No. 63.3094, Marshall
Boston Iron Mines Limited, 1972.
Technical File No. 2.836, Raymond.

YOST OCCURRENCE

Location

Latitude 48°02'10"N, Longitude 79°59'12"W.
Boston Township.

Remarks

The occurrence is located within a small body of Early Precambrian (Algoman) syenite that intrudes metavolcanics. Geiger counter readings were up to three times the background.

References

Pyke et al. (1973)

Ontario Ministry of Natural Resources Files
Source Mineral Deposits Record, Ontario
Geological Survey, Toronto

File No. 000324
Resident Geologist's Files, Kirkland Lake

BUCKE TOWNSHIP

HARRISON-HIBBERT OCCURRENCE

Location

Latitude 47°24'39"N, Longitude 79°37'20"W.
Bucke Township.

Remarks

Uranium (up to 4 pounds U₃O₈ per ton) occurs in a fault breccia in association with carbonate veins.

References

Card and Lumbers (1977)
Sergiades (1968)

CANE TOWNSHIP

CANE TOWNSHIP OCCURRENCE

Location

Latitude 47°36'05"N, Longitude 80°01'55"W.
Cane Township.

Remarks

Radioactivity was detected intermittently over a distance of 1.5 km, between diabase and quartzite of the Lorrain Formation. The uranium-bearing mineral was identified as pitchblende by R. Thomson, Ontario Department of Mines, (unpublished report in Resident Geologist's Files, Ontario Ministry of Natural Resources, Kirkland Lake). It is associated with carbonate veins and with granophyre differentiates (aplite and micropegmatite) and transitional rock (diabase of varied texture) near the upper contact of a Nipissing diabase basin 8 to 16 km in diameter (Lovell and Ploeger 1977).

References

Lovell and Ploeger (1977)
 Pyke et al. (1973)
 Robertson (1968a, p.90)

COLEMAN TOWNSHIP**KERR LAKE MINE****Location**

Latitude 47°22'20"N, Longitude 79°39'13"W.
 Coleman Township.

Remarks

Radioactivity is associated with veins in diabase that intrude metavolcanics and metasediments. A sample of cobalt-silver ore on display at the Cobalt Mining Museum was found to be radioactive.

References

Card and Lumbers (1977)
 Prasad (1981, p.65)
 Ruzicka (1979)

GILLIES LIMIT TOWNSHIP**SILVERMAQUE MINE****Location**

Latitude 47°20'37"N, Longitude 79°38'33"W.
 Gillies Limit Township

Remarks

Radioactive mineralization was found in argillitic rocks on a dump of the Silvermaque Mine. The radioactive minerals were pitchblende and allanite.

References

Card and Lumbers (1977)
 Ruzicka (1979)

HUDSON TOWNSHIP**SPENCER OCCURRENCE****Location**

Latitude 47°34'00"N, Longitude 79°52'00"W.
 Hudson Township.

Remarks

Uranium mineralization is in the Firstbrook argillite. Radioactive minerals are in bands of breccia consisting of argillite reddened and hardened in a dark green chloritic matrix. Seventeen samples sent to the Ontario Department of Mines showed

radioactivity ranging from nil to 0.18 percent U_3O_8 (radiometric equivalent).

References

Lovell and Fily (1970)
 Lovell and Ploeger (1977)
 Thomson (1966)

LEBEL TOWNSHIP**BIDGOOD KIRKLAND MINE****Location**

Latitude 48°10'05"W Longitude 79°54'45"W.
 Lebel Township.

Remarks

At the former gold-silver mine, radioactivity was detected by D. Lowe in 1966 across 90 m and along 360 m of intermittent outcrop. Radioactivity is more intense in interbedded conglomerate and greywacke and is most intense in rocks containing quartz veins. Assays were 0.2 pounds U_3O_8 per ton (radiometric equivalent).

References

Lovell and Ploeger (1977)
 Pyke et al. (1973)
 Robertson (1981)

HURD OCCURRENCE**Location**

Latitude 48°08'46"N, Longitude 79°58'04"W.
 Lebel Township.

Remarks

Widespread radioactivity is caused mainly by thorium, and little uranium is present.

References

Pyke et al. (1973)

Ontario Ministry of Natural Resources Files
 Resident Geologist's Files, Kirkland Lake

LEBEL OCCURRENCE**Location**

Latitude 48°09'50"N, Longitude 79°55'00"W.
 Lebel Township.

Remarks

The Lebel occurrence is associated with Timiskaming amygdaloidal trachyte cut by

Early Precambrian (Algoman) syenite.
Radioactivity was three to four times the
background over a 70-foot outcrop.
Representative samples assayed 0.032-0.05
percent U_3O_8 (Robertson 1968a, p.92).

References

Pyke et al. (1973)
Robertson (1968a, p.92)

Ontario Ministry of Natural Resources Files
Resident Geologist's Files, Kirkland Lake.

TUDHOPE TOWNSHIP**SAUVE-INCH OCCURRENCE****Location**

Latitude 47°45'17"N, Longitude 80°19'32"W.
Tudhope Township.

Remarks

This occurrence contains small amounts
of pitchblende in a carbonate vein along the
contact of a Nipissing Diabase dike and
Gowganda Formation conglomerate. A
sample assayed 1.56 percent U_3O_8 .

References

Johns (1980)
Pyke et al. (1973)

Ontario Ministry of Natural Resources Files
Regional Geologists File's, Kirkland Lake

REFERENCES

- Adams, S.S. and Button, A.
1981: Geology and Recognition Criteria for Uranium Deposits of the Quartz-Pebble Conglomerate Type, Final Report; National Uranium Resource Evaluation; Prepared for the United States Department of Energy, Grand Junction Office, Colorado. GTX-3(81).
- Airth, M.W. and Olson, E.R.
1958: Algom-Nordic Development to Production; The Canadian Mining and Metallurgical Bulletin, Canadian Institute of Mining and Metallurgy, Vol. 51, No. 559, p. 666-676.
- Amukun, S.E.
1977: Geology of the Tashota Area; Ontario Geological Survey, Report 167, 40p.
- Ayres, L.D., Randsepp, M., Averill, S.A., and Edwards, G.R.
1972: Favourable Lake - Poplar Hill Area; Indian Village - Moose Mountain Sheet, District of Kenora (Patricia Portion); Ontario Division of Mines, Preliminary Map P.769, Geological Compilation Series, scale 1 inch to 2 miles. Geological Compilation 1968-1972.
- 1973: Favourable Lake - Berens Lake, Kenora District; Ontario Division of Mines, Geological Compilation Map 2262, scale 1 inch to 4 miles.
- Beard, R.C.
1977: Kenora Uranium Deposits; Proceedings of the 33rd Annual Meeting of Lake Superior Geology, 1977.
- Beard, R.C., and Rivett, S.
1977: 1976 Report of the Northwestern Regional Geologist and Kenora Resident Geologist; p. 3-15, in Annual Report of the Regional and Resident Geologists 1976, edited by C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 71, 142p.
- 1978: 1977 Report of the Northwestern Regional Geologist and Kenora Resident Geologist; p. 1-13, in Annual Report of the Regional and Resident Geologists 1977, edited by C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 78, 121p.
- Beard, R.C., and Scott, W.R.
1976: 1975 Report of the Kenora Resident Geologist; p. 1-17, in Annual Report of the Regional and Resident Geologists, 1975, edited by C.R. Kustra, Ontario Division of Mines, Miscellaneous Paper 64, 146 p.
- Bennett, G.
1976: Huronian Volcanism in the Thessalon Area, District of Algoma; p. 111-113, in Summary of Fieldwork, 1976, by the Geological Branch, edited by V. G. Milne, W.R. Cowan, K.D. Card, and J.A. Robertson, Ontario Division of Mines, Miscellaneous Paper 67, 183p.
- Bennett, G., Brown, D.D., and George, P.T.
1968: Coral Rapids-Cochrane Sheet, Cochrane District; Ontario Department of Mines, Geological Compilation Series, Map 2161. Scale 1:253 440.
- Blackburn, C.E.
1973: Geology of the Otukamamoan Lake Area, District of Kenora; Ontario Division of Mines, Geological Report 109, 42p. Accompanied by Map 2266, scale 1 inch to 1 mile.
- 1981: Kenora-Fort Frances, Kenora and Rainy River Districts; Ontario Geological Survey, Map 2443, Geological Compilation Series, scale 1:253 440.
- Bond, W.D., and Breaks, F.W.
1978: Ground Evaluation of Airborne Radiometric Anomalies in Northwestern Ontario; p. 22-27 in Summary of Fieldwork, 1978, by the Ontario Geological Survey, edited by V.G. Milne, O.L. White, R.B. Barlow, and J.A. Robertson, Ontario Geological Survey, Miscellaneous Paper 82, 235p.
- Breaks, F.W., Bond, W.D., MacWilliams, G.H., Gower, C.F., and Stone, D.
1975: Operation Kenora-Sydney Lake, Eagle-Sydney Lakes Sheet, District of Kenora; Ontario Division of Mines, Preliminary Map P. 1026, Geological Series, scale 1 inch to 1 mile or 1:63 360. Geology 1974
- Card, K.D.
1975: Mongowin and Curtin Townships, District of Sudbury; Ontario Division of Mines, Geological Map 2312, scale 1 inch to 1/2 mile.

- Card, K.D., Innes, D.G., and Debicki, R.L.
1974: Stratigraphy, Sedimentology and Petrology of the Huronian Supergroup of the Sudbury-Espanola Area, Ontario; Ontario Division of Mines, Open File Report 5107.
- Card, K.D., and Lumbers, S.B.
1977: Sudbury-Cobalt, Districts of Cochrane, Sudbury, and Timiskaming; Ontario Division of Mines, Map 2361. Geological Compilation Series, Scale 1:253 440 or 1 inch to 4 miles.
- Card, K.D., McIlwaine, W.H., and Meyn, H.D.
1973: Geology of the Maple Mountain Area, District of Timiskaming, Nipissing, and Sudbury; Ontario Division of Mines, Geological Report 106, 133p. Accompanied by Map 2256, 2257, 2258, and 2259, scale 1 inch to 1 mile. Map 2260, scale 1 inch to 1/2 mile.
- Card, K.D. and Palonen, P.
1976: Geology of the Dunlop-Shakespeare Area, District of Sudbury, Ontario Division of Mines, Geoscience Report 139, 52p. Accompanied by Map 2313 scale 1:31 680 or 1 inch to 1/2 mile.
- Carter, M.W.
1977: Greenwich Lake Area, District of Thunder Bay; p. 64-68, in Summary of Fieldwork 1977, by the Geological Branch, edited by V.G. Milne, O.L. White, R.B. Barlow, and J.A. Robertson, Ontario Geological Survey Miscellaneous Paper 75, 208 p.
- Carter, M.W., McIlwaine, W.H., and Wisbey, P.A.
1973: Nipigon-Schreiber, Thunder Bay District; Ontario Division of Mines, Geological Map 2232, scale 1:253 440 or 1 inch to 4 miles.
- Chandler, F.W.
1973: Geology of McMahan and Morin Townships, District of Algoma; Ontario Division of Mines, Geological Report 112, 77p. Accompanied by Map 2272, scale 1 inch to 1/2 mile.
- Chisholm, E.O.
1951: Preliminary Report on Radioactive Occurrences in the Kenora Area; Ontario Department of Mines, Preliminary Report 1950-1.
- Coates, M.E.
1967a: Stevens Sheet, Thunder Bay District; Ontario Department of Mines, Geological Map 2141, scale 1:63 360 or 1 inch to 1 mile.
1967b: Shillabear Creek Sheet, Thunder Bay District; Ontario Department of Mines, Map 2236, scale 1:63 360 or 1 inch to 1 mile.
1972: Geology of Black Sturgeon Lake Area, District of Thunder Bay; Ontario Department of Mines and Northern Affairs, Geological Report 98, 41p. Accompanied by Map 2233, 2234, 2235 and 2236, scale 1 inch to 1 mile.
- Davies, J.C., Pryslak, A.P., and Pye, E.G.
1970: Sioux Lookout-Armstrong Sheet, Kenora and Thunder Bay Districts; Ontario Division of Mines, Geological Compilation Series, Map 2169, scale 1:253 440 or 1 inch to 4 miles.
- Dressler, B.O.
1979: Geology of McNish and Janes Townships, District of Sudbury; Ontario Geological Survey, Report 191, 91p. Accompanied by Map 2425, scale 1:31 680 or 1 inch to 1/2 mile.
- Fenwick, K.G., and Scott, J.F.
1977: 1976 Report of the North-Central Regional Geologist; p. 42-56 in Annual Report of the Regional and Resident Geologists, 1976, edited by C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 71, 142p.
1978: Report of the North Central Regional Geologist; p. 46, in Annual Report of the Regional and Resident Geologists, 1977, edited by C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 78, 121p.
- Ferguson, S.A., Brown, D.D., Davies, J.C., and Pryslak, A.P.
1967: Red Lake-Birch Lake Sheet, Kenora District; Ontario Department of Mines, Geological Compilation Sheet, Map 2175, scale 1:253 440.
- Financial Post
various: Financial Post Survey of Mines; published annually by MacLean Hunter.

- Franklin, J.M.
1978: Uranium Mineralization in the Nipigon Area, Thunder Bay District of Ontario; Current Research, Part A, Geological Survey Canada, Paper 78-1A, p. 275-282.
- Franklin, J.M., and Mitchell, R.H.
1977: Lead-Zinc-Barite Veins of the Dorion Area, Thunder Bay District, Ontario; Canadian Journal of Earth Sciences, Vol. 14, No. 9, p. 1963-1979.
- Frarey, M.J.
1962: Map 32-1962; Geological Survey of Canada.
1977: Geology of the Huronian Belt Between Sault Ste. Marie and Blind River; Geological Survey of Canada, Memoir 383.
- Giblin, P.E., and Leahy, E.J.
1979: 1978 Report of the Sault Ste. Marie Resident Geologist; p.89-98, in Annual Report of the Regional and Resident Geologists, 1978, edited by C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 84, 135p.
- Giblin, P.E., Leahy, E.J., and Robertson, J.A.
1979: Sault Ste. Marie - Elliot Lake, Algoma, Manitoulin, and Sudbury Districts; Ontario Geological Survey, Map 2419, Geological Compilation Series, scale 1:253 440.
- Gordon, J.B., Robertson, J.A., and Rybak, U.C.
1981: Radioactive Deposits of Southern Ontario; Ontario Geological Survey, Open File Report 5311.
- Griffith, J.W.
1967: The Uranium Industry - Its History, Technology and Prospects; Mineral Resources Division, Energy, Mines and Resources, Canada, Mineral Report No. 12, 335p.
- Harding, W.D.
1950: Geology along the Mississagi Road, Algoma and Sudbury Districts; Ontario Department of Mines, Preliminary Report 1950-6.
- Hewitt, D.F.
1967: Pegmatite Mineral Resources of Ontario, Ontario Department of Mines, Industrial Minerals Report 12.
- Hogg, N.
1948: Report on Mosher Discovery of Radioactive Minerals, Pitt Township, District of Cochrane; Ontario Department of Mines, Preliminary Report 1948-7, 10p.
- Houston, R.S. and Karlstrom, K.E.
1981: Uranium-Bearing Quartz-Pebble Conglomerate Type, Final Report; National Uranium Resource Evaluation; Prepared for the United States Department of Energy, Grand Junction Office, Colorado, GJBX-1, p. 1-509.
- Hurst, M.E.
1930: Geology of the Area between Favourable Lake and Sandy Lake, Kenora District; Ontario Department of Mines, Annual Report for 1929, Vol. 38, part 2, p.49-84. Accompanied by Map 38a, scale 1 inch to 2 miles.
- Johns, G.W.
1980: Hill Lake Area, District of Timiskaming, p. 89-91 in Summary of Field Work, 1980, by the Ontario Geological Survey, edited by V.G. Milne, O.L. White, R.B. Barlow, J.A. Robertson, and A.C. Colvine, Ontario Geological Survey, Miscellaneous Paper 96, 201p.
- Kelly, T.J.
1960: Statistical Review of the Mineral Industry for 1958; Ontario Department of Mines, Annual Report for 1959, Vol. 68, part 1, p.1-27.
1961: Statistical Review of the Mineral Industry for 1959; Ontario Department of Mines, Annual Report for 1960, Vol. 69, part 1, p.1-40.
- Kelly, T.J. and Riddell, G.S.
1962: Statistical Review of the Mineral Industry for 1960 combined with Mining Operations; Ontario Department of Mines, Annual Report, Vol.70.
1963: Statistical Review of the Mineral Industry and Mining Operations for 1961; Ontario Department of Mines, Annual Report for 1961, Vol.71.
1964: Statistical Review of the Mineral Industry and Mining Operations for 1962; Ontario Department of Mines, Annual Report for 1962, Vol.72.

- Lang, A.H.
1952: Canadian Deposits of Uranium and Thorium (Interim Account); Geological Survey of Canada, Economic Geology Series No. 16, 173p.
- Lang, A.H., Griffith, J.W., and Steacy, H.R.
1962: Canadian Deposits of Uranium and Thorium; Geological Survey of Canada, Economic Geology Series No. 16 (Second Edition), 324p.
- Leahy, E.J.
1973: Diamond Drilling in the Huronian Supergroup, Sault Ste. Marie - Elliot Lake Area, Volume 1: Summary of Diamond Drilling with Abstracts of Selected Drill Holes (accompanied by Vol. 2, Atlas of Maps showing locations of drilling); Ontario Division of Mines, Open File Report 5093.
- Little, H.W., Smith, E.E.N., Barnes, F.Q., et al.
1972: Uranium Deposits of Canada; International Geological Guidebook, No.24, Trip 67, 64p.
- Lovell, H.L., and Fily, E.D.
1970: Kerns and Hudson Townships, District of Timiskaming, Ontario Department of Mines, Map 2300. Geological Compilation Series, Scale 1:31 680 or 1 inch to 1/2 mile.
- Lovell, H.L. and Ploeger, F.R.
1977: 1976 Report of the Kirkland Lake Resident Geologist; p. 77-94 in Annual Report of the Regional and Resident Geologists, 1976; edited by C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 71, 142p.
- Lumbers, S.B.
1975: Geology of the Burwash Area, Districts of Sudbury, Nipissing, and Parry Sound; Ontario Division of Mines, Geological Report 116, 158p. Accompanied by Map 2271, scale 1 inch to 2 miles.
- McIlwaine, W.H.
1971: McTavish Township (West Part of North Half), District of Thunder Bay; Ontario Department of Mines and Northern Affairs, Preliminary Map P.720, scale 1 inch to 1/4 mile. Geology 1971.
- McIlwaine, W.H., and Tihor, L.A.
1975: Dorion-Wolf Lake Area (Eastern Part), District of Thunder Bay; Ontario Division of Mines, Preliminary Map P.995, Geological Series, scale 1 inch to 1/4 mile or 1:15 840. Geology 1972.
- McMillan, R.H.
1977: Metallogenesis of Canadian Uranium Deposits: A Review; in Geology, Mining and Extractive Processing of Uranium, edited by M.J. Jones, Proceedings of a Symposium sponsored by the Institute of Mining and Metallurgy, London, Jan. 17-19, 1977.
- Meyn, H.D.
1970: Geology of Hutton and Parkin Townships; Ontario Department of Mines, Geological Report 80, 78p.
- 1971: Geology of Roberts, Creelman, and Fraleck Townships, District of Sudbury; Ontario Department of Mines, Geology Report 91, 48p. Accompanied by Map 2212, scale 1 inch to 1/4 mile.
- 1972: Geology of Grigg and Stobie Townships, District of Sudbury; Ontario Department of Mines and Northern Affairs, Geological Report 100, 38p. Accompanied by Map 2238, Scale 1 inch to 1/2 mile.
- 1979: Uranium Deposits of the Cobalt Embayment; p.218-221, in Summary of Fieldwork, 1979, by the Ontario Geological Survey, edited by V.G. Milne, O.L. White, R.B. Barlow, and C.R. Kustra, Ontario Geological Survey, Miscellaneous Paper 90, 240p.
- Meyn, H.D., and Mathews, M.K.
1980: Uranium Deposits of the Cobalt Embayment; p.195-199, in Summary of Fieldwork, 1980, by the Ontario Geological Survey, edited by V.G. Milne, O.L. White, R.B. Barlow, J.A. Robertson, and A.C. Colvine, Ontario Geological Survey, Miscellaneous Paper 96, 201p.
- Milne, V.G., Giblin, P.E., Bennett, G., Thurston, P., Wolfe, W.J., Giguere, J.F., Leahy, E.J., and Rupert, R.J.
1972: Manitouwadge-Wawa Sheet, Districts of Algoma, Cochrane, Sudbury, and Thunder Bay; Ontario Division of Mines, Map 2220, Geological Compilation Series, scale 1:253 440.

Northern Miner Press

various: Canadian Mines Handbook;
published annually, Northern Miner Press.

various: The Northern Miner; published
weekly, Northern Miner Press.

Nuffield, E.W.

1956: Geology of the Montreal River Area;
Ontario Department of Mines, Annual
Report for 1955, Vol. 64, Part 3, 32p.
Accompanied by Map 1955-1, scale 1 inch
to 1/2 mile.

OECD

1979: Uranium: Resources, Production and
Demand; A Joint Report by the OECD
Nuclear Energy Agency and the
International Atomic Energy Agency,
December 1979; Organization for
Economic Cooperation and Development,
135p. Paris, France.

Parsons, G.E.

1961: Niobium-Bearing Complexes East of
Lake Superior; Ontario Department of
Mines, Geological Report 3, 73p.
Accompanied by Map 2006, scale 1 inch to
1/4 mile.

Pirie, J.

1978: Crooked Pine Lake, Rainy River
District; Ontario Geological Survey, Map
2405, scale 1:31 680 or 1 inch to 1/2 mile.

Prasad, N.

1981: Distribution of Uranium and Thorium
Occurrences in Canada; Geological Survey
of Canada, Open File Report 551, 124p.

Pryslak, A.P.

1974a: MacNicol and Tustin Townships,
Kenora District; Ontario Division of
Mines, Geological Map 2302, scale 1:31
680 or 1 inch to 1/2 mile.

1974b: MacNicol and Tustin Townships,
Kenora District; Ontario Division of
Mines, Geological Map 2303, scale 1:31
680 or 1 inch to 1/2 mile.

1976: Geology of the Bruin Lake - Edison
Lake Area, District of Kenora; Ontario
Division of Mines, Geoscience Report 130,
61p. Accompanied by Maps 2302 and
2303, scale 1:31 680.

Puskas, F.P.

1967: Geology of the Port Coldwell Area,
Ontario Department of Mines, Open File
Report 5014, 92p.

Pye, E.G., and Fenwick, K.G.

1963: Atikokan-Lakehead Sheet, Kenora,
Rainy River and Thunder Bay Districts;
Ontario Department of Mines, Map 2065,
scale 1:253 440 or 1 inch to 4 miles.

Pye, E.G., Harris, F.R., Fenwick, K.G., and
Baillie, J.

1966: Tashota-Geraldton Sheet, Thunder Bay
and Cochrane Districts; Ontario
Department of Mines, Geological
Compilation Series, Map 2102, scale 1:253
440 or 1 inch to 4 miles.

Pyke, D.R., Ayres, L.D., and Innes, D.G.

1973: Timmins-Kirkland Lake, Cochrane,
Sudbury, and Timiskaming Districts;
Ontario Division of Mines, Geological
Compilation Series Map 2205, scale 1:253
440 or 1 inch to 4 miles.

Riddell, G.S.

1965: Statistical Review of the Mineral
Industry and Mining Operations for 1963;
Ontario Department of Mines, Annual
Report for 1963, Vol. 73.

1966: Statistical Review of the Mineral
Industry and Mining Operations for 1964;
Ontario Department of Mines, Annual
Report for 1964, Vol. 74.

1968: Statistical Review of the Mineral
Industry and Mining Operations for 1965;
Ontario Department of Mines, Annual
Report for 1965, Vol. 75.

Rio Algom Mines Limited

1972: Uranium from Elliot Lake; booklet
published by Corporate Public Relations
Department of Rio Algom Mines Limited.

Robertson, J.A.

1957: Township 144, District of Algoma;
Ontario Department of Mines, Map 2002,
scale 1:15,840.

1961: Geology of Townships 143 and 144,
District of Algoma; Ontario Department
of Mines, Geological Report 4, 65p.
Accompanied by Maps 2001 and 2002,
scale 1 inch to 1/4.

1962: Geology of Townships 137 and 138,
District of Algoma; Ontario Department
of Mines, Geological Report 10, 94p.
Accompanied by Maps 2003 and 2004,
scale 1 inch to 1/4 mile.

- 1963: Townships 155, 156, 161, and 162, District of Algoma; Ontario Department of Mines, Geological Report 13, 88p. Accompanied by Maps 2014, 2015, 2026, and 2027, scale 1 inch to 1/4 mile.
- 1964: Geology of Scarfe, Mack, Cobden, and Striker Townships; Ontario Department of Mines, Geological Report 20, 89p. Accompanied by Maps 2028 and 2032, scale 1 inch to 1/2 mile.
- 1967: Township 150, Algoma District; Ontario Department of Mines, Map 2114, scale 1:15 840 or 1 inch to 1/4 mile.
- 1968a: Uranium and Thorium Deposits of Northern Ontario, Ontario Department of Mines, Mineral Resources Circular 9, 106p.
- 1968b: Geology of Township 149 and Township 150, District of Algoma; Ontario Department of Mines, Geological Report 57, 162p. Accompanied by Maps 2113 and 2114, scale 1 inch to 1/4 mile.
- 1970a: Geology of the Spragge Area; Ontario Department of Mines, Geological Report 76, 109p. Accompanied by Maps 2185 and 2186, scale 1 inch to 1/2 mile.
- 1970b: Townships 1A and 1B; p. 49, in Summary of Field Work, 1970, by the Geological Branch, edited by E.G. Pye, Ontario Department of Mines and Northern Affairs, Miscellaneous Paper 43, 96p.
- 1975: Uranium and Thorium Deposits of Ontario; Ontario Division of Mines, Preliminary Maps P.969, P.970, P.971, P.972, Scale 1 inch to 16 miles. Compilation 1973.
- 1976a: The Blind River Uranium Deposits - The Ores and Their Settings; Ontario Division of Mines, Miscellaneous Paper 65, 45p.
- 1976b: Geology of the Massey Area, Districts of Algoma, Manitoulin, and Sudbury; Ontario Division of Mines, Geoscience Report 136, 130p. Accompanied by Maps 2308 and 2309, scale 1 inch to 1/2 mile.
- 1977a: Poulin and Sagard Townships, Algoma District; Ontario Geological Survey, Map 2346, Scale 1:31,680.
- 1977b: Nicholas and Raimbault Townships, Algoma District; Ontario Division of Mines, Geological Map 2347, scale 1:31 680.
- 1978: Uranium Deposits of Ontario; p.229-289 in "Short Course in Uranium Deposits, Their Mineralogy and Origin" M.M. Kimberley, editor, Mineralogical Association of Canada.
- 1981: The Uranium Deposits of Ontario - Their Distribution and Classification; Ontario Geological Survey, Miscellaneous Paper 86, 37p.
- 1982: Uranium and Thorium Deposits of Ontario, Northwestern Sheet, Kenora District (Patricia Portion); Ontario Geological Survey Preliminary Map P.2425, Mineral Deposits Series; Scale 1:1 013 760 or 1 inch to 16 miles. Compilation 1973-1974, 1980-1981.
- in press: Uranium and Thorium Deposits of Ontario, West Central Sheet, Districts of Kenora (Patricia Portion); Thunder Bay, Algoma, and Cochrane, Ontario Geological Survey, Preliminary Map P.2427, Mineral Deposits Series. Scale 1:1 013 760 or 1 inch to 16 miles. Compilation 1981.
- Robertson, J.A., Gordon, J.B., and Rybak, U.C.
1981: Uranium and Thorium Deposits of Ontario, Southern Sheet, Southern Ontario and Nipissing, Muskoka, Parry Sound, Sudbury, and Manitoulin Districts; Ontario Geological Survey Preliminary Map P. 2424, Mineral Deposit Series. Scale 1:1 013 760 or 1 inch to 16 miles. Compilation 1980 and 1981.
- Robertson, J.A. and Gould, K.L.
1981: Uranium and Thorium Deposits of Ontario, East Central Sheet, Districts of Thunder Bay, Cochrane, Algoma, Sudbury, Timiskaming, and Nipissing; Ontario Geological Survey Preliminary Map P. 2426, Mineral Deposits Series. Scale 1:1 013 760 or 1 inch to 16 miles. Compilation 1981.
- Robertson, J.A., Siemiatkowska, K.M., and Cape, D.F.
1972: Harrow Township and Adjacent Islands, Districts of Sudbury, Algoma, and Manitoulin; Ontario Division of Mines, Preliminary Map P.793, Geological Series, scale 1 inch to 1/4 mile. Geology 1972.

- Rodgers, D.P.
1964: Metionga Lake Area, District of Thunder Bay; Ontario Department of Mines, Geological Report 24, 53p.
- Runnalls, J.C.
1981: Ontario's Uranium Industry - Past, Present and Future; Ontario Ministry of Natural Resources, Mineral Background Paper No. 13.
- Ruzicka, V.
1977: Assessment of Selected Uranium Occurrences and Areas Favourable for Uranium Mineralization in Canada; p. 27-29, in Report of Activities Part A, Geological Survey of Canada, Paper 77-1A, 527p.
- Sage, R.P.
1975: Prairie Lake Complex, District of Thunder Bay; p. 64-66, in Summary of Fieldwork, 1975 by the Geological Branch, edited by V.G. Milne, D.F. Hewitt, K.D. Card, and J.A. Robertson; Ontario Division of Mines, Miscellaneous Paper 63, 158p.
- 1976: Carbonatite-Alkalic Complexes; p. 56-79, in Summary of Fieldwork, 1976, by the Geological Branch, edited by V.G. Milne, W.R. Cowan, K.D. Card, and J.A. Robertson, Ontario Division of Mines, Miscellaneous Paper 67, 183p.
- 1977: Carbonatite-Alkalic Complex, District of Algoma; p. 71-72, in Summary of Fieldwork 1977, by the Geological Branch, edited by V.G. Milne, O.L. White, R.B. Barlow, and J.A. Robertson, Ontario Geological Survey, Miscellaneous Paper 75, 208p.
- 1978: Radioactive Diatremes North of Lake Superior; p. 53-66, in Summary of Fieldwork, 1978, by the Ontario Geological Survey, edited by V.G. Milne, O.L. White, R.B. Barlow, and J.A. Robertson, Ontario Geological Survey, Miscellaneous Paper 82, 235p.
- Sage, R.P., and Breaks, F.W.
1982: Geology of the Cat Lake - Pickle Lake Area, Districts of Kenora (Patricia Portion) and Thunder Bay; Ontario Geological Survey, Report 207, 238p. Accompanied by Map 2218, scale 1:253 440.
- Sage, R.P., and Wright, W.
1979: Schryburt Lake Carbonatite Complex, District of Kenora (Patricia Portion); Ontario Geological Survey, Preliminary Map P.2236, Geology Series, scale 1:15 840. Geology 1976.
- Satterly, J.
1955: Radioactive Occurrence in the Vicinity of Hawk and Richard Lakes; Ontario Department of Mines, Geological Circular 1.
- Sergiades, A.O.
1968: Silver Cobalt Calcite Vein Deposits of Ontario; Ontario Department of Mines, Mineral Resources Circular 10, 489p.
- Sherran, H.W.
1952: Geology of Lac Aux Sables-Shakawa Lake Area, Ontario Department of Mines, Preliminary Report 1952-2, p.4.
- Shklanka, R.
1969: Copper, Nickel, Lead, and Zinc Deposits of Ontario; Ontario Department of Mines, Mineral Resources Circular 12, 394p.
- Siemiatkowska, K.M., Grunsky, Eric, and Berger, B.R.
1975: Endikai Lake Area (Eastern Half), Algoma District; Ontario Division of Mines, Preliminary Map P.1002, Geology Series, scale 1:15,840.
- Tanton, T.L.
1931: Fort William and Port Arthur and Thunder Bay Cape Map Areas, Thunder Bay District, Ontario; Geological Survey of Canada, Memoir 167, 222p.
- Thomson, J.E.
1953: Geology of Baldwin Township, Sudbury District; Ontario Department of Mines, Annual Report for 1952, Vol. 61, part 4.
- 1960a: Uranium and Thorium Deposits at the Base of the Huronian System in the District of Sudbury; Ontario Department of Mines, Geological Report 1, 40p.
- 1960b: MacLennan and Scadding Townships, Ontario Department of Mines, Geological Report 2, 34p. Accompanied by Map 2009. Scale 1 inch to 1/2 mile.

Thomson, R.

1966: Geology of Henwood Township,
District of Timiskaming; Ontario
Department of Mines, Miscellaneous
Paper 5, 48p.

Thurston, P.C., Siragusa, G.M., and Sage,
R.P.

1977: Geology of the Chapleau Area,
Districts of Algoma, Sudbury, and
Cochrane; Ontario Division of Mines,
Geoscience Report 157, 293p.
Accompanied by Maps 2351, 2352, and
2221.

Tihor, L.A. and Hunt, D.S.

1979: Report of the Northern Regional
Geologist and Timmins Resident
Geologist; p. 56-67 in Annual Report of
the Regional and Resident Geologist,
1978, edited by C.R. Kustra, Ontario
Geological Survey, Miscellaneous Paper
84, 135p.

Vos, M.A.

1975: Economic Geology of the Cretaceous
Deposits, Moose River Basin, Ontario,
General Appraisal; Ontario Division of
Mines; Open File Report 5157, 70p.

Walker, J.W.R.

1954: Jackfish-Middleton Area, Thunder Bay
District; Ontario Department of Mines,
Map 2107, scale 1:31 680 or 1 inch to 1/2
mile.

Western Miner

various: Western Miner; published monthly
by Gordon Black Publications.

Wood, J.

1975: Geology of the Rawhide Lake Area,
District of Algoma; Ontario Division of
Mines, Geoscience Report 129, 67p.
Accompanied by Maps 2305 and 2306,
scale 1 inch to 1/2 mile.

INDEX

Abconore occurrence	33	Broulan Reef occurrence	
Aberdeen Tp.	8	Sutherland Creek occurrence	112
Abeta occurrence	28	Bucke Tp.	135
Aerobus Lake occurrence	73	Buckles mine (Rio Algom Ltd.)	23
Aggressive occurrence	103-104	Buckles Tp.	16-20
Agnew Lake mine	106	Buffana prospect	50
Agnew Lake Mines Ltd.	6	Bull Lake	
Alanen-Maki occurrence	101-102	see: Denison occurrence	
Albanel Tp.	8,58	Byberg-Campbell-MacFarlane occurrence	
Alexander occurrence	112	see: Hawk Lake occurrence	
Alur occurrence	59		
Anabar occurrence	49	Callinan Flin-Flon occurrence	9
Anuwon occurrence	69	Calmor occurrence	71
Arco Triller No. 2	58	Cameron-Alcock occurrence	75
Asarco occurrence	91	Camray occurrence	45
Ascot occurrence	86-87	Canadian Johns-Manville	
Assembly occurrence	104	occurrences	
Assembly IA occurrence	104-105	Baldwin Tp.	99
Assembly 17A occurrence	108	Grigg Tp.	103
Astrabrun radioactivity occurrence	123	Marconi Tp.	119-120
Aubinadong River occurrence	116	Canagan Dike occurrence	42
Aubrey Falls occurrence	42	Candore prospect	9
		Caneonti prospect	42-43
Balboa occurrence	102-103	Cane Tp.	135
Baldwin Tp.	98-100,116	Cane Tp. occurrence	135
Bamaji Lake occurrence	73-74	Can-Fer occurrences	
Batchawana Uranium Mines		Marconi Tp.	120
occurrence	44-45	Kenora Dist.	75-76
Beange Tp.	9-10	Can-Met mine	
Bearhead Lake prospect	74-75	Production	6
Bee Lake occurrence	91	Denison Mines Ltd.	16-17
Belair Lake radioactivity		Canuc occurrence	59
occurrence	88	Cassidy Tp.	116
Benchmark occurrence	87	Celotti occurrence	129
Benner occurrence	12	Charron Lake	
Bidgood Kirkland mine	136	see: Sandy Stone occurrence	
Big Game occurrence	49-50	Chase Lake radioactivity	
Bigwood Tp.	116	occurrence	88
Black Lake radioactivity occurrence	64	Chemical Research occurrence	109
Blind River - Elliot Lake -		Chevrette occurrence	119
Agnew Lake Camp		Clary Tp.	116-117
Production data	6-7	Coal Creek occurrence	90
Blind River occurrence	61	Cobalt Cons. occurrence	41
Blue occurrence	122	Coffee Lake occurrence	51
Bluett Lake occurrence	83	Coleman Tp.	136
Bobcam occurrence	45	Collins & Chewett Tps.	117
Bolger Tp.	11-12,58	Conecho prospect	17
Bomby Tp.	130	Conmee Tp.	130
Borden Lake occurrence	118	Consolidated Denison Mines Ltd.	
Boston Tp.	135	see: Denison mine	
Bottle Bay occurrence	86	Copper	44,55,93
Bouck Tp.	12-16	Coral Rapids occurrence	72
Boymar occurrence	65	Corner Lake occurrence	20
Bracemar radioactivity		Coulee occurrence	
occurrence	63	Kimber Lake - West occurrence	81
Brewis-White occurrence	108-109	Cramette occurrence	132
Bridges occurrence	117	Crazy Lake occurrence	37
Bridges Tp.	81-82,90	Creelman Tp.	100-101,117
Broulan Reef Central occurrence	98	Crestland occurrence	70
Broulan Reef Lookout Tower		Crotch Lake occurrence	23-24
occurrence	98	Cullis Lake occurrence	20

Curtin Tp.	117	Gaiashk Tp.	20-22,59
Cyr occurrence	68	Gaitwin occurrence	55-56
Daley Tp.	128	Gallagher Tp.	118
Damascus occurrence	38	Gaudette Tp.	59
Davidson Lake occurrence	76	Gemico prospect	13
Day Tp.	20,58	Genex prospect	24
Deadhorse Creek (Diatreme) occurrence	132-133	Giles Tp.	22
Deanville occurrence	120-121	Gillies Limit Tp.	136
Delcan occurrence	122	Gimby occurrence	36-37
Demorest Lake occurrence	65	God's Lake Mines occurrence	65
Demorest Tp.	117	Gold	93
Denison occurrences		Golden Arrow occurrence	40
Turner Tp.		Goldie Tp.	130
Bull Lake	123	Greenwich Lake occurrence	125-126
Seagram Tp.	121-122	Greenwood Tp.	22-23
Timmermans Tp.	51	Grey Trout Lake prospect	43
Denison mine	6,12-13	Grigg Tp.	103
Denison Mines Ltd. Production		Gui-Por prospect	11
Can-Met mine	16-17	Gunterman Tp.	23-28,59-60
Stanrock mine	19-20	Habel Tp.	72
Denvic Lake occurrence	44	Hallam Tp.	103-104
Desbarats radioactivity occurrence	60	Halliday occurrence	76-77
Docker Tp.	82-83	Harrison-Hibbert occurrence	135
Dolan occurrence	38	Harrison occurrence	115
Dominion Gulf occurrence		Harrow Tp.	118
May Tp.	107-108	Harvard prospect	44
Shakespeare Tp.	113	Hathaway occurrence	32
Baldwin Tp.	116	Hawk Lake occurrence	
Dominion prospect	51-52	Byberg-Campbell-MacFarlane occurrence	84
Dorion amethyst mine	130	Headvue occurrence	77
Dorion Tp.	128,130	Headway occurrence	82
Drope Tp.	83	Hele Tp.	131
Drury Tp.	101-102	Hemlo occurrence	130
Duncan Tp.	58-59	Hennessy occurrence	38
Duncan Tp. occurrence	58	Heward Lake occurrence	97
East Bay occurrence	106-107	Hollinger radioactivity occurrence	120
Enterprise mine	132	Hornby Lake occurrence	88
Ermatinger Tp.	102-103	Huclif Porcupine radioactivity occurrence	61
Fairservice occurrence	81-82	Hudson Tp.	136
Falconbridge occurrence		Hunter Lake occurrence	110
Rix Tp.	66	Hurd occurrence	136
Fano radioactivity occurrences		Huston Lake radioactivity occurrence	88
Juillette Tp.	60	Hutton Tp.	104-105
Timmermans Tp.	69	Hyman Tp.	105-107,118-119
Fausten occurrence	47-48	Ingamar occurrence	118
Firesand River Carbonatite Complex	64	Innes Lake occurrence	128
Foot Lake occurrence	84	Inspiration occurrence	40-41
Fort Norman radioactivity occurrence	69	International occurrence	59
Fraleck Tp.	118	Iron	35-36
Franz occurrence	33-34	Iso Mines occurrence	
French River occurrence		Whiskey zone	20-21
South side of main channel	116	Jalore occurrence	61-62
Frye radioactivity occurrence	63	James occurrence	66
		Jarvis Tp.	60
		Jay-Dee occurrence	48

Jeanette prospect	52	McCombe J.G. occurrence	38-39
Jellicoe occurrence	99-100	McCombe Synd. occurrence	34
Espanola Bay	99	McCoy Tp.	131
Jenson occurrence	77	McDevitt occurrence	62
Johnson Lake radioactivity occurrence	88-89	McDonald Bay occurrence	113-114
Johnson Tp.	60	McDonough occurrence	35
Jollineau Tp.	28	MacFie occurrence	121
Jonsmith occurrence		McIntyre Porcupine occurrence	37
McKinnon occurrence	114	McIntyre Tp.	131
Joubin Tp.	28-32	McKellar Creek (Diatreme) occurrence	133
Juillette Tp.	60	McKinnon occurrence	
		see: Jonsmith occurrence	
Kakabeka Falls occurrence	130	Mack Tp.	64
Kamichisitit Tp.	32,61	MacLennan Tp.	107,119
Kamis occurrence	24	McMaster Tp.	91
Kassagimini Lake		McMurray Tp.	64
see: Sandy Stone occurrence		MacNicol Tp.	84-86,91
Kee No. 2 occurrence	32	McNish Tp.	120
Kenoratomic occurrence	85	McTavish Tp.	132
Kerr Addison prospect	102	Madawanson Lake occurrence	68
Kerr Lake Mine	136	Maeck Tp.	64
Kerr-McGee 150-151 occurrence	14	Mainville Lake occurrence	96
Kimber Lake (East) occurrence		Manwood radioactivity occurrence	67
Stephens, C.S.	82-83	Marcia Lake occurrence	
Kimber Lake - West occurrence		see: Noranda occurrence	
see: Coulee occurrence		Marconi Tp.	119-120
Kincaid Tp.	32	Martin occurrence	52-53
Kirkpatrick Lake radioactivity occurrence	63	Massey Bay occurrence	119
Kirkwood Tp.	33,61	Mattaini occurrence	56
Kirkwood Tp. radioactivity occurrence	61	Matinenda radioactivity occurrence	60
Kram-Glowaski occurrence	119	Maud Lake occurrence	59
		May Tp.	107-108
Labelle Tp.	33-34	Medicine Stone Lake occurrence	
Laberge Tp.	131	see: Pirson occurrence	
Labine-McCarthy occurrence	48	Meehan occurrence	
Labonte Tp.	34-35,61-62	see: Quibell occurrence	
Lacnor mine	25	Milliken mine	
Langlois Tp.	119	Rio Algom Ltd.	25-26
Langton Tp.	83-84	Milliken Lake Uranium Mines Ltd.	
Lapaska occurrence	62	see: Rio Algom (Milliken mine)	
Larson Tp.	35	Milliqua prospect	
Lebel occurrence	136-137	Pardee reef of Pardee zone	28-29
Lebel Tp.	136-137	Molson Lake occurrence	131
Lecaron Tp.	63	Moncrieff Tp.	120
Le Clerc occurrence		Monestime Tp.	64
see: Picton occurrence		Montgomery Tp.	64
Lefroy Tp.	63	Moon Lake occurrence	
Lehman Tp.	63	Timmermans Tp.	53
Leslie occurrence	100-101	Moon Lake Mines occurrence	
White River occurrence	8	Bolger Tp.	
Location X occurrence	35-36	Moon Lake zone	11
Long Lac occurrence	128	Moon Lake zone	11-12
Long Tp.	35-36	Morin Tp.	37,65
Loudon prospect	83	Mosher-Byles occurrence	46
Lunge occurrence	135	Mount Lake area	67
McAughey Tp.	36-37	Nairn Tp.	120-121
McBrien Tp.	90-91	Napray occurrence	35
		Nasco occurrence	29
		Neil occurrence	95

Nelson occurrence	130	Phillips occurrence	22-23
Nemegosenda Lake occurrence	117	Pic Bamooos occurrence	
New Campbell Island		Port Coldwell Complex	129
see: Richard Lake prospect		Pic Tp.	132
New Insko prospect		Piche Tp.	40-41
Prairie Lake Complex	126	Pick occurrence	122
New Mylamaque (Mymar)		Pickle Crow occurrence	93
occurrence	110-111	Picton occurrences	
New Quirke mine No.2		MacLennan Tp.	
Rio Algom Ltd.	15	Le Clerc occurrence	107
New Santiago occurrence		Timmermans Tp.	53-54
Sturgeon Narrows	126-127	Pilgrim Creek	
Nicholas Tp.	37	see: Noranda occurrence	
Niobium	33-34,79,126-127	Pineneedle Lake radioactivity	
Noranda occurrences		occurrence	89
Clary Tp.		Pine Road occurrence	89
Marcia Lake occurrence	116-117	Pirson occurrence	
Demorest Tp.		Medicine Stone Lake occurrence	78
Yorston Lake occurrence	117	Pistol Lake prospect	54
Hyman Tp.	105	Pitchore prospect	55
Turner Tp.		Pitt Tp.	71,72
Pilgrim Creek	123-124	Plum occurrence	100
Nordic Group West prospect		Port Arthur occurrence	131
Moon Lake zone	12	Port Coldwell Complex	
Nordic mine (Rio Algom Ltd.)	26-27	see: Pic Bamooos occurrence	
Nordic Resources prospect	111	Port Monroe occurrence	
North Creelman occurrence	117	Potvine	131
Northspan mine	6	Porter Tp.	108-111,121
Northspan Uranium Mines Ltd.		Potvine	
see: Northspan mine		see: Port Monroe occurrence	
Oak Lake occurrence	89	Poulin Tp.	41
Octopus Lake radioactivity		Prairie Lake Complex	
occurrence	89	see: New Insko prospect	
O'Neill T#32		Preston mine (Stanleigh)	6
Ottawa Assoc. occurrence	34	Preston occurrence	83
Paddington Lake occurrence	91	Pronto mine (Rio Algom Ltd.)	36
Pancer occurrence	77-78	Pronto Uranium Mines Ltd.	
Panel mine (Rio Algom Ltd.)	17-18	see: Rio Algom (Pronto mine)	
Pardee prospect		Proscop prospect	93
Pardee zone	29-30	Provencher occurrence	72
Pardee zone	28-31	Punkari occurrence	131
Pardo Tp.	93	Quibell occurrence	
Parkin Tp.	108	Meehan occurrence	92
Parth occurrence	90	Quirke Group West	
Paterson Lake occurrence	89	Rio Algom	10
Patrick occurrence	39	Quirke mine No. 1 (Rio Algom Ltd.)	14-15
Pearl Lake occurrence	132	Raimbault Tp.	65-66
Pecors Lake East prospect		Ranger occurrence	41-42
Whiskey zone	21	Ranrouyn occurrence	68
Pecors Lake West prospect		Ranwick prospect	39
Pardee zone	30	Rare earths	14
Pecors zone	31-32	Rawhide occurrence	56-57
Peerless Uranium Mining Corp. Ltd.		Reilly Tp.	41-42
Exploration area	58	Reserve Lake occurrence	60
Peever Tp.	38-40	Reynar Lake occurrence	90
Pennbec occurrence	121	Richard Lake prospect	
Pennwood occurrence	67	New Campbell Island	85-86
Perch Lake occurrence	78	Richore occurrence	118-119
Petursson Lake occurrence	87		

Rio Algom Ltd.	6	Squaw Lake occurrence	129
Buckles mine	23	Stancan occurrence	30-31
Lacnor mine	7	Stanleigh mine (Rio Algom Ltd.)	27-28
Milliken mine	7,25-26	Stanleigh Uranium Mining Corp. Ltd.	
New Quirke mine No.2	15	See: Preston mine	
Nordic mine	7,26-27	Stanrock mine	7
Panel mine	7,17-18	Denison Mines Ltd.	19-20
Pronto mine	7,36	Stencill occurrence	118
Quirke Group West occurrence	10	Stephens, C.S.	
Quirke mine	7	see: Kimber Lake (East) occurrence	
Quirke mine No.1	14-15	Stetham Tp.	114,123
Spanish American mine	16	Stobie Tp.	114-115
Span-North occurrence	10	Stobie Tp. occurrence	114-115
Stanleigh mine	27-28	Strain Tp.	68
Rio Tinto prospect	8	Strathcona Tp.	95
Rix Tp.	42,66	Striker Tp.	68
Robb-Murmac prospect	46-47	Sturgeon Narrows	
Roberts Lake occurrence	111	see: New Santiago occurrence	
Roberts Tp.	111	Sugar Lake occurrence	124
Roche Long Lac prospect	18	Surluga occurrence	64
Roche occurrence	66	Sutherland Creek occurrence	
Rollins Tp.	66	see: Broulan Reef occurrence	
Roman Island occurrence	19	Sutherland occurrence	65
Root Tp.	42	Sydney Lake radioactivity occurrence	90
Russian Lake radioactivity occurrence	64		
		Tarbutt Mines occurrence	68
St. Mary's occurrence		Tarbutt Tp.	68
Pardee zone	31	Temple Tp.	86
Sagard Tp.	42-44,67	Tessier-Williamson occurrence	131
Salter Tp.	67	Theano Point prospect	46
Sandy Stone occurrence		Thessalon Tp.	68-69
Charron Lake	127	Thorncrest radioactivity occurrence	60
Kassagimini Lake	127-128	Timmermans Tp.	49-55,69
Satellite occurrence	122	Tooker radioactivity occurrence	62
Sauve-Inch occurrence	137	Tourist Lake prospect	79-80
Saville occurrence	120	Towle-Humpage occurrence	124
Schlesinger occurrence	123	Treelined Lake occurrence	80
Schryburt Lake occurrence	79	Trottier Tp.	97
Seabrook Lake Carbonatite Complex	64	Tudhope Tp.	137
Seagram Tp.	121-122	Tupper Tp.	55
Shackleton Tp.	72	Turner Tp.	115,123-124
Shakespeare Tp.	112-114,122-123	Tustin Tp.	86-87,91
Shedden Tp.	44		
Shunck occurrence	67	United Cobalt occurrence	116
Silvermaque Mine	136	Univex occurrence	129
Silvermaque occurrence	27		
Slater Tp.	44-47,67	Vankoughnet occurrence	55
Smilsky Tp.	47-48,68	Vankoughnet Tp.	55
Smith occurrence	132	Van Lake occurrence	39-40
Snook Lake occurrence	79	Vermilion Lake occurrence	80
Soo-Tomic occurrences		Viceroy occurrence	91
Giles Tp.	22	Victoria Tp.	124
Slater Tp.	47	Viel Tp.	55-57,70
Sowerby occurrence	58	Vite occurrence	
Spanish American mine (Rio Algom Ltd.)	16	Pecors zone	31-32
Span-North occurrence (Rio Algom Ltd.)	10	Vogt Tp.	93
Spencer occurrence	136		
Spragge Creek occurrence	49	Wabigoon Tp.	87,92
Spragge Tp.	49	Wadge occurrence	121
		Walsh Tp.	132-133

Webb occurrence	63
Weston radioactivity occurrence	70
West Thessalon occurrence	68
Whiskey prospect	
Whiskey zone	21-22
Whiskey zone	20-22
Whitman Tp.	70
Wilson occurrence	90
Wolf Island occurrence	80-81
Yorston Lake occurrence	
see: Noranda occurrence	
Yost occurrence	135
Yttrium	12-13, 19-20, 26-27, 106
Yzerdraat occurrence	129-130
Zenmac occurrence	
Timmermans Tp.	54
Zenmac radioactivity occurrence	
Raimbault Tp.	66
Zirconium	115

7476
450 P.R., 83/11
ISSN 0706-4551
ISBN 0-7743-8439-5