

THESE TERMS GOVERN YOUR USE OF THIS DOCUMENT

Your use of this Ontario Geological Survey document (the “Content”) is governed by the terms set out on this page (“Terms of Use”). By downloading this Content, you (the “User”) have accepted, and have agreed to be bound by, the Terms of Use.

Content: This Content is offered by the Province of Ontario’s *Ministry of Northern Development and Mines* (MNDM) as a public service, on an “as-is” basis. Recommendations and statements of opinion expressed in the Content are those of the author or authors and are not to be construed as statement of government policy. You are solely responsible for your use of the Content. You should not rely on the Content for legal advice nor as authoritative in your particular circumstances. Users should verify the accuracy and applicability of any Content before acting on it. MNDM does not guarantee, or make any warranty express or implied, that the Content is current, accurate, complete or reliable. MNDM is not responsible for any damage however caused, which results, directly or indirectly, from your use of the Content. MNDM assumes no legal liability or responsibility for the Content whatsoever.

Links to Other Web Sites: This Content may contain links, to Web sites that are not operated by MNDM. Linked Web sites may not be available in French. MNDM neither endorses nor assumes any responsibility for the safety, accuracy or availability of linked Web sites or the information contained on them. The linked Web sites, their operation and content are the responsibility of the person or entity for which they were created or maintained (the “Owner”). Both your use of a linked Web site, and your right to use or reproduce information or materials from a linked Web site, are subject to the terms of use governing that particular Web site. Any comments or inquiries regarding a linked Web site must be directed to its Owner.

Copyright: Canadian and international intellectual property laws protect the Content. Unless otherwise indicated, copyright is held by the Queen’s Printer for Ontario.

It is recommended that reference to the Content be made in the following form:

Gao, C. and Yeung, K.H. 2018. Surficial geology of the Kenogami River area southwest, northern Ontario; Ontario Geological Survey, Preliminary Map P.3752, scale 1:100 000

Use and Reproduction of Content: The Content may be used and reproduced only in accordance with applicable intellectual property laws. *Non-commercial* use of unsubstantial excerpts of the Content is permitted provided that appropriate credit is given and Crown copyright is acknowledged. Any substantial reproduction of the Content or any *commercial* use of all or part of the Content is prohibited without the prior written permission of MNDM. Substantial reproduction includes the reproduction of any illustration or figure, such as, but not limited to graphs, charts and maps. Commercial use includes commercial distribution of the Content, the reproduction of multiple copies of the Content for any purpose whether or not commercial, use of the Content in commercial publications, and the creation of value-added products using the Content.

Contact:

FOR FURTHER INFORMATION ON	PLEASE CONTACT:	BY TELEPHONE:	BY E-MAIL:
The Reproduction of the EIP or Content	MNDM Publication Services	Local: (705) 670-5691 Toll-Free: 1-888-415-9845, ext. 5691 (inside Canada, United States)	Pubsales.ndm@ontario.ca
The Purchase of MNDM Publications	MNDM Publication Sales	Local: (705) 670-5691 Toll-Free: 1-888-415-9845, ext. 5691 (inside Canada, United States)	Pubsales.ndm@ontario.ca
Crown Copyright	Queen’s Printer	Local: (416) 326-2678 Toll-Free: 1-800-668-9938 (inside Canada, United States)	Copyright@ontario.ca

Marginal Notes

In 2008, the Ontario government announced plans to permanently protect half of the Far North region of Ontario and launched a planning process to support the goal (Far North Information and Knowledge Management Program). During the initial stages of planning, the need for primary landscape data became apparent. A terrain mapping project to remotely predict surficial materials was initiated by the Ontario Geological Survey in response to this information need.

SPOT imagery (4 colour bands and the panchromatic band), a digital elevation model and its derivatives and the Ontario Hydro Network vector drainage shape files (Ontario Ministry of Natural Resources and Forestry 2015) are the primary data sources for this remote predictive mapping exercise. A multiresolution segmentation algorithm, using different image layer weights, scale parameters and homogeneity criterion, within an object-based image analysis software is used to achieve meaningful objects representing various surficial material types. Objects are then classified based on digital signature, internal variability of signature and proximity to certain vector layers and certain adjacent material types.


Limited helicopter-supported field work combined with the examination of archival information (Riley and Boissonneau unpublished field notes and photographs; Geological Survey of Canada, Operation Wink; unpublished field notes; Skinner 1973; Sanford, Norris and Bostock 1968; Sanford and Norris 1975; and Thorleifson, Wyatt and Warman 1993, and their unpublished field notes) provided the ground control on the classification of objects. In addition, information from the various other Far North Information and Knowledge Management Program projects, such as base data and land cover information (Ontario Ministry of Natural Resources and Forestry 2014), has been used in the interpretation and classification of the surficial materials.

The maps show only the surface material distribution. For better definition of wetland types, please see Ontario Ministry of Natural Resources and Forestry (2014). Older deposits that occur along many of the deeply incised river valleys are only depicted where their scale is suitable for cartographic presentation. The reader is encouraged to review publications by McDonald (1969), Skinner (1973), Nielsen et al. (1986), Dredge, Morgan and Nielsen (1990), Thorleifson, Wyatt and Warman (1993), Allard et al. (2012), Dubé-Loubert et al. (2013), Gao et al. (2012), Gao and Crabtree (2016) and Dalton et al. (2016, 2017a, 2017b) for additional information and details on the older sediments exposed along rivers within the Hudson Bay Lowland.

This project is funded by the Ministry of Northern Development and Mines. Interaction with the First Nation community members greatly enhanced the map products of the Far North Information and Knowledge Management Program Terrain Mapping Project.

REFERENCES

- Allard, G., Roy, M., Chabot, B., Richard, P.J.M., Larouche, A.C., Veilleux, J.J., and Parent, M., 2012. Constraining the age of the last interglacial-glacial transition in the Hudson Bay Lowlands (Canada) using U-Th dating of buried wood; *Quaternary Geochronology*, v.7, p.37-47.
- Dalton, A.S., Finkelstein, S.A., Barnett, P.J. and Forman, S.L., 2016. Constraining the Late Pleistocene history of the Laurentide ice Sheet by dating the Missinabi Formation, Hudson Bay Lowlands, Canada; *Quaternary Science Reviews*, v.146, p.288-299.
- Dalton, A.S., Valiranta, M., Barnett, P.J. and Finkelstein, S.A., 2017a. Pollen and macrofossil-inferred palaeoclimate at the Ridge Site, Hudson Bay Lowlands, Canada: evidence for a dry climate and significant recession of the Laurentide ice Sheet during Marine Isotope Stage 3; *Boreas*, v.46, p.388-401.
- Dalton, A.S., Finkelstein, S.A., Barnett, P.J., Valiranta, M. and Forman, S.L., 2017b. Late Pleistocene chronology, palaeoecology and stratigraphy at a suite of sites along the Albany River, Hudson Bay Lowlands, Canada; *Palaeogeography, Palaeoclimatology, Palaeoecology*, v.492, p.50-63. doi:10.1016/j.palaeo.2017.12.011
- Dredge, L.A., Morgan, A. and Nielsen, E., 1990. Sangamon and pre-Sangamon interglaciations in the Hudson Bay Lowlands of Manitoba; *Géographie physique et Quaternaire*, v.44, no.3, p.319-336.
- Dubé-Loubert, H., Roy, M., Allard, G., Lamothe, M. and Veilleux, J.J., 2013. Glacial and nonglacial events in the eastern James Bay Lowlands, Canada; *Canadian Journal of Earth Sciences*, v.50, p.379-396.
- Gao, C. and Crabtree, D.C., 2016. Results of regional till and modern alluvium sampling in the McFaulds Lake ("Ring of Fire") area, northern Ontario; *Ontario Geological Survey, Open File Report 6369*, 16p.
- Gao, C., McAndrews, J.H., Wang, X., Mercieca, J., Turton, C.L., Wood, B.D., Pei, J. and Kodors, C., 2012. Glaciation of North America in the James Bay Lowland, Canada, 3.5 Ma; *Geology*, v.40, no.11, p.975-978.
- Geological Survey of Canada. Operation Wink; unpublished field notes, Geological Survey of Canada, Ottawa, Canada.
- McDonald, B.C., 1969. Glacial and interglacial stratigraphy, Hudson Bay Lowlands; in *Earth Science Symposium on Hudson Bay*, Geological Survey of Canada, Paper 68-53, p.78-99.
- Nielsen, E., Morgan, A.V., Morgan, A., Mott, R.J., Rutter, N.W. and Causee, C., 1986. Stratigraphy, paleoecology, and glacial history of the Gillam area, Manitoba; *Canadian Journal of Earth Sciences*, v.23, p.1641-1661.
- Ontario Ministry of Natural Resources and Forestry 2014. Far North Land Cover; Ministry of Natural Resources and Forestry, Land Information Ontario, Peterborough, Ontario.
- Ontario Ministry of Natural Resources and Forestry 2015. Ontario Hydro Network; Ministry of Natural Resources and Forestry, Land Information Ontario, Peterborough, Ontario.
- Riley, J. and Boissonneau, A. Unpublished field notes and photographs; Ontario Ministry of Natural Resources, Peterborough, Ontario.
- Sanford, B.V. and Norris, A.W., 1975. Devonian stratigraphy of the Hudson Bay Platform; *Geological Survey of Canada, Memoir 379*, 124p.
- Sanford, B.V., Norris, A.W. and Bostock, H.H., 1968. Geology of the Hudson Bay Lowlands (Operation Wink); *Geological Survey of Canada, Paper 67-60*, p.1-45.
- Skinner, R.G., 1973. Quaternary stratigraphy of the Moose River Basin, Ontario; *Geological Survey of Canada, Bulletin 225*, 77p.
- Thorleifson, L.H., Wyatt, P.H. and Warman, T.A., 1993. Quaternary stratigraphy of the Severn and Winkisk drainage basins, northern Ontario; *Geological Survey of Canada, Bulletin 442*, 69p.



LEGEND*

PHANEROZOIC

CENOZOIC

QUATERNARY

- 21 **Anthropogenic Deposits:** Mine tailings, waste rock piles and other man-made features.
- 20 **Marine Mud-Flat Deposits:** Sand, silt and clay; deposited in mud flats and offshore bars within the tidal zone.
- 19 **Marine Salt-Marsh Deposits:** Silt and clay with organic matter, minor sand; deposited in salt marshes along present-day coast.

- 18 **Organic Deposits:** Peat, muck and/or marl; deposited in fen wetlands.[†]
- 17 **Organic Deposits:** Peat, muck and/or marl; deposited in bog wetlands.[†]
- 16 **Fluvial Deposits (recent):** Stratified sand and gravel; may include silt, minor clay. Deposited in bars and channels of present-day rivers and creeks.
- 15 **Fluvial Deposits (abandoned):** Stratified sand and gravel; may include silt, minor clay. Deposited in bars and channels along abandoned terraces of rivers and creeks.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

- 14 **Marine Deltaic Deposits:** Stratified sand and gravel; silt, minor clay. Deposited at the mouths of rivers and creeks entering a postglacial sea.
- 13 **Marine Beach and Nearshore Deposits:** Stratified sand and gravel, minor silt. Deposited primarily in the shore zone and nearshore zone of a postglacial sea.
- 12 **Marine Basin Deposits:** Massive to stratified silt and clay; minor sand. Deposited in basinal areas of a postglacial sea.
- 11 **Glaciolacustrine Deltaic Deposits:** Stratified sand and gravel; may include minor silt and clay. Deposited at the mouths of rivers and creeks entering a proglacial lake.
- 10 **Glaciolacustrine Beach and Nearshore Deposits:** Stratified sand and gravel; minor silt. Deposited primarily in the shore zone and nearshore zone of a proglacial lake.
- 9 **Glaciolacustrine Basin Deposits:** Stratified silt and clay; minor sand. Deposited in basinal areas of a proglacial lake.
- 8 **Glaciofluvial Outwash Deposits:** Stratified sand and gravel; including minor silt, clay. Deposited in bars and channels in rivers flowing from a glacier.
- 7 **Glaciofluvial Ice-Contact Deposits:** Stratified sand and gravel; including minor silt, clay, till and flow till. Deposited in eskers, kames, deltas and subaqueous fans and in end and recessional moraines along a glacier margin.

- 6 **Till:** Massive to bedded diamicton; clayey silt to clay matrix, clast content low to moderate, moderate to high matrix carbonate content, may contain discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, includes flow tills. Deposited at the base or along the margins of a glacier.
- 5 **Till:** Massive to bedded diamicton; sandy silt to silt matrix (clay texture locally), clast content moderate to high, low to high matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 4 **Till:** Massive to bedded diamicton; silty sand to sand matrix, clast content moderate to high, low to moderate matrix carbonate content, rare discontinuous layers or lenses of stratified gravel, sand, silt and/or clay, or flow tills. Deposited at the base or along the margins of a glacier.
- 3 **Thin Sediment over Bedrock:** Thin, near-continuous cover of Quaternary sediments overlying undifferentiated bedrock. Quaternary sediments are commonly less than 1 m thick; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 2 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.
- 1 **Bedrock:** Undifferentiated bedrock with a thin, discontinuous cover of Quaternary sediments. Quaternary sediments rarely exceed 0.5 m in thickness; however, areas of outcrop indicated on the map may contain small areas where the cover exceeds 1 m in thickness, but are too small to delineate at this map scale.

SOURCES OF INFORMATION

The digital base map is derived from the Land Information Ontario Data Warehouse, Land Information Ontario, Ministry of Natural Resources and Forestry, scale 1:50 000, with modifications by staff of the Ministry of Northern Development and Mines. The digital base data is current to January 2017. The map co-ordinates are in UTM zone 16, North American Datum 1983 (NAD83).

Magnetic declination at the centre of the map area was approximately 7°40'W in 2018. Mean annual change is 0.7°E.

Barnett, P.J., Webb, J.L. and Hill, J.L., 2009. Flow indicator map of the Far North of Ontario; *Ontario Geological Survey, Preliminary Map P-3610*, scale 1:1 000 000.

Norris, A.W., Sanford, B.V. and Bostock, H.H., 1967. Geology, Hudson Bay Lowlands, Manitoba, Ontario, Quebec and District of Keewatin, compiled; *Geological Survey of Canada, Preliminary Map 17-1967*. Accompanies Geological Survey of Canada, Paper 67-60.

Ontario Centre for Remote Sensing. Surficial geology, Fort Albany area, Ontario; unpublished map, Ontario Ministry of Natural Resources, OCRS-43A, scale 1:250 000.

Ontario Centre for Remote Sensing. Surficial geology, Kapisikau River area, Ontario; unpublished map, Ontario Ministry of Natural Resources, OCRS-43B, scale 1:250 000.

Ontario Ministry of Natural Resources 2012. Ontario radar digital surface model; Ministry of Natural Resources, Land Information Ontario, Peterborough, Ontario.

Ontario Ministry of Natural Resources and Forestry 2014. Far North land cover; Ministry of Natural Resources and Forestry, Land Information Ontario, Peterborough, Ontario.

Sanford, B.V. and Norris, A.W., 1975. Devonian stratigraphy of the Hudson Bay Platform; *Geological Survey of Canada, Memoir 379*, 124p.



Ontario Geological Survey

MAP P.3752

SURFICIAL GEOLOGY

KENOGAMI RIVER AREA SOUTHWEST, NORTHERN ONTARIO

Scale 1:100 000

2000 m 0 2 4 km

NTS References: 42 K/3, 4, 5, 6

© Queen's Printer for Ontario, 2018.

This map is published with the permission of the Director, Ontario Geological Survey.

